

## WORKSHEET 5: APPLICATIONS OF DERIVATIVES

Name: \_\_\_\_\_ Section No: \_\_\_\_\_

### Derivatives and Graphs

- (1) For the following functions, use first and second derivatives to determine (i) all points  $x$  where  $f(x)$  is a local maximum or minimum value, (ii) all open intervals where  $f(x)$  is increasing, decreasing, concave up, and concave down. Give all answers **exactly**, not as numerical approximations.
- (a)  $f(x) = x^5 - 2x^3$
- (b)  $f(x) = x + \sin x$  for  $-2\pi \leq x \leq 2\pi$
- (c)  $f(x) = e^{-x} - e^{-3x}$
- (2) Let  $f(x) = x^{100} + (100 - x)^{100}$ . Determine where  $f(x)$  is increasing and decreasing for  $0 \leq x \leq 100$  and use this information to decide which of  $33^{100} + 67^{100}$  or  $41^{100} + 59^{100}$  is larger.
- (3) Sketch a possible graph of a function  $y = f(x)$  over the interval  $[-2, 2]$  when its derivatives have the following properties:
- $f'(x) = 0$  for  $x = 1$  and  $x = -1$ ,  $f'(x) > 0$  for  $x$  in  $(-2, -1)$ , and  $f'(x) < 0$  for  $x$  in  $(-1, 1)$  and  $(1, 2)$ .
  - $f''(x) = 0$  for  $x = 0$  and  $x = 1$ ,  $f''(x) > 0$  for  $x$  in  $(0, 1)$ , and  $f''(x) < 0$  for  $x$  in  $(-2, 0)$  and  $(1, 2)$ .

### L'Hospital's Rule

- (4) Describe each limit's indeterminate form and then compute the limit. If l'Hospital's Rule is needed more than once, try to simplify the expression before applying it.
- (a)  $\lim_{x \rightarrow 0} \frac{(x+1)^{11} - 11x - 1}{x^2}$
- (b)  $\lim_{x \rightarrow 0} \frac{\sin(3x)}{e^{9x} - e^{2x}}$
- (c)  $\lim_{x \rightarrow \infty} \frac{\ln(2013x^2 + 1)}{\ln x}$
- (d)  $\lim_{x \rightarrow 0^+} \frac{\ln(\sin(2x))}{\ln(\sin(3x))}$
- (e)  $\lim_{x \rightarrow 0} \frac{\ln(\cos(2x))}{\ln(\cos(3x))}$
- (5) Try to use l'Hospital's Rule on  $\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}}$  and explain what goes wrong. Then evaluate this limit using earlier methods from the course.

### Optimization

- (6) Find the points on  $y = x^2$  that are closest to  $(0, 3)$  by the following steps.
- (a) For each point  $(x, x^2)$  on the parabola, find a formula for its distance to  $(0, 3)$ . Call this  $f(x)$ . (See Figure 1.)
- (b) Let  $g(x) = f(x)^2$ , which is the *squared distance* between  $(x, x^2)$  and  $(0, 3)$ . Show  $g'(x)$  is 0, positive, or negative exactly when  $f'(x)$  is 0,

positive, or negative. So finding out where  $f(x)$  is minimal is the same as finding out where  $g(x)$  is minimal.

- (c) Find all  $x$  where  $g'(x) = 0$  and determine among such  $x$  where  $g(x)$  (and thus  $f(x) = \sqrt{g(x)}$ ) has a local minimum value. Explain why  $g(x)$  has an absolute minimum value at such  $x$ . The points  $(x, x^2)$  for such  $x$  are the closest points to  $(0, 3)$  on  $y = x^2$ .
- (d) Where does  $g'(x) = 0$  and  $g(x)$  *not* have a local minimum value? Is there something special about the distance of such points  $(x, x^2)$  to the point  $(0, 3)$ ?




FIGURE 1. Distance to  $(0, 3)$  on  $y = x^2$ .

- (7) Three line segments of length 1 are joined together at endpoints to form a base and the legs of an isosceles trapezoid, as in Figure 2. Let  $\theta$  be the common angle measurement between the legs and the line passing through the base of length 1.




FIGURE 2. An isosceles trapezoid with base and legs of length 1.

- (a) Compute the area  $A(\theta)$  of the trapezoid. The general area formula for a trapezoid is  $\frac{1}{2}h(b_1 + b_2)$ , where  $h$  is the height and  $b_1$  and  $b_2$  are the lengths of the bases. (Hint: Break up the trapezoid into a rectangle with two right triangles at both ends. Use trigonometry to compute the height and the length of the longer base in terms of  $\theta$ .)
- (b) Find all solutions to  $A'(\theta) = 0$  with  $0 < \theta < \pi/2$ . (The answer is *not*  $\pi/4 = 45^\circ$ .)
- (c) Verify that the area  $A(\theta)$  is a maximum at the angle found in part b and compute this maximum area.