

Name: _____

Discussion Section: _____

Solutions should show all of your work, not just a single final answer.

3.3: Derivatives of Trigonometric Functions

1. Compute the derivative of each function below using differentiation rules.

(a) $f(x) = x^3 \cos x$

(b) $f(x) = \frac{1 + \sin x}{1 + \cos x}$

(c) $f(x) = e^x \tan x$

(d) $f(x) = \frac{\sec x}{\sqrt{x}}$ (Compute (d) in **two ways**, using (i) the quotient rule and (ii) the product rule.)

2. Find the equation of the tangent line to the curve $y = \sin x \cos x$ at $x = \frac{\pi}{4}$. (Your coefficients must be exact, not approximations.)

3. Find the higher derivative $\frac{d^{2017}}{dx^{2017}}(2 \cos x)$ by finding the first eight derivatives and observing the pattern that occurs.

4. Determine the following limits by making a change of variables to allow you to use the relation $\lim_{t \rightarrow 0} \frac{\sin t}{t} = 1$.

(a) $\lim_{x \rightarrow 0} \frac{\sin 4x}{x}$

(b) $\lim_{x \rightarrow 0} \frac{\sin 7x}{5x}$

3.4: The Chain Rule

5. Compute the derivative with respect to x of each function below using differentiation rules.

(a) $f(x) = (x^3 - x + 1)^{10}$

(b) $f(x) = \sqrt{x^3 + 4x}$

(c) $f(x) = e^{ax} \cos(bx)$ for constants a and b

(d) $f(x) = \left(\frac{e^x}{3-x}\right)^8$

(e) $f(x) = \sin^2(x) - \sin(x^2)$

6. Differentiate the functions below **with respect to** t , where $r = r(t)$ is a function of t .

(a) $(r^2 + 1)^4$

(b) $\sin(2r) - 2 \sin r$

(c) e^{r^2+ar+b} for constants a and b .

7. If $f'(0) = 5$ and $F(x) = f(3x)$, what is $F'(0)$?

8. T/F (with justification) If $f(x)$ is differentiable, then $\frac{d}{dx}(f(\sqrt{x})) = \frac{f'(x)}{2\sqrt{x}}$.

3.5: Implicit Differentiation

9. Find $\frac{dy}{dx}$ using implicit differentiation. Your final answer may involve both x and y .

(a) $x^2y - axy^2 = x + y$ where a is a constant.

(b) $\sin(x + y) = x + \cos(3y)$


(c) $e^{xy} = x^2 + y^2$

(d) $x = \arctan(y^2)$

10. Use implicit differentiation to find an equation of the tangent line to the curve

$$x^2 + y^2 = (2x^2 + 2y^2 - x)^2$$

at the point $(0, 1/2)$. **Note.** The graph of this equation is known as a cardioid, shown below. It's not the graph of a function, and this is where implicit differentiation can be helpful to us.



11. On the ellipse $x^2 + 9y^2 = 9$, find $\frac{d^2y}{dx^2}$ using implicit differentiation. Your final answer may involve both x and y .

