


# Homomesies Lurking in the Twelfefold Way

Tom Roby (UConn)

*Describing joint research with  
Michael Joseph & Michael LaCroix*

Special Session on Algebraic and Enumerative Com  
AMS Eastern Sectional Meeting  
Bowdoin College Brunswick, ME USA

25 September 2016 (Sunday)



Slides for this talk are available online (or will be soon) at

<http://www.math.uconn.edu/~troby/research.html>

# Homomesies Lurking in the Twelfold Way

Tom Roby (UConn)

*Describing joint research with  
Michael Joseph & Michael LaCroix*

Special Session on Algebraic and Enumerative Combinatorics  
AMS Eastern Sectional Meeting  
Bowdoin College Brunswick, ME USA

25 September 2016 (Sunday)

Slides for this talk are available online (or will be soon) at

<http://www.math.uconn.edu/~troby/research.html>

**Abstract:** Given a group acting on a finite set of combinatorial objects, one can often find natural statistics on these objects which are *homomesic*, i.e., over each orbit of the action, the average value of the statistic is the same. Since the notion was codified a few years ago, homomesic statistics have been uncovered in a wide variety of situations within dynamical algebraic combinatorics. We discuss several examples lurking in Rota's *Twelvefold Way* related to actions on injections, surjections (joint work with Michael Joseph), and bijections/permutations (joint work with Michael LaCroix) of finite sets.

## Acknowledgments

This seminar talk discusses joint work with Michael Joseph and Michael La Croix. Thanks to James Propp for suggesting the study of whirling and of Foatic actions, as well as earlier collaborations on the homomesy phenomenon.

Please feel free to interrupt with questions or comments.

- Actions, orbits, and homomesy;
- The Twelfefold Way;
- Foatic actions on  $\mathfrak{S}_n$ .
- Whirling injections and surjections;

Among the most basic objects counted by the Twelfold Way are subsets, counted by binomial coefficients.

Set  $[n] := \{1, 2, \dots, n\}$ .

$\binom{[n]}{k} = \{k\text{-element subsets of } [n]\}$ .

For example,  $\binom{[7]}{3}$  consists of 35 subsets (dropping braces & commas): 123, 124, 125, 126, 127, 134, 135, 136, 137, 145, 146, 147, 156, 157, 167, 234,...

The map that adds 1 to each element mod  $n$  acts on  $\binom{[n]}{k}$ .

156  $\mapsto$  267  $\mapsto$  137  $\mapsto$  124  $\mapsto$  235  $\mapsto$  346  $\mapsto$  457  $\mapsto$  156

It's easy to see that the cardinality of each orbit of this action is a divisor of  $n$ .

We find another unexpected structure in the orbits if we consider  $S = \binom{[n]}{k}$ , as length  $n$  binary strings with  $k$  1's as we cyclically shift them.  $\tau := C_R : S \rightarrow S$  by  $b = b_1 b_2 \cdots b_n \mapsto b_n b_1 b_2 \cdots b_{n-1}$ , and count  $f(b) = \#\text{inversions}(b) = \#\{i < j : b_i > b_j\}$ .

We find another unexpected structure in the orbits if we consider  $S = \binom{[n]}{k}$ , as length  $n$  binary strings with  $k$  1's as we cyclically shift them.  $\tau := C_R : S \rightarrow S$  by  $b = b_1 b_2 \cdots b_n \mapsto b_n b_1 b_2 \cdots b_{n-1}$ , and count  $f(b) = \#\text{inversions}(b) = \#\{i < j : b_i > b_j\}$ .  
EG:  $n = 4, k = 2$  gives us two orbits:

0011 0101



We find another unexpected structure in the orbits if we consider  $S = \binom{[n]}{k}$ , as length  $n$  binary strings with  $k$  1's as we cyclically shift them.  $\tau := C_R : S \rightarrow S$  by  $b = b_1 b_2 \cdots b_n \mapsto b_n b_1 b_2 \cdots b_{n-1}$ , and count  $f(b) = \# \text{inversions}(b) = \#\{i < j : b_i > b_j\}$ .  
EG:  $n = 4, k = 2$  gives us two orbits:

0011	0101
<hr/>	
1001	1010
1100	0101
0110	
0011	

We find another unexpected structure in the orbits if we consider  $S = \binom{[n]}{k}$ , as length  $n$  binary strings with  $k$  1's as we cyclically shift them.  $\tau := C_R : S \rightarrow S$  by  $b = b_1 b_2 \cdots b_n \mapsto b_n b_1 b_2 \cdots b_{n-1}$ , and count  $f(b) = \# \text{inversions}(b) = \#\{i < j : b_i > b_j\}$ .  
EG:  $n = 4, k = 2$  gives us two orbits:

0011	0101
1001 $\mapsto 2$	1010 $\mapsto 3$
1100 $\mapsto 4$	0101 $\mapsto 1$
0110 $\mapsto 2$	
0011 $\mapsto 0$	

We find another unexpected structure in the orbits if we consider  $S = \binom{[n]}{k}$ , as length  $n$  binary strings with  $k$  1's as we cyclically shift them.  $\tau := C_R : S \rightarrow S$  by  $b = b_1 b_2 \cdots b_n \mapsto b_n b_1 b_2 \cdots b_{n-1}$ , and count  $f(b) = \#inversions(b) = \#\{i < j : b_i > b_j\}$ .  
 EG:  $n = 4, k = 2$  gives us two orbits:

0011	0101
1001 $\mapsto$ 2	1010 $\mapsto$ 3
1100 $\mapsto$ 4	0101 $\mapsto$ 1
0110 $\mapsto$ 2	AVG = $\frac{4}{2} = 2$
0011 $\mapsto$ 0	
AVG = $\frac{8}{4} = 2$	

EG:  $n = 6, k = 2$  gives us three orbits:

000011 000101 001001

EG:  $n = 6, k = 2$  gives us three orbits:

000011	000101	001001
100001	100010	100100
110000	010001	010010
011000	101000	001001
001100	010100	
000110	001010	
000011	000101	

EG:  $n = 6, k = 2$  gives us three orbits:

000011	000101	001001
100001	100010	100100
110000	010001	010010
011000	101000	001001
001100	010100	
000110	001010	
000011	000101	

EG:  $n = 6, k = 2$  gives us three orbits:

000011	000101	001001
100001 $\mapsto$ 4	100010 $\mapsto$ 5	100100 $\mapsto$ 6
110000 $\mapsto$ 8	010001 $\mapsto$ 3	010010 $\mapsto$ 4
011000 $\mapsto$ 6	101000 $\mapsto$ 7	001001 $\mapsto$ 2
001100 $\mapsto$ 4	010100 $\mapsto$ 5	
000110 $\mapsto$ 2	001010 $\mapsto$ 3	
000011 $\mapsto$ 0	000101 $\mapsto$ 1	

EG:  $n = 6, k = 2$  gives us three orbits:

000011	000101	001001
100001 $\mapsto$ 4	100010 $\mapsto$ 5	100100 $\mapsto$ 6
110000 $\mapsto$ 8	010001 $\mapsto$ 3	010010 $\mapsto$ 4
011000 $\mapsto$ 6	101000 $\mapsto$ 7	001001 $\mapsto$ 2
001100 $\mapsto$ 4	010100 $\mapsto$ 5	
000110 $\mapsto$ 2	001010 $\mapsto$ 3	
000011 $\mapsto$ 0	000101 $\mapsto$ 1	
AVG = $\frac{24}{6} = 4$	AVG = $\frac{24}{6} = 4$	AVG = $\frac{12}{3} = 4$



## More rotation

EG:  $n = 6, k = 2$  gives us three orbits:

000011	000101	001001
100001 $\mapsto$ 4	100010 $\mapsto$ 5	100100 $\mapsto$ 6
110000 $\mapsto$ 8	010001 $\mapsto$ 3	010010 $\mapsto$ 4
011000 $\mapsto$ 6	101000 $\mapsto$ 7	001001 $\mapsto$ 2
001100 $\mapsto$ 4	010100 $\mapsto$ 5	
000110 $\mapsto$ 2	001010 $\mapsto$ 3	
000011 $\mapsto$ 0	000101 $\mapsto$ 1	
AVG = $\frac{24}{6} = 4$	AVG = $\frac{24}{6} = 4$	AVG = $\frac{12}{3} = 4$

Note that in each of the three orbits average of the statistic  $f$  is the same.

### Definition ([PrRo15])

Given an (invertible) action  $\tau$  on a finite set of objects  $S$ , call a statistic  $f : S \rightarrow \mathbb{C}$  **homomesic** with respect to  $(S, \tau)$  iff the average of  $f$  over each  $\tau$ -orbit  $\mathcal{O}$  is the same constant  $c$  for all  $\mathcal{O}$ , i.e.,  $\frac{1}{\#\mathcal{O}} \sum_{s \in \mathcal{O}} f(s) = c$  does not depend on the choice of  $\mathcal{O}$ .

(Call  $f$   $c$ -mesic for short.)

## Definition ([PrRo15])

Given an (invertible) action  $\tau$  on a finite set of objects  $S$ , call a statistic  $f : S \rightarrow \mathbb{C}$  **homomesic** with respect to  $(S, \tau)$  iff the average of  $f$  over each  $\tau$ -orbit  $\mathcal{O}$  is the same constant  $c$  for all  $\mathcal{O}$ , i.e.,  $\frac{1}{\#\mathcal{O}} \sum_{s \in \mathcal{O}} f(s) = c$  does not depend on the choice of  $\mathcal{O}$ .

(Call  $f$   $c$ -mesic for short.)

Equivalently: the average of  $f$  over each  $\tau$ -orbit  $\mathcal{O}$  is the same as the average over the entire set  $S$ :

$$\frac{1}{\#\mathcal{O}} \sum_{s \in \mathcal{O}} f(s) = \frac{1}{\#S} \sum_{s \in S} f(s).$$



So we can compute what the average should be before checking whether a statistic is homomesic.

Since its initial codification about 5 years ago, a large number of examples of the homomesy phenomenon have been identified across dynamical algebraic combinatorics. These include:

- Promotion of SSYT; Rowmotion of “nice” (e.g., minuscule heap) posets [PrRo15, StWi11, RuWa15+] ;

Since its initial codification about 5 years ago, a large number of examples of the homomesy phenomenon have been identified across dynamical algebraic combinatorics. These include:

- Promotion of SSYT; Rowmotion of “nice” (e.g., minuscule heap) posets [PrRo15, StWi11, RuWa15+] ;
- In general, composing certain involutions called “toggles” on the set leads to operations with interesting homomesy [Str15+];

Since its initial codification about 5 years ago, a large number of examples of the homomesy phenomenon have been identified across dynamical algebraic combinatorics. These include:

- Promotion of SSYT; Rowmotion of “nice” (e.g., minuscule heap) posets [PrRo15, StWi11, RuWa15+] ;
- In general, composing certain involutions called “toggles” on the set leads to operations with interesting homomesy [Str15+];
- Toggling the “arcs” in noncrossing partitions [E+15+];

Since its initial codification about 5 years ago, a large number of examples of the homomesy phenomenon have been identified across dynamical algebraic combinatorics. These include:

- Promotion of SSYT; Rowmotion of “nice” (e.g., minuscule heap) posets [PrRo15, StWi11, RuWa15+] ;
- In general, composing certain involutions called “toggles” on the set leads to operations with interesting homomesy [Str15+];
- Toggling the “arcs” in noncrossing partitions [E+15+];
- Toggling independent sets of a path graph [JoRo16+]; and

Since its initial codification about 5 years ago, a large number of examples of the homomesy phenomenon have been identified across dynamical algebraic combinatorics. These include:

- Promotion of SSYT; Rowmotion of “nice” (e.g., minuscule heap) posets [PrRo15, StWi11, RuWa15+] ;
- In general, composing certain involutions called “toggles” on the set leads to operations with interesting homomesy [Str15+];
- Toggling the “arcs” in noncrossing partitions [E+15+];
- Toggling independent sets of a path graph [JoRo16+]; and
- Liftings of homomesy from combinatorial actions to piecewise linear and birational maps [EiPr13, GrRo16, GrRo15b].



Since its initial codification about 5 years ago, a large number of examples of the homomesy phenomenon have been identified across dynamical algebraic combinatorics. These include:

- Promotion of SSYT; Rowmotion of “nice” (e.g., minuscule heap) posets [PrRo15, StWi11, RuWa15+] ;
- In general, composing certain involutions called “toggles” on the set leads to operations with interesting homomesy [Str15+];
- Toggling the “arcs” in noncrossing partitions [E+15+];
- Toggling independent sets of a path graph [JoRo16+]; and
- Liftings of homomesy from combinatorial actions to piecewise linear and birational maps [EiPr13, GrRo16, GrRo15b].
- There are many others.

The Twelfold Way is a framework for basic combinatorial counting problems, conceived by G.-C. Rota. It counts  $\#\{f : [n] \rightarrow [x]\}$ , where objects may be considered distinct or indistinct, and the functions arbitrary, injective, or surjective.

Quoting *EC1ed2*: [Stan11, § 1.9]

We are now ready to present the Twelfold Way. The twelve entries are numbered and will be discussed individually. The table gives the number of inequivalent functions  $f : N \rightarrow X$  of the appropriate type, where  $\#N = n$  and  $\#X = x$ .

**The Twelfold Way**

Elements of $N$	Elements of $X$	Any $f$	Injective $f$	Surjective $f$
dist.	dist.	<sup>1.</sup> $x^n$	<sup>2.</sup> $(x)_n$	<sup>3.</sup> $x!S(n, x)$
indist.	dist.	<sup>4.</sup> $\binom{x}{n}$	<sup>5.</sup> $\binom{x}{n}$	<sup>6.</sup> $\binom{x}{n-x}$
dist.	indist.	<sup>7.</sup> $S(n, 0) + S(n, 1) + \cdots + S(n, x)$	<sup>8.</sup> 1 if $n \leq x$ 0 if $n > x$	<sup>9.</sup> $S(n, x)$
indist.	indist.	<sup>10.</sup> $p_0(n) + p_1(n) + \cdots + p_x(n)$	<sup>11.</sup> 1 if $n \leq x$ 0 if $n > x$	<sup>12.</sup> $p_x(n)$

The Twelfold Way is a framework for basic combinatorial counting problems, conceived by G.-C. Rota. It counts  $\#\{f : [n] \rightarrow [x]\}$ , where objects may be considered distinct or indistinct, and the functions arbitrary, injective, or surjective. Quoting *EC1ed2*: [Stan11, § 1.9]

We are now ready to present the Twelfold Way. The twelve entries are numbered and will be discussed individually. The table gives the number of inequivalent functions  $f : N \rightarrow X$  of the appropriate type, where  $\#N = n$  and  $\#X = x$ .

The Twelfold Way

Elements of $N$	Elements of $X$	Any $f$	Injective $f$	Surjective $f$
dist.	dist.	<sup>1.</sup> $x^n$	<sup>2.</sup> $(x)_n$	<sup>3.</sup> $x!S(n, x)$
indist.	dist.	<sup>4.</sup> $\left(\binom{x}{n}\right)$	<sup>5.</sup> $\binom{x}{n}$	<sup>6.</sup> $\left(\binom{x}{n-x}\right)$
dist.	indist.	<sup>7.</sup> $S(n, 0) + S(n, 1) + \cdots + S(n, x)$	<sup>8.</sup> $\begin{matrix} 1 & \text{if } n \leq x \\ 0 & \text{if } n > x \end{matrix}$	<sup>9.</sup> $S(n, x)$
indist.	indist.	<sup>10.</sup> $p_0(n) + p_1(n) + \cdots + p_x(n)$	<sup>11.</sup> $\begin{matrix} 1 & \text{if } n \leq x \\ 0 & \text{if } n > x \end{matrix}$	<sup>12.</sup> $p_x(n)$

In the rest of this talk we will consider actions on some of the basic things counted: permutations, injections, and surjections.

## Definition

For  $w \in \mathfrak{S}_n$ , its **canonical (disjoint) cycle decomposition (CCD)** satisfies:

- (a) each cycle is written with its largest element first; and
- (b) the cycles are written in increasing order of largest element.

The map  $\mathcal{F} : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$  simply removes the parentheses from the CCD of  $w$  and regards the resulting word as a permutation in one-line notation.

$$w = 847296513 = (42)(6)(81)(9375) \xrightarrow{\mathcal{F}} 426819375 = (2)(951487369),$$

Note that here  $w$  has 4 cycles, and  $\mathcal{F}(w)$  has 4 **records** (i.e., left-to-right maxima) viz., 4, 6, 8, and 9.

It is easy to see that  $\mathcal{F}$  is a bijection.

We consider actions  $\mathfrak{S}_n$  of the following form:

$$\mathfrak{S}_n \xrightarrow{\mathcal{F}} \mathfrak{S}_n \xrightarrow{\mathcal{A}} \mathfrak{S}_n \xrightarrow{\mathcal{F}^{-1}} \mathfrak{S}_n \xrightarrow{\mathcal{B}} \mathfrak{S}_n$$

where  $\mathcal{A}$  and  $\mathcal{B}$  are **dihedral involutions**, defined below.

- Ⓐ  $\mathcal{C} : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ , which takes a permutation  $w = w_1 \dots w_n$  to its **complement** whose value in position  $i$  is  $n + 1 - w_i$ ;
- Ⓑ  $\mathcal{R} : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ , which takes a permutation  $w = w_1 \dots w_n$  to its **reversal** whose value in position  $i$  is  $w_{n+1-i}$ ;
- Ⓒ  $\mathcal{Q}^2 : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ , which takes a permutation  $w = w_1 \dots w_n$  to its **rotation by 180-degrees**, whose value in position  $i$  is  $n + 1 - w_{n+1-i}$ .
- Ⓓ  $\mathcal{I} : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ , which takes a permutation  $w$  to its **inverse**  $w^{-1}$ ;
- Ⓔ  $\mathcal{D} : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ , which takes a permutation  $w$  to its **rotated-inverse**  $\mathcal{Q}^2(\mathcal{I}(w))$ .

We call such fourfold compositions, where  $\mathcal{A}$  and  $\mathcal{B}$  are from the above list, **Foatic**.

## Foatic example

$\mathcal{A} = \mathcal{C}$  and  $\mathcal{B} = \mathcal{I}$  gives the Foatic map  $\gamma := \mathcal{I} \circ \mathcal{F}^{-1} \circ \mathcal{C} \circ \mathcal{F}$ .

If  $n = 5$ , then  $\gamma[(4213)(5)] = (2)(4)(513)$  as follows

$$(4213)(5) \xrightarrow{\mathcal{F}} 42135 \xrightarrow{\mathcal{C}} 24531 \xrightarrow{\mathcal{F}^{-1}} (2)(4)(531) \xrightarrow{\mathcal{I}} (2)(4)(513)$$

The orbit (of size six) generated by  $(4213)(5)$  is

$$(4213)(5) \xrightarrow{\gamma} (2)(4)(513) \xrightarrow{\gamma} (412)(53) \xrightarrow{\gamma} (2)(5314) \xrightarrow{\gamma} (431)(52) \xrightarrow{\gamma} (2)(3)(541)$$

## Foatic example

$\mathcal{A} = \mathcal{C}$  and  $\mathcal{B} = \mathcal{I}$  gives the Foatic map  $\gamma := \mathcal{I} \circ \mathcal{F}^{-1} \circ \mathcal{C} \circ \mathcal{F}$ .

If  $n = 5$ , then  $\gamma[(4213)(5)] = (2)(4)(513)$  as follows

$$(4213)(5) \xrightarrow{\mathcal{F}} 42135 \xrightarrow{\mathcal{C}} 24531 \xrightarrow{\mathcal{F}^{-1}} (2)(4)(531) \xrightarrow{\mathcal{I}} (2)(4)(513)$$

The orbit (of size six) generated by  $(4213)(5)$  is

$$(4213)(5) \xrightarrow{\gamma} (2)(4)(513) \xrightarrow{\gamma} (412)(53) \xrightarrow{\gamma} (2)(5314) \xrightarrow{\gamma} (431)(52) \xrightarrow{\gamma} (2)(3)(541)$$

Define the statistic “Fix  $w$ ” to count the number of fixed points (equiv. 1-cycles) of  $w$ .

## Foatic example

$\mathcal{A} = \mathcal{C}$  and  $\mathcal{B} = \mathcal{I}$  gives the Foatic map  $\gamma := \mathcal{I} \circ \mathcal{F}^{-1} \circ \mathcal{C} \circ \mathcal{F}$ .

If  $n = 5$ , then  $\gamma[(4213)(5)] = (2)(4)(513)$  as follows

$$(4213)(5) \xrightarrow{\mathcal{F}} 42135 \xrightarrow{\mathcal{C}} 24531 \xrightarrow{\mathcal{F}^{-1}} (2)(4)(531) \xrightarrow{\mathcal{I}} (2)(4)(513)$$

The orbit (of size six) generated by  $(4213)(5)$  is

$$(4213)(5) \xrightarrow{\gamma} (2)(4)(513) \xrightarrow{\gamma} (412)(53) \xrightarrow{\gamma} (2)(5314) \xrightarrow{\gamma} (431)(52) \xrightarrow{\gamma} (2)(3)(541)$$

Define the statistic “Fix  $w$ ” to count the number of fixed points (equiv. 1-cycles) of  $w$ .

### Conjecture

*The statistic Fix is homomesic with respect to the Foatic complement-inversion action on  $\mathfrak{S}_n$ .*



The Foatic action with the nicest orbit structures and properties appears to be *Reversal-Inversion*:

$$\bar{\varphi} : \mathfrak{S}_n \xrightarrow{\mathcal{F}} \mathfrak{S}_n \xrightarrow{\mathcal{R}} \mathfrak{S}_n \xrightarrow{\mathcal{F}^{-1}} \mathfrak{S}_n \xrightarrow{\mathcal{I}} \mathfrak{S}_n.$$

$$\varphi : \mathfrak{S}_n \xrightarrow{\mathcal{R}} \mathfrak{S}_n \xrightarrow{\mathcal{F}^{-1}} \mathfrak{S}_n \xrightarrow{\mathcal{I}} \mathfrak{S}_n \xrightarrow{\mathcal{F}} \mathfrak{S}_n$$

$$\begin{aligned} w = (2)(43)(51) &\mapsto 24351 \mapsto 15342 \mapsto (1)(5342) \mapsto (1)(5243) = \bar{\varphi}(w) \\ (1)(5243) &\mapsto 15243 \mapsto 34251 \mapsto (3)(42)(51) \mapsto (3)(42)(51) = \bar{\varphi}^2(w) \\ (3)(42)(51) &\mapsto 34251 \mapsto 15243 \mapsto (1)(5243) \mapsto (1)(5342) = \bar{\varphi}^3(w) \\ (1)(5342) &\mapsto 15342 \mapsto 24351 \mapsto (2)(43)(51) \mapsto (2)(43)(51) = \bar{\varphi}^4(w) \end{aligned}$$

This example also displays (down the second column) the conjugate orbit of  $\varphi$ , also of size 4.

$$24351 \xrightarrow{\varphi} 15243 \xrightarrow{\varphi} 34251 \xrightarrow{\varphi} 15342 \uparrow$$

# Data on Orbit Sizes for Foatic Reversal-Inversion


$n$	1	2	3	4	5	6	7	8	9	10
# of orbits:	1	1	2	5	19	84	448	2884	21196	174160
LCM of orbit sizes:	1	2	4	8	16	32	64	128	256	512
GCD of orbit sizes:	1	2	2	4	4	4	4	8	8	8
Longest orbit size:	1	2	4	8	16	32	64	128	256	512
Shortest orbit size:	1	2	2	4	4	4	4	8	8	8
Size of id's orbit:	1	2	4	8	16	32	64	128	256	512

### Definition

Recursively define the **heap** of  $w \in \mathfrak{S}_n$ ,  $H(w)$  as follows: Set  $H(\emptyset) = \emptyset$  (the empty word). If  $w \neq \emptyset$ , let  $m$  be the largest element of  $w$ , so  $w$  can be written uniquely as  $umv$ , where  $u$  and  $v$  are partial permutations (possibly empty). Set  $m$  to be the root of  $H(w)$ , with  $H(u)$  its left subtree and  $H(v)$  its right subtree.


The heap of a permutation will turn out to be a *decreasing binary tree*, (labels decrease along any path from root).

The heap associated with  $w = 314975826$  is shown below.



# Orbits represented as heaps

Two orbits (one for  $\mathfrak{S}_7$ , one for  $\mathfrak{S}_9$ ) of the Foatic reversal-inversion map  $\bar{\varphi}$  with associated heaps, with **fixed points** marked in red. Each orbit has an average of one fixed point per permutation.



## Theorem (La Croix-R.)

The orbits of the action of  $\bar{\varphi}$  (or  $\varphi$ ) on  $\mathfrak{S}_n$ , satisfy the following:

- ① The size of a  $\varphi$ -orbit  $\mathcal{O}$  (equivalently  $\bar{\varphi}$ -orbit) is  $2^h$ , where  $h$  is the number of edges in a maximal path from the root (to a leaf) for any  $w \in \mathcal{O}$ .
- ② Let  $\text{Fix } w$  denote the number of fixed points, i.e., 1-cycles, of  $w$ . Then the statistic  $\text{Fix}$  is 1-mesic with respect to the action of  $\bar{\varphi}$ ; (Equivalently,  $\text{Rasc} = \# \text{record-ascents}$  is 1-mesic with respect action of  $\varphi$ .)
- ③ For fixed  $i \neq j$  in  $[n]$ , let  $\mathbb{1}_{i < j}(u)$  denote the indicator statistic of whether  $i$  occurs to the left of  $j$  in the one-line notation of  $u$ . Then  $\mathbb{1}_{i < j}$  is  $\frac{1}{2}$ -mesic with respect to the action of  $\varphi$ .
- ④ Similarly for fixed  $i \in [n]$ , let  $\mathbb{1}_{(i,n)}$  denote the indicator statistic of whether  $i$  and  $n$  lie in the same cycle of  $w$ . Then  $\mathbb{1}_{(i,n)}$  is  $\frac{1}{2}$ -mesic with respect to the action of  $\bar{\varphi}$ .

## Reveral-Inversion Recursion (Key Lemma)

All the results listed above follow without difficulty from the following key lemma.

### Lemma

*Let  $w \in \mathfrak{S}_n$  have the form  $AnB$  (in one-line notation), where  $A$  and  $B$  are (possibly empty) partial permutations of  $n$ . Then the action of  $\varphi$  satisfies  $\varphi(AnB) = \varphi(B)nA$ . Thus,  $H(\varphi(AnB))$  is the heap interchanging the left and right subtrees at  $v$ , leaving the former unchanged and applying  $\varphi$  recursively to the latter. In particular, the action of  $\varphi$  preserves the underlying unlabeled graph of the corresponding heaps.*

## Whirling action on injections/surjections

We write functions  $f \in [k]^{[n]}$  in one-line notation  $f(1)f(2)\cdots f(n)$ .

### Definition

Let  $\mathcal{J}$  denote either  $\text{Inj}(n, k)$  or  $\text{Sur}(n, k)$  for a given  $n, k \in \mathbb{P}$ . Define a map  $\text{wh}_i : \mathcal{J} \rightarrow \mathcal{J}$ , called **whirling at index  $i$**  in the following way. Given  $f \in \mathcal{J}$ , repeatedly add 1 (mod  $k$ ) to the value of  $f(i)$  until we get a function in  $\mathcal{J}$ . The new function is  $\text{wh}_i(f)$ .

EG:  $f = 124 \in \text{Inj}(3, 6) \implies \text{wh}_1(124) = 324, \text{wh}_2(124) = 134,$   
and  $\text{wh}_3(124) = 125$ .

These generalize *toggle operations*, which are involutions. The composition  $\text{wh} := \text{wh}_n \text{wh}_{n-1} \cdots \text{wh}_2 \text{wh}_1$  is called **whirling**.

EG:  $\text{wh}(124) = (324 \cdots 354 \cdots 356) = 356$ . 124 generates the whirling orbit

$124 \mapsto 356 \mapsto 412 \mapsto 534 \mapsto 651 \mapsto 263 \mapsto 415 \mapsto 621 \mapsto 342 \mapsto 563$

EG: Let  $v = 21444323 \in \text{Sur}(8, 4)$ . Then  $\text{wh}_1(v) = 31444323$ , while  $\text{wh}_2(v) = v$ . The orbit generated by  $v$  is:

$21444323 \mapsto 31114424 \mapsto 32211134 \mapsto 43222141 \mapsto 13332242 \mapsto 14433312 \mapsto \dots$

### Theorem (Joseph-R.)

*Fix  $\mathcal{J}$  to be either  $\text{Inj}(n, k)$  or  $\text{Sur}(n, k)$  for a given  $n, k \in \mathbb{P}$ . For  $i \in [k]$ , define  $\eta_i(f) = \#f^{-1}(\{i\})$  to be the number of times  $i$  appears as an output of the function  $f$ . Then  $\eta_i$  is  $\frac{n}{k}$ -mesic for any  $i \in [k]$ .*

Equivalently,  $\eta_i - \eta_j$  is 0-mesic for any  $i, j \in [k]$ , i.e.,  $i$  and  $j$  appear as outputs of functions the same number of times across any orbit.



## Proving homomesy for injections

**Key Idea:** Partition the orbit into  $[k]$ -chunks. If a value  $j$  appears in  $i$ th spot, then  $j + 1 \bmod k$  must occur directly below, unless it was already in the row when  $w_i$  was applied. Thus, the next  $j + 1$  occurs no later than the  $n$ th letter after  $j$ . Color these the same.

124

356

412

534

651

263

415

621

342

563

It's easy to see this relation goes backwards as well as forwards, so partitions the orbit into chunks each containing all of  $[k]$ . (The chunks can wrap around from bottom to top.)

## Proving homomesy for surjections

This uses a somewhat different partitioning argument. Since  $v_i$  and  $\text{wh}(v)_i$  either agree or differ by one, we could just partition into vertical chunks, except when  $v_i = \text{wh}(v)_i$  (i.e., same values on top of one another). So it suffices to show that the number of pairs of a  $j$  directly below another  $j$  is the same for all  $j \in [k]$ . The tops of such pairs are circled below in red.

Finally, one shows that every circled  $j$  is followed within the next  $n$  slots by a circled  $j + 1$ , allowing these to be partitioned as well.

2	①	4	4	④	3	②	3
③	1	1	①	4	4	2	④
3	2	②	1	1	①	3	4
4	③	2	2	②	1	④	1
①	3	3	③	2	2	4	②
1	4	④	3	3	③	1	2

## A consequence of homomesy for orbits

From this homomesy we can deduce information about orbit sizes (that we currently don't know by any other means).

Let  $\ell(\mathcal{O})$  be the length of the orbit  $\mathcal{O}$ .

If we consider surjective functions from  $[7]$  to  $[4]$ , then across every orbit, the numbers 1, 2, 3, 4 all appear as outputs the same number of times; hence,  $4 \mid 7\ell(\mathcal{O}) \implies 4 \mid \ell(\mathcal{O})$ .

## A consequence of homomesy for orbits

From this homomesy we can deduce information about orbit sizes (that we currently don't know by any other means).

Let  $\ell(\mathcal{O})$  be the length of the orbit  $\mathcal{O}$ .

If we consider surjective functions from  $[7]$  to  $[4]$ , then across every orbit, the numbers 1, 2, 3, 4 all appear as outputs the same number of times; hence,  $4 \mid 7\ell(\mathcal{O}) \implies 4 \mid \ell(\mathcal{O})$ .

On the other hand, if we consider surjective functions from  $[8]$  to  $[4]$ , then across every orbit,  $4 \mid 8\ell(\mathcal{O})$ , which gives no new information.

A function  $f : S \rightarrow T$  between two sets  $S$  and  $T$  is  **$m$ -injective** if  $\#f^{-1}(t) \leq m$  for every  $t \in T$  and  **$m$ -surjective** if  $\#f^{-1}(t) \geq m$  for every  $t \in T$ . Let  $\text{Inj}_m(n, k)$  and  $\text{Sur}_m(n, k)$  denote the set of  $m$ -injective (resp.  $m$ -surjective) functions from  $[n]$  to  $[k]$ .

### Conjecture (Joseph)

Fix  $\mathcal{J}$  to be either  $\text{Inj}_m(n, k)$  or  $\text{Sur}_m(n, k)$  for fixed  $n, k, m \in \mathbb{P}$ . For  $i \in [k]$ , define  $\eta_i(f) = \#f^{-1}(\{i\})$  to be the number of times  $i$  appears as an output of the function  $f$ . Then  $\eta_i$  is  $\frac{n}{k}$ -mesic for any  $i \in [k]$ .

We're happy to talk about this further with anyone who's interested.

Slides for this talk are available online (or will be soon) at

<http://www.math.uconn.edu/~troby/research.html>

We're happy to talk about this further with anyone who's interested.

Slides for this talk are available online (or will be soon) at






<http://www.math.uconn.edu/~troby/research.html>






Thanks very much for coming to this talk!

## References

-  Jonathan Bloom, Oliver Pechenik, and Dan Saracino, *Proofs and generalizations of a homomesy conjecture of Propp and Roby*, Discrete Math., **339** (2016), 194–206.
-  Melody Chan, Shahrzad Haddadan, Sam Hopkins, and Luca Moci, *The expected jaggedness of order ideals*, 2015, arXiv:1507.00249v1.
-  David Einstein, Miriam Farber, Emily Gunawan, Michael Joseph, Matthew Macauley, James Propp, and Simon Rubinstein-salzedo, *Noncrossing partitions, toggles, and homomesies*, in preparation.
-  David Einstein and James Propp, *Combinatorial, piecewise-linear, and birational homomesy for products of two chains*, 2013, arXiv:1310.5294.
-  David Einstein and James Propp, *Piecewise-linear and birational toggling (Extended abstract)*, DMTCS proc. FPSAC 2014, <http://www.dmtcs.org/dmtcs-ojs/index.php/proceedings/article/view/dmAT0145/4518>.



-  Darij Grinberg and Tom Roby, *Iterative properties of birational rowmotion I: generalities and skeletal posets*, Electron. J. of Combin. **23**(1), #P1.33 (2016). <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v23i1p33>
-  Darij Grinberg and Tom Roby, *Iterative properties of birational rowmotion II: rectangles and triangles*, Elec. J. Combin. **22**(3), #P3.40, 2015. <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v22i3p40>
-  Shahrzad Haddadan, *Some Instances of Homomesy Among Ideals of Posets*, 2014, arXiv:1410.4819v2.
-  Michael Joseph and Tom Roby, *Homomesy in toggling independent sets of a path graph*, in preparation.
-  Michael La Croix and Tom Roby, *Foatic actions of the symmetric group and fixed-point homomesy*, in preparation.

-  James Propp and Tom Roby, *Homomesy in products of two chains*, Electronic J. Combin. **22(3)** (2015), #P3.4, <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v22i3p4>. Also available at arXiv:1310.5201v5.
-  David B. Rush and Kelvin Wang, *On orbits of order ideals of minuscule posets II: Homomesy*, arXiv:1509.08047.
-  Richard P. Stanley, *Enumerative Combinatorics, volume 1, 2nd edition*, no. 49 in Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2011. Also available at <http://math.mit.edu/~rstan/ec/ec1/>.
-  Jessica Striker, *Rowmotion and generalized toggle groups*, arXiv:1601.03710.
-  Jessica Striker and Nathan Williams, *Promotion and Rowmotion*, Europ. J. of Combin. 33 (2012), 1919–1942, <http://www.sciencedirect.com/science/article/pii/S0195669812000972>. Also available at arXiv:1108.1172v3.