

Survival Models

Lecture: Weeks 2-3

Chapter summary

- Survival models
 - Age-at-death random variable
 - Time-until-death random variables
 - Force of mortality (or hazard rate function)
 - Some parametric models
 - De Moivre's (Uniform), Exponential, Weibull, Makeham, Gompertz
 - Generalization of De Moivre's
 - Curtate future lifetime
- Chapter 2 (Dickson, Hardy and Waters = DHW)

Age-at-death random variable

- X is the **age-at-death random variable**; continuous, non-negative
- X is interpreted as the lifetime of a newborn (individual from birth)
- Distribution of X is often described by its survival distribution function (SDF):

$$S_0(x) = \Pr[X > x]$$

- other term used: **survival function**
- Properties of the survival function:
 - $S_0(0) = 1$: probability a newborn survives 0 years is 1.
 - $S_0(\infty) = \lim_{x \rightarrow \infty} S_0(x) = 0$: all lives eventually die.
 - non-increasing function of x : not possible to have a higher probability of surviving for a longer period.

Cumulative distribution and density functions

- Cumulative distribution function (CDF): $F_0(x) = \Pr[X \leq x]$
 - nondecreasing; $F_0(0) = 0$; and $F_0(\infty) = 1$.
- Clearly we have: $F_0(x) = 1 - S_0(x)$
- Density function: $f_0(x) = \frac{dF_0(x)}{dx} = -\frac{dS_0(x)}{dx}$
 - non-negative: $f_0(x) \geq 0$ for any $x \geq 0$
 - in terms of CDF: $F_0(x) = \int_0^x f_0(z) dz$
 - in terms of SDF: $S_0(x) = \int_x^\infty f_0(z) dz$

Force of mortality

- The **force of mortality** for a newborn at age x :

$$\mu_x = \frac{f_0(x)}{1 - F_0(x)} = \frac{f_0(x)}{S_0(x)} = -\frac{1}{S_0(x)} \frac{dS_0(x)}{dx} = -\frac{d \log S_0(x)}{dx}$$

- Interpreted as the conditional instantaneous measure of death at x .
- For very small Δx , $\mu_x \Delta x$ can be interpreted as the probability that a newborn who has attained age x dies between x and $x + \Delta x$:

$$\mu_x \Delta x \approx \Pr[x < X \leq x + \Delta x | X > x]$$

- Other term used: **hazard rate** at age x .

Some properties of μ_x

Some important properties of the force of mortality:

- non-negative: $\mu_x \geq 0$ for every $x > 0$
- divergence: $\int_0^{\infty} \mu_x dx = \infty$.
- in terms of SDF: $S_0(x) = \exp\left(-\int_0^x \mu_z dz\right)$.
- in terms of PDF: $f_0(x) = \mu_x \exp\left(-\int_0^x \mu_z dz\right)$.

Moments of age-at-death random variable

- The mean of X is called the **complete expectation of life** at birth:

$$\dot{e}_0 = \mathbf{E}[X] = \int_0^{\infty} x f_0(x) dx = \int_0^{\infty} S_0(x) dx.$$

- The RHS of the equation can be derived using integration by parts.
- Variance:

$$\text{Var}[X] = \mathbf{E}[X^2] - (\mathbf{E}[X])^2 = \mathbf{E}[X^2] - (\dot{e}_0)^2.$$

- The median age-at-death m is the solution to

$$S_0(m) = F_0(m) = \frac{1}{2}.$$

Some special parametric laws of mortality

| Law/distribution | μ_x | $S_0(x)$ | Restrictions |
|---------------------------------|------------------|--|--|
| De Moivre (uniform) | $1/(\omega - x)$ | $1 - (x/\omega)$ | $0 \leq x < \omega$ |
| Constant force (exponential) | μ | $\exp(-\mu x)$ | $x \geq 0, \mu > 0$ |
| Gompertz | Bc^x | $\exp\left[-\frac{B}{\log c}(c^x - 1)\right]$ | $x \geq 0, B > 0, c > 1$ |
| Makeham | $A + Bc^x$ | $\exp\left[-Ax - \frac{B}{\log c}(c^x - 1)\right]$ | $x \geq 0, B > 0, c > 1,$ $A \geq -B$ |
| Weibull | kx^n | $\exp\left(-\frac{k}{n+1}x^{n+1}\right)$ | $x \geq 0, k > 0, n > 1$ |







Figure: Makeham's law: $A = 0.002$, $B = 10^{-4.5}$, $c = 1.10$

Illustrative example 1

Suppose X has survival function defined by

$$S_0(x) = \frac{1}{10}(100 - x)^{1/2}, \quad \text{for } 0 \leq x \leq 100.$$

- 1 Explain why this is a legitimate survival function.
- 2 Find the corresponding expression for the density of X .
- 3 Find the corresponding expression for the force of mortality at x .
- 4 Compute the probability that a newborn with survival function defined above will die between the ages 65 and 75.

Solution to be discussed in lecture.

Practice problem - SOA MLC Spring 2016 Question #2

You are given the survival function:

$$S_0(x) = \left(1 - \frac{x}{60}\right)^{1/3}, \quad \text{for } 0 \leq x \leq 60.$$

Calculate $1000\mu_{35}$.

2.2 Future lifetime random variable

- For a person now age x , its **future lifetime** is $T_x = X - x$. For a newborn, $x = 0$, so that we have $T_0 = X$.
- Life-age- x is denoted by (x) .
- SDF: It refers to the probability that (x) will survive for another t years.

$$S_x(t) = \Pr[T_0 > x + t | T_0 > x] = \frac{S_0(x+t)}{S_0(x)} = {}_t p_x = 1 - {}_t q_x$$

- CDF: It refers to the probability that (x) will die within t years.

$$F_x(t) = \Pr[T_0 \leq x + t | T_0 > x] = \frac{S_0(x) - S_0(x+t)}{S_0(x)} = {}_t q_x$$

- continued

- Density:

$$f_x(t) = \frac{dF_x(t)}{dt} = -\frac{dS_x(t)}{dt} = \frac{f_0(x+t)}{S_0(x)}.$$

- Remark: If $t = 1$, simply use p_x and q_x .
- p_x refers to the probability that (x) survives for another year.
- $q_x = 1 - p_x$, on the other hand, refers to the probability that (x) dies within one year.

Conditions to be valid

To reiterate, these are the conditions for a survival function to be considered valid:

- $S_x(0) = 1$: probability a person age x survives 0 years is 1.
- $S_x(\infty) = \lim_{t \rightarrow \infty} S_x(t) = 0$: all lives, regardless of age, eventually die.
- The survival function $S_x(t)$ for a life (x) must be a non-increasing function of t .

2.3 Force of mortality of T_x

- In deriving the force of mortality, we can use the basic definition:

$$\begin{aligned}\mu_x(t) &= \frac{f_x(t)}{S_x(t)} = \frac{f_0(x+t)}{S_0(x)} \cdot \frac{S_0(x)}{S_0(x+t)} \\ &= \frac{f_0(x+t)}{S_0(x+t)} = \mu_{x+t}.\end{aligned}$$

- This is easy to see because the condition of survival to age $x+t$ supercedes the condition of survival to age x .
- This results implies the following very useful formula for evaluating the density of T_x :

$$f_x(t) = {}_t p_x \times \mu_{x+t}$$

Special probability symbol

- The probability that (x) will survive for t years and die within the next u years is denoted by ${}_t|uq_x$. This is equivalent to the probability that (x) will die between the ages of $x + t$ and $x + t + u$.
- This can be computed in several ways:

$$\begin{aligned}
 {}_t|uq_x &= \Pr[t < T_x \leq t + u] \\
 &= \Pr[T_x \leq t + u] - \Pr[T_x < t] \\
 &= {}_{t+u}q_x - {}_tq_x \\
 &= {}_tp_x - {}_{t+u}p_x \\
 &= {}_tp_x \times {}_uq_{x+t}.
 \end{aligned}$$

- If $u = 1$, prefix is deleted and simply use ${}_t|q_x$.

Other useful formulas

- It is easy to see that

$$F_x(t) = \int_0^t f_x(s) ds$$

which in actuarial notation can be written as

$${}_tq_x = \int_0^t {}_s p_x \mu_{x+s} ds$$

- See Figure 2.3 for a very nice interpretation.
- We can generalize this to

$${}_{t|u}q_x = \int_t^{t+u} {}_s p_x \mu_{x+s} ds$$

2.6 Curtate future lifetime

- Curtate future lifetime of (x) is the number of future years completed by (x) prior to death.
- $K_x = \lfloor T_x \rfloor$, the greatest integer of T_x .
- Its probability mass function is

$$\begin{aligned} \Pr[K_x = k] &= \Pr[k \leq T_x < k + 1] = \Pr[k < T_x \leq k + 1] \\ &= S_x(k) - S_x(k + 1) = {}_{k+1}q_x - {}_kq_x = {}_k|q_x, \end{aligned}$$

for $k = 0, 1, 2, \dots$

- Its distribution function is

$$\Pr[K_x \leq k] = \sum_{h=0}^k {}_h|q_x = {}_{k+1}q_x.$$

2.5/2.6 Expectation of life

- The expected value of T_x is called the **complete expectation of life**:

$$\overset{\circ}{e}_x = \mathbb{E}[T_x] = \int_0^{\infty} t f_x(t) dt = \int_0^{\infty} t {}_t p_x \mu_{x+t} dt = \int_0^{\infty} {}_t p_x dt.$$

- The expected value of K_x is called the **curtate expectation of life**:

$$e_x = \mathbb{E}[K_x] = \sum_{k=0}^{\infty} k \cdot \Pr[K_x = k] = \sum_{k=0}^{\infty} k \cdot {}_k q_x = \sum_{k=1}^{\infty} k p_x.$$

- Proof can be derived using discrete counterpart of integration by parts (summation by parts). Alternative proof will be provided in class.
- Variances of future lifetime can be similarly defined.

Illustrative example 2

Let X be the age-at-death random variable with

$$\mu_x = \frac{1}{2(100 - x)}, \quad \text{for } 0 \leq x < 100.$$

- 1 Give an expression for the survival function of X .
- 2 Find $f_{36}(t)$, the density function of future lifetime of (36).
- 3 Compute ${}_{20}p_{36}$, the probability that life (36) will survive to reach age 56.
- 4 Compute $\overset{\circ}{e}_{36}$, the average future lifetime of (36).

Illustrative example 3

Suppose you are given that:

- $\overset{\circ}{e}_0 = 30$; and
- $S_0(x) = 1 - \frac{x}{\omega}$, for $0 \leq x \leq \omega$.

Evaluate $\overset{\circ}{e}_{15}$.

Solution to be discussed in lecture.

Illustrative example 4

For a group of lives aged 40 consisting of 30% smokers (sm) and the rest, non-smokers (ns), you are given:

- For non-smokers, $\mu_x^{\text{ns}} = 0.05$, for $x \geq 40$
- For smokers, $\mu_x^{\text{sm}} = 0.10$, for $x \geq 40$

Calculate q_{65} for a life randomly selected from those who reach age 65.

Temporary (partial) expectation of life

We can also define **temporary (or partial) expectation of life**:

$$E[\min(T_x, n)] = \overset{\circ}{e}_{x:\overline{n}|} = \int_0^n {}_t p_x dt$$

This can be interpreted as the average future lifetime of (x) within the next n years.

Suppose you are given:

$$\mu_x = \begin{cases} 0.04, & 0 < x < 40 \\ 0.05, & x \geq 40 \end{cases}$$

Calculate $\overset{\circ}{e}_{25:\overline{25}|}$

Generalized De Moivre's law

The SDF of the so-called **Generalized De Moivre's Law** is expressed as

$$S_0(x) = \left(1 - \frac{x}{\omega}\right)^\alpha \text{ for } 0 \leq x \leq \omega.$$

Derive the following for this special type of law of mortality:

- 1 force of mortality
- 2 survival function associated with T_x
- 3 expectation of future lifetime of x
- 4 can you find explicit expression for the variance of T_x ?

Illustrative example

- We will do **Example 2.6** in class.

Example 2.3

Let $\mu_x = Bc^x$, for $x > 0$, where B and c are constants such that $0 < B < 1$ and $c > 1$.

Derive an expression for $S_x(t)$.

Typical mortality pattern observed

- High (infant) mortality rate in the first year after birth.
- Average lifetime (nowadays) range between 70-80 - varies from country to country.
- Fewer lives/deaths observed after age 110 - **supercentenarian** is the term used to refer to someone who has reached age 110 or more.
- The highest recorded age at death (I believe) is 122.
- Different male/female mortality pattern - females are believed to live longer.

Substandard mortality

- A **substandard** risk is generally referred to someone classified by the insurance company as having a higher chance of dying because of:
 - some physical condition
 - family or personal medical history
 - risky occupation
 - dangerous habits or lifestyle (e.g. skydiving)
- Mortality functions are superscripted with s to denote substandard: q_x^s and μ_x^s .
- For example, substandard mortality may be obtained from a standard table using:
 - ① adding a constant to force of mortality: $\mu_x^s = \mu_x + c$
 - ② multiplying a fixed constant to probability: $q_x^s = \min(kq_x, 1)$
- The opposite of a substandard risk is **preferred** risk where someone is classified to have better chance of survival.

Practice problem - SOA MLC Fall 2000 Question #4

Mortality for Audra, age 25, follows De Moivre's law with $\omega = 100$. If she takes up hot air ballooning for the coming year, her assumed mortality will be adjusted so that for the coming year only, she will have a constant force of mortality of 0.1.

Calculate the decrease in the 11-year temporary complete life expectancy for Audra if she takes up hot air ballooning.

Illustrative example 5

You are given:

- Mortality for standard lives follows the Standard Ultimate Life Table (SULT).
- The force of mortality for standard lives age $45 + t$ is represented as μ_{45+t}^{SULT} .
- The force of mortality for substandard lives age $45 + t$, μ_{45+t}^{sub} , is defined by

$$\mu_{45+t}^{\text{sub}} = \begin{cases} \mu_{45+t}^{\text{SULT}} + 0.05, & \text{for } 0 \leq t < 1 \\ \mu_{45+t}^{\text{SULT}}, & \text{for } t \geq 1 \end{cases}$$

Calculate the probability that a substandard 45-year-old will die within the next two years.

Practice problem - SOA LTAM Spring 2019 Question #3

You are given:

- A life table uses a Makeham's mortality model with parameters $A = 0.00022$, $B = 2.7 \times 10^{-6}$, $c = 1.124$
- ${}_{10}p_{50} = 0.9803$

Calculate $\frac{d}{dt} {}_tq_{50}$ at $t = 10$.

Final remark - other contexts

- The notion of a lifetime or survival learned in this chapter can be applied in several other contexts:
 - engineering: lifetime of a machine, lifetime of a lightbulb
 - medical statistics: time-until-death from diagnosis of a disease, survival after surgery
 - finance: time-until-default of credit payment in a bond, time-until-bankruptcy of a company
 - space probe: probability radios installed in space continue to transmit
 - biology: lifetime of an organism
 - other actuarial context: disability, sickness/illness, retirement, unemployment

Other symbols and notations used

| Expression | Other symbols used |
|--------------------------------|---------------------------|
| probability function | $P(\cdot)$ $\Pr(\cdot)$ |
| survival function of newborn | $S_X(x)$ $S(x)$ $s(x)$ |
| future lifetime of x | $T(x)$ T |
| curtate future lifetime of x | $K(x)$ K |
| survival function of x | $S_{T_x}(t)$ $S_T(t)$ |
| force of mortality of T_x | $\mu_{T_x}(t)$ $\mu_x(t)$ |