


absorbing

A  
a

basic survival model



## Multiple State Models

Lecture: Weeks 3-4

health sickness  
model



state  $\rightarrow$  countable  
 $n+1$  states

$\{0, 1, \dots, n\}$

reserves for  
where you begin

permanent disability  
or  
critical illness



# Chapter summary

- Multiple state models (also called transition models)

- what are they?
- ✓ • actuarial applications - some examples

- State space - finite  $\{0, 1, \dots, n\}$

- Transition probabilities


- continuous and discrete time space

- Markov chains

- time homogeneous versus non-homogeneous Markov chains

- Cash flows and actuarial present value calculations in multiple state models

- Chapter 8 (Dickson, et al.)




# Introduction


probability

- Multiple state models are probability models that describe the random movements of:
  - a subject (often a person, but could be a machinery, organism, etc.)
  - among various states
- Discrete time or continuous time and discrete state space
- Examples include:
  - basic survival model ✓
  - multiple decrement models
  - health-sickness model ✓
  - disability model ✓
  - pension models
  - multiple life models
  - long term care (or continuing care retirement communities, CCRCs) models


# The basic survival model




# The withdrawal-death model



# The permanent disability model




# The HIV-AIDS progression model



## Notation

Continuous time



- Assume a finite state space (total of  $n + 1$  states):  $\{0, 1, \dots, n\}$
- In most actuarial applications, we need a reference age.
  - Denote by  $x$  the age at which the multiple state process begins.
  - $x$  is the age at time  $t = 0$ .
- Denote by  $Y_x(t)$  the state of the process at time  $t$ .
  - This can take on possible values in the state space.
  - The process can be denoted by  $\{Y_x(t), t \geq 0\}$ .





## Continuous time Markov chain models

# Transition probabilities and forces of transition

- **Transition probabilities:**

$${}_t p_x^{ij} = \Pr[Y_x(t) = j | Y_x(0) = i]$$



- This is the probability that a life age  $x$  at time 0 is in state  $i$  and will be in state  $j$  after  $t$  periods.
- **Force of transition** (also called **transition intensity**):

$$\mu_x^{ij}$$

$$\mu_x^{ij} = \lim_{h \rightarrow 0^+} \frac{1}{h} \underline{{}_h p_x^{ij}}, \quad \text{for } i \neq j$$

$$\lim_{h \rightarrow 0} \frac{{}_h p_x^{ij}}{h} = \mu_x^{ij}$$

- This is defined only in the case where we have a continuous time process.
- Analogous to the force of mortality in the basic survival model.
- It is understood that  $\mu_x^{ij} = 0$  if it is not possible to transition from state  $i$  to state  $j$  at any time.

$$\lim_{h \rightarrow 0} \frac{1}{h} h P_x^{ij} = \mu_x^{ij}$$

~

$$h P_x^{ij} \approx h \cdot \mu_x^{ij} + \underbrace{\text{"small"}}_{o(h)}$$

$$h P_x^{ij}$$

$$\mu_x^{ij}$$

force of transition  
- intensity -

$$\lim_{h \rightarrow 0} \frac{1}{h} o(h) \rightarrow 0$$

/

## Some assumptions



- Assumption 1: The **Markov property** holds.

*history of my visit*

$$\begin{aligned} & \Pr[Y_x(s+t) = j | Y_x(s) = i, Y_x(u) = k, 0 \leq u < s] \\ &= \Pr[Y_x(s+t) = j | Y_x(s) = i] \end{aligned}$$

- Assumption 2: For any positive interval of time length (generally very small)  $h$ ,

$$\Pr[2 \text{ or more transitions within a time period of length } h] = o(h)$$

*"small"*

- Assumption 3: For all states  $i$  and  $j$  and all ages  $x \geq 0$ ,  ${}_t p_x^{ij}$  is a differential function of  $t$ .

*$\frac{d}{dt} p_x^{ij}$  exists*

## Some useful approximation

We can express the transition probabilities in terms of the forces of transition as

$${}_h p_x^{ij} = h \mu_x^{ij} + o(h),$$

so that for very small values of  $h$ , we have the approximation

$${}_h p_x^{ij} \approx h \mu_x^{ij}.$$

# The occupancy probability




When a person currently age  $x$  and is currently in state  $i$ , the probability that the person continuously remains in the same state for a length of  $t$  periods is called an **occupancy probability**.

For any state  $i$  in a multiple state model, the probability that ( $x$ ) now in state  $i$  will remain in state  $i$  for  $t$  years can be computed using:

$${}_t p_x^{\bar{i}i} = \exp \left[ - \int_0^t \sum_{j=0, j \neq i}^n \mu_{x+s}^{ij} ds \right].$$

Sketch of proof will be done in class - also on pages 239 - 240.

$$e^{- \int_0^t \sum_{j \neq i} \mu_{x+s}^{ij} ds}$$



$$t p^{\bar{h}h} \neq t p^{hh}$$

$$t p^{hh} = t p^{\bar{h}h}$$

$$t p_x^{ii} \stackrel{\checkmark}{=} t p_x^{\bar{i}\bar{i}}$$

$$t p_x^{ii} \stackrel{\checkmark}{<} t p_x^{\bar{i}\bar{i}} \checkmark$$

Think!

$$tP_x^{\bar{ii}} = e^{-\int_0^t \sum_{j \neq i} \mu_{x+s}^{ij} ds}$$

start with small  $h$



$$tP_x^{\bar{ii}} = 1 + \sum_{j \neq i} tP_x^{ij} \quad \checkmark$$

$\downarrow$   
 $(h \cdot \mu_x^{ij} + o(h))$

no more than 2 transitions in small  $h$

$$tP_x^{ij} = h \cdot \mu_x^{ij} + o(h)$$

$$= 1 - \sum_{j \neq i} h \cdot \mu_x^{ij} + o(h)$$

$$\frac{d}{dt} tP_x^{\bar{ii}} = \lim_{h \rightarrow 0} \frac{tP_x^{\bar{ii}} - tP_x^{\bar{ii}}}{h} = \lim_{h \rightarrow 0} \frac{1}{h} \left( tP_x^{\bar{ii}} \cdot tP_{x+t}^{\bar{ii}} - tP_x^{\bar{ii}} \right) + o(h)$$

$$= tP_x^{\bar{ii}} * \lim_{h \rightarrow 0} \frac{1}{h} \left( tP_{x+t}^{\bar{ii}} - 1 \right) \quad \checkmark$$



$$\frac{d}{dt} P_{x^{ii}} = -P_{x^{ii}} \times \lim_{h \rightarrow 0} \frac{1}{h} \sum_{j \neq i} h P_{x+t}^{ij}$$

$$\sum_{j \neq i} \lim_{h \rightarrow 0} \frac{1}{h} h P_{x+t}^{ij} = \sum_{j \neq i} \mu_{x+t}^{ij}$$

$$e^{\log_t P_{x^{ii}} - \log_0 P_{x^{ii}}} = e^{-\sum \dots}$$

replace t by s

$$\int_0^t \frac{d}{ds} P_{x^{ii}} = \int_0^t -P_{x^{ii}} \cdot \sum_{j \neq i} \mu_{x+s}^{ij} ds$$

$$\int_0^t \frac{d P_{x^{ii}}}{P_{x^{ii}}} = - \int_0^t \sum_{j \neq i} \mu_{x+s}^{ij} ds$$

$$\log P_{x^{ii}} \Big|_0^t = - \int_0^t \sum_{j \neq i} \mu_{x+s}^{ij} ds$$

$$t p_x^{\bar{ii}} = e^{-\int_0^t \sum_{j \neq i} \mu_{x+s}^{ij} ds}$$

occupancy formula


Illustration

permanent disability

$$\mu_x^{01} = .0279 \quad x \geq 0$$

$$\mu_x^{02} = .0229 \quad x \geq 0$$

$$\mu_x^{12} = \mu_x^{21} = .0229 \quad x \geq 0$$




- ① Calculate the probability that a healthy life (x) will be healthy at the end of 10 years.

$$\begin{aligned}
 {}_{10}p_x^{\infty} &= {}_{10}p_x^{\bar{00}} = e^{-\int_0^{10} (-0.0279 + 0.0229) ds} = e^{-0.0508(10)} \\
 &= \underline{.60170}
 \end{aligned}$$

(2)

$${}_{10}P_x^{01} = ?$$

$${}_{10}P_x^{00} = {}_{10}P_x^{\overline{00}}$$

(3)  ~~${}_{10}P_x^{02}$~~  ?

$$= \int_0^{10} {}_5P_x^{\overline{00}} \cdot \mu_{x+t}^{01} \cdot {}_{10-5}P_{x+t}^{\overline{11}} dt$$



.19363

Wednesday Feb 19

end here

- bring calculator

- two sheets -

10 gunti

occupancy probability

$ij$  status

$${}_t P_x^{\bar{ii}} = \exp \left[ - \int_0^t \sum_{j \neq i} \mu_{x+s}^{ij} ds \right]$$

very important

$$\mu^{01} = .0279$$

$$\mu^{02} = .0229$$

$$\mu^{12} = \mu^{21} = .0229$$



$$\textcircled{1} \quad {}_{10}P_x^{\bar{00}} = e^{-\int_0^{10} (\mu^{01} + \mu^{02}) ds} = e^{-10(.0508)} \approx \underline{.60170}$$

$$\textcircled{2} \quad {}_{10}P_x^{\bar{01}} = \int_0^{10} \underbrace{{}_sP_x^{\bar{00}} \cdot \mu_{x+s}^{01}}_{\text{rate}} \cdot {}_{10-s}P_{x+s}^{\bar{11}} ds$$

$${}_{10}p_x^{01} = \int_0^{10} e^{-.0508s} \cdot .0279 \cdot e^{-.0229(10-s)} ds$$




$$.0508 = .0279 + .0229$$

$$= .0279 e^{-.229} \int_0^{10} e^{-.0279s} ds$$

$$= 0.19363$$


$${}_{10}p_x^{02}$$



$$= \int_0^{10} \int_s^{10} e^{-.0508s} \cdot \mu_{s|}^{01} e^{-.0229(t-s)} \cdot \mu_{t|}^{12} \cdot 1 dt ds + \int_0^{10} e^{-.0508s} \cdot \mu_{s|}^{02} ds \} = \underline{.20467}$$

$$\mu^{HD} = .017$$

$$\mu^{CD} = .055$$



$${}_{10}P^{\overline{HH}} = {}_{10}P^{HH} = .64$$

$$\mu^{HC} = ?$$


$${}_{10}P^{HH} = {}_{10}P^{\overline{HH}} =$$

$$e^{-\int_0^{10} (\mu^{HC} + \mu^{HD}) ds} = .64$$

$$\log = \ln$$

$$e^{-\mu^{HC} \cdot 10} * e^{-.17} = .64$$

$$\Rightarrow \mu^{HC} = -\frac{1}{10} \log \left( \frac{.64}{e^{-.17}} \right) = .02762871 \approx \underline{\underline{.0276}}$$



discoun  
programi

$$\begin{aligned} \mu^{01} &= .005 & \mu^{03} &= .01 \\ \mu^{12} &= .08 & \mu^{13} &= .05 \\ \mu^{23} &= .40 \end{aligned}$$

$${}_{10}P^{02} = ?$$

$$\int_0^{10} \int_s^{10} s P^{00} \cdot \mu^{01} \cdot \underbrace{t-s P^{11}}_{\mu^{12}} \cdot \underbrace{10-t P^{22}}_{\mu^{23}} dt ds$$

$$= \int_0^{10} \int_s^{10} e^{-.0155s} (-.005) e^{-.130(t-s)} \cdot .08 e^{-.40(10-t)} dt ds$$

$$= \underline{\underline{.00433562}}$$



Spring 2015 CT1

Q8, Q10 -

~~Q2~~ not include gains/loss -

Q3


Whole life insurance (40)

SULT -  
 $\delta = .05$

no expenses  
 $B = 1000$  payable at e.o.y.  
Single premium =  $P = 125$

$L_{10}$  = prospective loss at end of year 10

$L_{10} = PVFB_{10} - PVFP_{10}$



Calculate  $\Pr[L_{10} > 200]$

$L_{10} = PVFB_{10} - K_{10}$   
 $= 1000 \cdot v$

$$L_{10} > 200 \iff 1000 v^{k+1} > 200$$

$$\iff v^{k+1} > \frac{200}{1000} = .20$$

$$\iff (k+1) \underbrace{\log v}_{-.05} > \log(.20)$$

$$\iff k < \underbrace{\frac{\log(.20)}{-.05} - 1}_{31.18876}$$

$$Pr[k < 31.18876] = Pr[k \leq 31]$$

$$= {}_{32}p_{50} = 1 - {}_{32}q_{50}$$

$$= 1 - \frac{l_{82}}{l_{50}} = 1 - \frac{70507.2}{98576.4}$$

$${}_n p_x = \frac{l_{x+n}}{l_x}$$



$$= 0.285$$

Q4

fully discrete whole life (40)

- 1st year DB = 20
- other years DB = 10

$$q_{40} = .0020 \quad q_{50} = .0025$$

$$i = .03$$

$$A_{40} = .37$$

$$A_{50} = .48$$

UDD within each year

no expenses

$$\ddot{O}_{40} = \frac{1 - A_{40}}{d}$$

$$d = .03/1.03$$



Calculate  $10.5V = ?$

①  $P = ?$

$$APVFB_0 = APVFB_0$$

$$\dot{P}\ddot{A}_{40} = 10 \dot{A}_{40} + 10 \cdot V \cdot \left( \frac{i}{1.03} \right) q_{40}$$

$$21.63 = 37 + \frac{10 \cdot V \cdot .0020}{1.03}$$

$$P = .1719564$$

$$\begin{aligned}
 \textcircled{2} \quad 10V &= APVFB_{10} - APVFP_{10} \\
 &= 10 \cdot A_{50} - P \check{A}_{50} \\
 &\quad \underbrace{\hspace{10em}}_{1.730005 \approx 1.73}
 \end{aligned}$$

$$\begin{aligned}
 B &= 10 \\
 \check{A}_{50} &= \frac{1 - A_{50}}{d} \\
 &\quad \underbrace{\hspace{2em}}_{17.85333} \\
 P &= .1719564 -
 \end{aligned}$$

$$\textcircled{3} \quad 10.5V = \frac{(10V + P)(1.03)^{.5} - 0.5 \times f_{50} \times 10 \cdot V^{.5}}{1 - .5 \times f_{50}}$$

$$\begin{aligned}
 10V &= 1.73 \\
 P &= .1719564 \\
 f_{50} &= .0025
 \end{aligned}$$

$$\underbrace{\hspace{10em}}_{1.939037}$$

Study again -

$$\underbrace{\hspace{10em}}_{10.4V = ?}$$

Q5

3-year term (62)

$$g_{k+1} = \underline{.025} \quad k=0,1,\dots$$

$$DB = 10 + V \checkmark$$

endowment ~~is~~ benefit = 40 ✓

$$i = .05$$

terminal



$${}_0V = 0$$

3 years

$${}_1V = ({}_0V + P)(1.05) - (10 + \cancel{V} - \cancel{V}) \cdot g$$

$$\frac{P(1.05) - 10 \checkmark}{g}$$

$$\begin{aligned} {}_2V &= ({}_1V + P)(1.05) - (10 + \cancel{2V} - \cancel{V}) \cdot g \\ &= P[1.05^2 + 1.05] - 10(g) \left[ \cancel{1.05} + 1 \right] \end{aligned}$$

$$\therefore {}_3V = \frac{P[1.05^3 + 1.05^2 + 1.05] - 10(g)[1.05^2 + 1.05 + 1]}{g} = 40$$

$$P = 12.3 \checkmark$$

no expenses



$${}_{t+1}V = \frac{({}_tV + P)(1+i) - B \cdot g_{x+t}}{1 - g_{x+t}}$$




$${}_{t+1}V = ({}_tV + P)(1+i) - \left( B - \frac{{}_{t+1}V}{i} \right) g_{x+t}$$

Complement

$$10p^{00} + 10p^{01} + 10p^{12} = 1$$

$\cdot 60170$        $\cdot 19363$        $\cdot 20467$

Example




Multiple Decrement Models -

$$\mu^{01} = .003$$

$$\mu^{02} = .005$$

$$\mu^{03} = .010$$

$$\Rightarrow .019$$



Calculate the probability that

① you stay alive within 4 years

$${}_4p^{00} = {}_4p^{0\bar{0}} = e^{-\int_0^4 (\underbrace{\mu^{01} + \mu^{02} + \mu^{03}}_{.019}) dt} = e^{-.019(4)} = .9268162$$

② you die of cancer within 4 years

$${}_4p^{02} = \int_0^4 {}_s p^{00} \mu^{02} ds = \int_0^4 e^{-.019s} (.006) ds = \frac{.006}{.019} \left( 1 - \underbrace{e^{-.019(4)}}_{.9268162} \right) = ??$$





$$\frac{\mu^{0j} (1 - t p^{60})}{\sum_{j=1}^m \mu^{0j}}$$

$j = 1, 2, \dots, m$   
 = prob of dying from cause  $j$ ,  $j = 1, \dots, m$

③ given you die within 4 years, you die of cancer?  
 $\Pr(A|B) = \frac{\Pr(AB)}{\Pr(B)}$

$$\frac{\mu^{02}}{\mu^{01} + \mu^{02} + \mu^{03}} \Pr(\text{cancer} | \text{die}) = \frac{\Pr(\text{cancer, die})}{\Pr(\text{die})} = \frac{.006 (1 - e^{-.019(4)})}{.019 (1 - e^{-.011(4)})} = \frac{.006}{.019}$$

$$\Pr(\text{heart disease} | \text{die}) = \frac{.003}{.019}$$


$$\Pr(\text{others} | \text{die}) = \frac{.010}{.019}$$

MD  $j = 1, 2, \dots, m$      0 = alive

$$\Pr(\text{cause } j | \text{die}) = \frac{\mu^{0j}}{\sum_{j=1}^m \mu^{0j}}, \quad j = 1, 2, \dots, m$$

$$tP_x^{01}$$

$$\frac{d}{dt} tP_x^{00}$$



$$tP_x^{02}$$

$$\frac{d}{dt} tP_x^{01}$$


$$tP_x^{00} \neq tP_x^{0\bar{0}}$$

$$\frac{d}{dt} tP_x^{02}$$

motivate Kolmogorov

Forward Equations -

For small  $h$ , we assume small probability you transition 2 or more states -



$$t+hP_x^{00} = \frac{tP_x^{00} \cdot hP_{x+t}^{00} + tP_x^{01} \cdot hP_{x+t}^{10}}{h}$$

$$\frac{d}{dt} tP_x^{00} = \lim_{h \rightarrow 0} \left( \frac{t+hP_x^{00} - tP_x^{00}}{h} \right) = \lim_{h \rightarrow 0} \frac{1}{h} \left[ tP_x^{00} (P_{x+t}^{00} - P_{x+t}^{01} - P_{x+t}^{02}) + tP_x^{01} P_{x+t}^{10} - tP_x^{00} \right]$$

$$\sim \lim_{h \rightarrow 0} \left[ tP_x^{01} \frac{P_{x+t}^{10}}{h} - tP_x^{00} \left( \frac{P_{x+t}^{01}}{h} + \frac{P_{x+t}^{02}}{h} \right) \right]$$

$$\frac{d}{dt} \tau P_x^{00} = \underbrace{\tau P_x^{01} \mu_{x+t}^{10}}_{\text{terms out of 0}} - \underbrace{\tau P_x^{00} (\mu_{x+t}^{01} + \mu_{x+t}^{02})}_{\text{stay in 0} \rightarrow \text{then out of 0}}$$

terms out of  
0  $\rightarrow$  back to 0



stay in 0  $\rightarrow$  then out of 0

$$\frac{d}{dt} \tau P_x^{01} = \tau P_x^{00} \mu_{x+t}^{01} - \tau P_x^{01} (\mu_{x+t}^{10} + \mu_{x+t}^{12})$$



$$\frac{d}{dt} \tau P_x^{02} = (\tau P_x^{00} \mu_{x+t}^{02} + \tau P_x^{01} \mu_{x+t}^{12})$$

~~$\tau P_x^{01} \mu_{x+t}^{12}$~~

cannot get out  
of 2!  
=



Kolmogorov forward equations (KFE's)


$i, j$  are two states  
say  $0, 1, 2, \dots, n$

$$\frac{d}{dt} p_x^{ij} = \sum_{k=0, k \neq j}^n p_x^{ik} \mu_{x+t}^{kj} - \sum_{k=0, k \neq j}^n p_x^{ij} \mu_{x+t}^{jk}$$

$+$ 's consist of all those  
you get out of  $i$  & later  
return to  $j$

$-$ 's consist of all those  
you get from  $i \rightarrow j$   
and leave  $j$  thereafter

not needed to memorize, but  
must know how to write KFE's for  
any MS models



$$\frac{d}{dt} {}_tP_x^{00} = ({}_tP_x^{01} \mu_{x+t}^{10} + {}_tP_x^{02} \mu_{x+t}^{20}) - {}_tP_x^{00} (\mu_{x+t}^{01} + \mu_{x+t}^{02})$$

$t \rightarrow 0 \rightarrow \frac{x}{z} \rightarrow 0$

---

$$\frac{d}{dt} {}_tP_x^{01} \quad \text{OR} \quad \frac{d}{dt} {}_tP_x^{02} \quad \text{OR} \quad \frac{d}{dt} {}_tP_x^{12}$$

how to solve?



SEV

- no exact solution  
(in many situations)
- numerically approximate

$$- \frac{d}{dt} p_x^{ij} \approx \frac{1}{h} (p_x^{ij}(t+h) - p_x^{ij}(t)) \quad \text{--- provide } h \text{ is small}$$

-  $h$  is called a step-  
e.g.  $h = 0.01$   
 $h = 0.10$   
 $h = 0.25$

$$\frac{d}{dt} p_x^{ij}$$

- boundary conditions  
at time 0

$$p_x^{ij} = \begin{cases} 1, & i=j \\ 0, & i \neq j \end{cases}$$

Remarks: smaller  $h$  gets you better approximation  
but requires more steps to solve.

# Kolmogorov's forward equations

For a Markov process, transition probabilities can be expressed as

$${}_{t+h}p_x^{ij} = {}_t p_x^{ij} + h \sum_{k=0, k \neq j}^n \left( {}_t p_x^{ik} \mu_{x+t}^{kj} - {}_t p_x^{ij} \mu_{x+t}^{jk} \right) + o(h).$$

This leads us to the **Kolmogorov's Forward Equations** (KFE):

$$\frac{d}{dt} {}_t p_x^{ij} = \sum_{k=0, k \neq j}^n \left( {}_t p_x^{ik} \mu_{x+t}^{kj} - {}_t p_x^{ij} \mu_{x+t}^{jk} \right).$$

This set of differential equations is used to solve for transition probabilities.

## Numerical evaluation of transition probabilities

To solve for the set of KFE's for the transition probabilities, we can equate  $o(h) \rightarrow 0$ , especially if  $h$  is small, or equivalently use the approximation

$$\frac{d}{dt} {}_t p_x^{ij} \approx \frac{1}{h} \left( {}_{t+h} p_x^{ij} - {}_t p_x^{ij} \right) \quad \checkmark$$

This is a similar approach used to approximate the solution to the Thiele's differential equation for reserves

Method is called the **Euler's method**. The primary differences are:

- solution is performed recursively going forward with the boundary conditions:

$${}_0 p_x^{ij} = \begin{cases} 1, & \text{if } i = j \\ 0, & \text{if } i \neq j \end{cases}$$


- the process usually requires solving a number of equations.



## Illustrative example from book

- Consider Example 8.4 on pages 254-255

# The health-sickness model



## Example 8.5 from the book

Consider the health-sickness insurance model illustrated in Example 8.5 with

$$\mu_x^{01} = a_1 + b_1 \exp(c_1 x)$$

$$\mu_x^{10} = 0.10 \mu_x^{01}$$

$$\mu_x^{02} = a_2 + b_2 \exp(c_2 x)$$

$$\mu_x^{12} = \mu_x^{02}$$

where

$$a_1 = 4 \times 10^{-4}, \quad b_1 = 3.4674 \times 10^{-6}, \quad c_1 = 0.138155$$

$$a_2 = 5 \times 10^{-4}, \quad b_2 = 7.5868 \times 10^{-5}, \quad c_2 = 0.087498$$

Verify the calculations of  ${}_{10}p_{60}^{00}$  and  ${}_{10}p_{60}^{01}$ , and follow the same procedure to calculate  ${}_{10}p_{60}^{02}$ .

## Numerical process of solutions

One can verify that to solve for the desired probabilities, one solves the set of Kolmogorov's forward equations

$$\begin{aligned}
 \frac{d}{dt} {}_t p_{60}^{00} &= {}_t p_{60}^{01} \mu_{60+t}^{10} - {}_t p_{60}^{00} (\mu_{60+t}^{01} + \mu_{60+t}^{02}) \\
 \frac{d}{dt} {}_t p_{60}^{01} &= {}_t p_{60}^{00} \mu_{60+t}^{01} - {}_t p_{60}^{01} (\mu_{60+t}^{10} + \mu_{60+t}^{12}) \\
 \frac{d}{dt} {}_t p_{60}^{02} &= {}_t p_{60}^{00} \mu_{60+t}^{02} + {}_t p_{60}^{01} \mu_{60+t}^{12}
 \end{aligned}$$

*Handwritten notes:* The first equation is circled in orange. To its left, the expression  $\frac{{}_t p_{60}^{00} - {}_t p_{60}^{00}}{h}$  is written in orange, with an arrow pointing to the circled term.

Then use the numerical approximations:

$$\begin{aligned}
 {}_{t+h} p_{60}^{00} &\approx {}_t p_{60}^{00} + h [{}_t p_{60}^{01} \mu_{60+t}^{10} - {}_t p_{60}^{00} (\mu_{60+t}^{01} + \mu_{60+t}^{02})] \\
 {}_{t+h} p_{60}^{01} &\approx {}_t p_{60}^{01} + h [{}_t p_{60}^{00} \mu_{60+t}^{01} - {}_t p_{60}^{01} (\mu_{60+t}^{10} + \mu_{60+t}^{12})] \\
 {}_{t+h} p_{60}^{02} &\approx {}_t p_{60}^{02} + h [{}_t p_{60}^{00} \mu_{60+t}^{02} + {}_t p_{60}^{01} \mu_{60+t}^{12}]
 \end{aligned}$$

with initial boundary conditions:  ${}_0 p_{60}^{00} = 1, {}_0 p_{60}^{01} = {}_0 p_{60}^{02} = 0$

Detailed results with step size  $h = 1/12$ *verify in excel.*

$t$	$\mu_{60+t}^{01}$	$\mu_{60+t}^{02}$	$\mu_{60+t}^{10}$	$\mu_{60+t}^{12}$	${}_tP_{60}^{00}$	${}_tP_{60}^{01}$	${}_tP_{60}^{02}$
0	0.01420	0.01495	0.00142	0.01495	1.00000	0.00000	0.00000
1/12	0.01436	0.01506	0.00144	0.01506	0.99757	0.00118	0.00125
2/12	0.01453	0.01517	0.00145	0.01517	0.99512	0.00238	0.00250
3/12	0.01469	0.01527	0.00147	0.01527	0.99266	0.00358	0.00376
4/12	0.01485	0.01538	0.00149	0.01538	0.99018	0.00479	0.00503
5/12	0.01502	0.01549	0.00150	0.01549	0.98769	0.00601	0.00630
6/12	0.01519	0.01560	0.00152	0.01560	0.98518	0.00723	0.00759
7/12	0.01536	0.01571	0.00154	0.01571	0.98265	0.00847	0.00888
8/12	0.01554	0.01582	0.00155	0.01582	0.98011	0.00972	0.01017
9/12	0.01571	0.01593	0.00157	0.01593	0.97755	0.01097	0.01148
10/12	0.01589	0.01605	0.00159	0.01605	0.97497	0.01224	0.01279
11/12	0.01607	0.01616	0.00161	0.01616	0.97238	0.01351	0.01411
1	0.01625	0.01628	0.00162	0.01628	0.96977	0.01479	0.01544
2	0.01860	0.01772	0.00186	0.01772	0.93713	0.03089	0.03198
3	0.02129	0.01929	0.00213	0.01929	0.90200	0.04833	0.04967
4	0.02439	0.02101	0.00244	0.02101	0.86432	0.06712	0.06856
5	0.02794	0.02289	0.00279	0.02289	0.82407	0.08722	0.08872
6	0.03202	0.02493	0.00320	0.02493	0.78127	0.10855	0.11018
7	0.03671	0.02717	0.00367	0.02717	0.73601	0.13100	0.13299
8	0.04209	0.02961	0.00421	0.02961	0.68846	0.15435	0.15719
9	0.04826	0.03227	0.00483	0.03227	0.63886	0.17835	0.18279
10	0.05535	0.03517	0.00554	0.03517	0.58756	0.20263	0.20981

## Additional problem

When you have the moment, try to calculate (using some software or a spreadsheet) to estimate the transition probabilities given that at age 60, the person is sick:  $\underbrace{{}_{10}p_{60}^{10}}$  and  $\underbrace{{}_{10}p_{60}^{11}}$ , and  $\underbrace{{}_{10}p_{60}^{12}}$

Kolmogorov forward equations:

$$\frac{d}{dt} tP_x^{ij} = \underbrace{\sum_{k=0, k \neq j}^n tP_x^{ik} \mu_{x+t}^{kj}}_{\substack{i \rightarrow k \rightarrow j \\ \downarrow \\ k \neq j}} - \sum_{k=0, k \neq j}^n tP_x^{ij} \mu_{x+t}^{jk}$$

$i \rightarrow j \rightarrow k$   
 $k \neq j$

# Illustrative example 1

Consider the health-sickness insurance model with:



$$\mu_{50+t}^{hs} = 0.040,$$

$$\mu_{50+t}^{sh} = 0.005,$$

$$\mu_{50+t}^{hd} = 0.010, \text{ and}$$

$$\mu_{50+t}^{sd} = 0.020,$$

$${}_{10}p_{50}^{\overline{hh}} = .605307$$

$${}_{10}p_{50}^{\overline{ss}} = .7788008$$

for all  $t \geq 0$ . Do the following:

① Calculate  ${}_{10}p_{50}^{\overline{hh}}$  and  ${}_{10}p_{50}^{\overline{ss}}$ .

② Write out the Kolmogorov's forward equations for solving the  $t$ -year transition probabilities for a person age 50 who is currently healthy.

✓ (consider all possible transitions; do not solve)  $\frac{d}{dt} p_{50}^{sh}, \frac{d}{dt} p_{50}^{ss}, \frac{d}{dt} p_{50}^{sd}$

③ Write out the Kolmogorov's forward equations for solving the  $t$ -year transition probabilities for a person age 50 who is currently sick.

✓ (consider all possible transitions; do not solve)



instant  
ann

←  $\bar{A}$

insurance

So long  
as you  
stay in  
a state.

←  $\bar{a}$

annuities.



## Illustrative example 2

Suppose that an insurer uses the health-sickness model to price a policy that provides both sickness and death benefits to healthy lives aged 40.

You are given:


- The term of the policy is 25 years.
- If the individual dies during the term of the policy, there is a death benefit of \$20,000 payable at the moment of death. An additional \$10,000 is payable if the individual is sick at the time of death.
- If the individual becomes sick during the term of the policy, there is a sickness benefit at the rate of \$3,000 per year. No waiting period before benefits are payable.
- The premium rate is \$600 payable annually by healthy policyholders.

Express the following in integral form using standard notation of transition probabilities and forces of transitions:

- 1 the actuarial present value at issue of future premiums;
- 2 the actuarial present value at issue of future death benefits; and
- 3 the actuarial present value at issue of future sickness benefits.


$$APV(FP_0) = \int_0^{25} 600v^t + P_{40}^{hh} dt$$

$$= 600 \int_0^{25} v^t + P_{40}^{hh} dt$$



$$APV(FDB_0) = \int_0^{25} 20000v^t + P_{40}^{hh} \mu_{40:t}^{hd} dt$$


$$+ \int_0^{25} 30000v^t + P_{40}^{hs} \mu_{40:t}^{sd} dt$$




$$APV(FSB_0) = \int_0^{25} 30000v^t + P_{40}^{hs} dt$$


$$\underbrace{\hspace{10em}}_{-hs}$$

$$A_{40:\overline{25}|}$$



$$\bar{A}_{x:\overline{n}|} = \int_0^n \left( v^t p_x^{hh} \mu_{x+t}^{hd} + v^t p_x^{hs} \mu_{x+t}^{sd} \right) dt$$





## Policy values and Thiele's differential equations

Consider the health-sickness insurance model where we have a disability income policy with a term for  $n$  years issued to a healthy life ( $x$ ):

- Premiums are payable continuously throughout the policy term at the rate of  $P$  per year, while healthy.
- Benefit in the form of an annuity is payable continuously at the rate of  $B$  per year, while sick.
- A lump sum benefit of  $S$  is payable immediately upon death within the term of the policy.


Give an expression for the:

- 1 policy value at time  $t$  for a healthy policyholder;
- 2 policy value at time  $t$  for a sick policyholder; and
- 3 Thiele's differential equations for solving these policy values.

$P =$  premium ( $x$ )  $n$  years

$B =$  sickness benefit

$S =$  death benefit




① healthy at time  $t$  <sup>dead</sup>

$${}_tV^{(h)} = \underbrace{APV(FDB_t) + APV(FSB_t)}_{\text{all benefit}} - \underbrace{APV(FP_t)}_{\text{all premiums}}$$

$$= S \times \underbrace{\bar{A}_{x+t:\overline{n-t}|}^{hd}}_{\text{try this in integral form}} + B \underbrace{\bar{a}_{x+t:\overline{n-t}|}^{hs}}_{\int_0^{n-t} v^r r p_{x+t}^{hs} dr} - P \times \underbrace{\bar{a}_{x+t:\overline{n-t}|}^{hh}}_{\int_0^{n-t} v^r r p_{x+t}^{hh} dr}$$

try this in integral form

leave for you to do





# Generalization of Thiele's differential equations

- Section 8.7.2, pages 266-267
- General situation of an insurance contract issued within a more general multiple state model

## SOA question #10, Spring 2018


A whole life policy with Critical Illness benefits is issued to a Healthy life age 60. The insurer uses the following multiple state model to value the benefits:



The premium  $\dot{P}$  is payable continuously while the policyholder is Healthy. A benefit of 50,000 is paid immediately on diagnosis of Critical Illness (CI), with another 50,000 paid on death after CI. If the policyholder dies without a diagnosis, the full 100,000 is paid immediately on death.



- continued



03 → 1 <sup>at sum</sup> ~~at sum~~

You are given the following information:

$x$	$\bar{a}_x^{00}$	$\bar{A}_x^{01}$	$\bar{A}_x^{02}$	$\bar{A}_x^{03}$	$\bar{A}_x^{13}$
60	10.989	0.390	0.181	0.280	0.546

Calculate  $P$ .

$$\begin{aligned}
 \text{APV}(FP_0) &= \text{APV}(FDB_0) + \text{APV}(FCI) \\
 P \bar{a}_{60}^{00} &= 50000 \bar{A}_{60}^{02} + 50000 \bar{A}_{60}^{01}
 \end{aligned}$$

$$P = 4695.605$$

## SOA question #12, Spring 2012

Employees in Company ABC can be in: **State 0**: Non-executive employee; **State 1**: Executive employee; or **State 2**: Terminated from employment.

John joins Company ABC as a non-executive employee at age 30.


You are given:

- $\mu^{01} = 0.01$  for all years of service
- $\mu^{02} = 0.006$  for all years of service
- $\mu^{12} = 0.002$  for all years of service
- Executive employees never return to the non-executive employee state.
- Employees terminated from employment never get rehired.
- The probability that John lives to age 65 is 0.9, regardless of state.

Calculate the probability that John will be an executive employee of Company ABC at age 65.

## SOA question #10, Fall 2013

For a multiple state model, you are given:



The following forces of transition:

$$\mu^{01} = 0.02 \quad \mu^{02} = 0.03 \quad \mu^{12} = 0.05$$

Calculate the conditional probability that a Healthy life on January 1, 2004 is still Healthy on January 1, 2014, given that this person is not Dead on January 1, 2014.

## SOA question #3, Fall 2015

Johnny Vegas performs motorcycle jumps throughout the year and has injuries in the course of his shows according to the following three-state model:

State 0: No injuries

State 1: Exactly one injury

State 2: At least two injuries

You are given:

- Transition intensities between States are per year.
- $\mu_t^{01} = 0.03 + 0.06 \times 2^t$ , for  $t > 0$
- $\mu_t^{02} = 2.718 \mu_t^{01}$ , for  $t > 0$
- $\mu_t^{12} = 0.025$ , for  $t > 0$

Calculate the probability that Johnny, who currently has no injuries, will sustain at least one injury in the next year.

annual  
monthly  
semi annually

## Discrete time Markov chain models

---

# Transition probabilities - Markov Chains

- Assume a finite state space:  $\{0, 1, 2, \dots, n\}$  and let  $Y_x(k)$  be the state at time  $k$ .
- Basic **Markov chain** assumption:

$$\begin{aligned} \Pr[Y_x(k+1) = j | Y_x(k) = i, Y_x(k-1), \dots, Y_x(0)] \\ = \Pr[Y_x(k+1) = j | Y_x(k) = i] \end{aligned}$$

- Notation of transition probabilities:

$$\Pr[Y_x(k+1) = j | Y_x(k) = i] = Q_k^{(i,j)} = Q_k^{ij}.$$

- Transition probability matrix:

*Q* homogeneous  $\leftarrow$   $Q_k$   $\leftarrow$  *total*

$$Q_k = \begin{matrix} & \begin{matrix} 0 & 1 & \dots & n \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ \vdots \\ n \end{matrix} & \begin{pmatrix} Q_k^{00} & Q_k^{01} & \dots & Q_k^{0,n} \\ Q_k^{10} & Q_k^{11} & \dots & Q_k^{1,n} \\ \vdots & \vdots & \ddots & \vdots \\ Q_k^{n,0} & Q_k^{n,1} & \dots & Q_k^{n,n} \end{pmatrix} \end{matrix}$$

# Homogeneous and non-homogeneous Markov chains

- If the transition probability matrix  $Q_k$  depends on the time  $k$ , it is said to be a **non-homogeneous** Markov Chain.
- Otherwise, it is called a **homogeneous** Markov Chain, and we shall simply denote the transition probability matrix by  $Q$ .
- Define

$${}_r Q_k = \begin{matrix} & \begin{matrix} 0 & 1 & \dots & n \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ \vdots \\ n \end{matrix} & \begin{pmatrix} r Q_k^{00} & r Q_k^{01} & \dots & r Q_k^{0,n} \\ r Q_k^{10} & r Q_k^{11} & \dots & r Q_k^{1,n} \\ \vdots & \vdots & \ddots & \vdots \\ r Q_k^{n,0} & r Q_k^{n,1} & \dots & r Q_k^{n,n} \end{pmatrix} \end{matrix} = Q_k * Q_{k+1} * \dots * Q_{k+r-1}$$

where

$${}_r Q_k^{ij} = \Pr[Y_x(k+r) = j | Y_x(k) = i]$$

is the probability of going from state  $i$  to state  $j$  in  $r$  steps. It is sometimes written as  ${}_r Q_k^{(i,j)}$ .

# Chapman-Kolmogorov equations

- Discrete analogue of the Kolmogorov's forward equations.
- Theorem:

$${}_r Q_k = Q_k \times Q_{k+1} \times \cdots \times Q_{k+r-1}$$

- Chapman-Kolmogorov equations:

$${}_{m+p} Q_k^{ij} = \sum_s {}_m Q_k^{is} \times {}_p Q_{k+m}^{sj}$$

- In the case of homogeneous Markov Chains, we drop the subscript  $k$  and simply write

$${}_r Q = Q \times \cdots \times Q = \underbrace{Q^r}$$





## Example 1

- Consider a critical illness model with 3 states: healthy (H), critically ill (C) and dead (D).
- Suppose you have the homogeneous Markov Chain with transition matrix

$$\begin{array}{c}
 \text{H} \quad \text{C} \quad \text{D} \\
 \text{H} \begin{pmatrix} 0.92 & 0.05 & 0.03 \\ 0.00 & 0.76 & 0.24 \\ 0.00 & 0.00 & 1.00 \end{pmatrix} \\
 \text{C} \\
 \text{D}
 \end{array}$$

- What are the probabilities of being in each of the state at times  $t = 1, 2, 3$ ?

$$P_0 = P = \begin{pmatrix} .92 & .05 & .03 \\ 0 & .76 & .24 \\ 0 & 0 & 1 \end{pmatrix}$$



t=2 2 periods

Chapman  
Kilmer

$${}^2Q = Q * Q = \begin{pmatrix} .92 & .05 & .03 \\ 0 & .76 & .24 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} .92 & .05 & .03 \\ 0 & .76 & .24 \\ 0 & 0 & 1 \end{pmatrix}$$
$$= \begin{pmatrix} .8464 & .0840 & .0696 \\ 0 & .5776 & .4224 \\ 0 & 0 & 1 \end{pmatrix}$$

prob that C person will be D in two years = 0.4224

$${}^3Q = \begin{pmatrix} .778688 & .106160 & .115152 \\ 0 & .438976 & .56104 \\ 0 & 0 & 1 \end{pmatrix}$$

0.438976

0.438976

$${}_3Q^{CD} = \text{Pr. ob}(C \text{ at } t=0 | D \text{ at } t=3)$$

	C		D	
	0	1	2	3
C		D	→	.24
C	←	C		D
			→	.76 × .24
C		C		C
				D
				.76 × .76
				× .24
				<u>Σ = .56104</u>

Now suppose we have 10 policyholders now at time 0  
 Find the probability that exactly 4 of them will be critically ill  
 at the end of 3 years.

$$\text{Pr}(N=4) = \binom{10}{4} (.106160)^4 (1 - .106160)^6$$

$$= .2878518$$

$${}_3Q^{CC} = .106160$$

$N = \text{number} \sim 1$

$$E(N) = nq = 10(.106160)$$

$$\text{Var}(N) = nq(1-q) = 10(.106160)(1 - .106160)$$

now suppose 1000 polyhydrazes

probability that at least 120 will be critically ill in 3 years

$$Pr(N \geq 120) = \binom{1000}{120} (.106160)^{120} (1 - .106160)^{880} + \dots$$

$N = \text{number}$   
 $\sim \text{binomial } (n=1000, p=.106160)$

Recall  $N$  can be approximately Normal

with mean  $E(N) = 1000(.106160) = 106.16$

$\text{Var}(N) = 1000(.106160)(1 - .106160) = 94.89005$

$$Pr[N \geq 120] = Pr\left[ \frac{N - E(N)}{\sqrt{\text{Var}(N)}} \geq \frac{120 - 106.16}{\sqrt{94.89005}} \right] = Pr[Z \geq a] = .0777$$

$$Pr[N \geq 120] = Pr[N \geq 119.5] = Pr\left[ Z \geq \frac{119.5 - 106.16}{\sqrt{94.89005}} \right] = .0854$$

ccf

Exact Value .08713564



## Example 2

- Suppose that an auto insurer classifies its policyholders according to **Preferred** (State #0) or **Standard** (State #1) status, starting at time 0 at the start of the first year when they are first insured, with reclassifications occurring at the start of each new policy year.
- You are given the following  $t$ -th year non-homogeneous transition matrix:

$$Q_t = \begin{pmatrix} 0.65 & 0.35 \\ 0.50 & 0.50 \end{pmatrix} + \frac{1}{t+1} \begin{pmatrix} 0.15 & -0.15 \\ -0.20 & 0.20 \end{pmatrix}$$

- Given that an insured is Preferred at the start of the second year:
  - 1 Find the probability that the insured is also Preferred at the start of the third year.
  - 2 Find the probability that the insured transitions from being Preferred at the start of the third year to being Standard at the start of the fourth year.

## Cash flows and actuarial present values

- We are interested in the actuarial present value of cash flows

$${}_{t+k+1}C^{ij}$$

which are the cash flows at time  $t + k + 1$  for movement from state  $i$  (at time  $t + k$ ) to state  $j$  (at time  $t + k + 1$ ).

- Discount typically by  $v^{k+1}$ .
- Theorem: Suppose that the subject is in state  $s$  at time  $t$ . The **actuarial present value** (APV) of cash flows from state  $i$  to state  $j$  is given by

$$\text{APV}_{s@t} = \sum_{k=0}^{\infty} \left( {}_kQ_t^{si} \cdot Q_{t+k}^{ij} \right) {}_{t+k+1}C^{ij} \times v^{k+1}.$$

time homogeneous

$$Q \begin{matrix} & H & C & D \\ H & .92 & .05 & .03 \\ C & 0 & .76 & .24 \\ D & 0 & 0 & 1 \end{matrix}$$

$Q^{-1}$        $Q^2$

You are healthy now

• benefit is 100 each time you are critically ill

• no other benefits

•  $i = 5\%$

• 3 year term

discount  
cash flow

$$100(v + v^2 + v^3) = 7.86474$$

$$100(v^2 + v^3) = 6.190551$$

$$100v^3 = 3.655701$$

$$100v = 1.142857$$

$$100v^2 = 1.095782$$

$$100(v + v^2) = 1.001361$$

$$\Sigma = 21.55$$

possible  
transitions

H → C → C

H → H → C → C

H → H → H → C

H → C → D → D

H → H → C → D

H → C → C → D

probabilities

$$.05 \times .76 \times .76$$


$$.92 \times .05 \times .76$$

$$.92^2 \times .05$$

$$.05 \times .24$$

$$.92 \times .05 \times .24$$

$$.05 \times .76 \times .24$$



# Another approach

## transitions

in C 1st year

$$1Q^{HC} = .05$$

$$\times \frac{\text{discounted CF}}{100 v} = 4.761905$$

in C 2nd year

$$2Q^{HC} = .089$$

$$\times 100 v^2 = 7.619048$$

in C 3rd year

$$3Q^{HC} = .10616$$

$$\times 100 v^3 = 7.170500$$

---


$$\Sigma = 21.5595$$



---

$$2Q^{HC} \left( \begin{array}{l} H \rightarrow H \rightarrow C \quad .92 \times .05 \\ H \rightarrow C \rightarrow C \quad .15 \times .76 \end{array} \right)$$

$$3Q^{HC} \left( \begin{array}{l} H \rightarrow H \rightarrow H \rightarrow C \\ H \rightarrow H \rightarrow C \rightarrow C \\ H \rightarrow C \rightarrow C \rightarrow C \end{array} \right) + = .10616$$

quiz  
in  
Velocity





## Illustrative example no. 1

An insurer issues a special 3-year insurance contract to a high risk individual with the following homogeneous Markov Chain model:

- States: 0 = active, 1 = disabled, 2 = withdrawn, and 3 = dead.
- Transition probability matrix:

$$\begin{array}{c} \rightarrow 1 \\ 2 \\ 3 \end{array} \begin{pmatrix} 0 & 1 & 2 & \textcircled{3} \\ 0.4 & 0.2 & 0.3 & 0.1 \\ 0.2 & 0.5 & 0.0 & 0.3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$i=5\%$

- Changes in state occur only at the end of the year.
- The death benefit is \$1,000 payable at the end of the year of death.
- The insured is disabled at the end of year 1.
- Assuming interest rate of 5% p.a., Calculate the actuarial present value of the prospective death benefits at the beginning of year 2.

Death  
in  
2nd  
yr

Transitions

① → ③

① → ① → ③

① → ② → ③

Prob

.30 \*

.5(.3) = .15 } -17

.2(.1) = .02

Benefit

1000 v

1000

Discout


v

v<sup>2</sup>

Present

285.7143

154.1950



+ = 439.9093

APV  $\approx$  439.91  
future  
lump benefit

3rd  
yr

## Illustrative example no. 2

Consider a special three-year term insurance:

- Insureds may be in one of three states at the beginning of each year: active, disabled or dead. All insureds are initially active.
- The annual transition probabilities are as follows:

	0	1	2	
0	Active	0.8	0.1	0.1
1	Disabled	0.1	0.7	0.2
2	Dead	0.0	0.0	1.0

- A \$100,000 benefit is payable at the end of the year of death whether the insured was active or disabled.
- Premiums are paid at the beginning of each year when active. Insureds do not pay annual premiums when they are disabled.
- Interest rate  $i = 10\%$ .

Calculate the level annual net premium for this insurance.

$P = \text{premium}$



$$APV(\text{premiums}) = P + PV(.8) + PV^2(.8^2 + .1(1))$$

$0 \rightarrow 0$                        $0 \rightarrow 0 \rightarrow 0$   
 $0 \rightarrow 1 \rightarrow 0$   
 $\underbrace{\quad \quad}_{.1}$     $\underbrace{\quad \quad}_{.1}$

$$= P(1 + .8v + .65v^2)$$

$$APV(\text{Death}) = 100,000 \left( v(.1) + v^2(.8(1) + .1(.2)) \right. \\ \left. + v^3(.064 + .016 + .014 + .001) \right)$$

Equate the two,  $\boxed{P = 10,816.19}$



## Illustrative example no. 3



- A machine can be in one of four possible states, labeled  $a$ ,  $b$ ,  $c$ , and  $d$ . It migrates annually according to a Markov Chain with transition probabilities:

$$\begin{array}{c} \rightarrow a \\ b \\ c \\ d \end{array} \begin{array}{c} \checkmark \\ \\ \\ \end{array} \begin{array}{c} a \\ b \\ c \\ d \end{array} \begin{pmatrix} 0.25 & 0.75 & 0.00 & 0.00 \\ 0.50 & 0.00 & 0.50 & 0.00 \\ 0.80 & 0.00 & 0.00 & 0.20 \\ 1.00 & 0.00 & 0.00 & 0.00 \end{pmatrix}$$

500 V<sup>2</sup> $i = 5\%$ 

- At time  $t = 0$ , the machine is in State  $a$ . A salvage company will pay 500 at the end of 2 years if the machine is in State  $a$ .
- Assuming  $i = 0.05$ , calculate the actuarial present value at time  $t = 0$  of this payment.

Illustration no. 3

$$a \rightarrow a \rightarrow a \quad .25 \times .25 =$$

$$a \rightarrow b \rightarrow c \quad .75 \times .50 =$$

$$a \rightarrow c \rightarrow a \quad 0$$

$$c \rightarrow d \rightarrow c \quad 0$$



---

$$\times 500 \text{ V}^2 = \underline{\underline{198.4127}}$$

a 0 0 0

Suppose payments are 100 in a

200 in b


300 in c

400 in d

state vector \* Q \* Q

(1000) \* Q \* Q

$$(1000) \begin{pmatrix} .25 & .75 & 0 & 0 \\ .5 & & & \\ .8 & & & \\ .1 & & & \end{pmatrix} \begin{pmatrix} .25 & .75 & 0 & 0 \\ .5 & & & \\ .8 & & & \\ .1 & & & \end{pmatrix}$$




$$\begin{pmatrix} .25 & .75 & 0 & 0 \\ .5 & & .5 & 0 \\ .8 & & 0 & .2 \\ 1 & & 0 & 0 \end{pmatrix} = \begin{pmatrix} .25(.25) & .25(.75) & 0 & 0 \\ +.75(.5) & +0 & .75(.5) & 0 \end{pmatrix}$$

$$APV = .4375(100v) + .1875(200v^2) + .375(300v^3) + 0 = 193.75$$

$$a(.4375, .1875, .375, 0)$$

## SOA essay question #1, Fall 2016

You are given the following 3-state Markov model:



For all states  $i$  and  $j$ , and for all ages  $x \geq 0$ ,  ${}_t p_x^{ij}$  is a differentiable function of  $t$ , and for  $i \neq j$ :

$$\mu_x^{ij} = \lim_{h \rightarrow 0^+} \frac{1}{h} h p_x^{ij},$$

- (a) Define the symbols  ${}_t p_x^{00}$  and  $\overline{{}_t p_x^{00}}$ , and explain why these probabilities are not equal for this model.



## - continued

(b) The probability  ${}_{t+h}p_x^{00}$  can be expressed as  
 ${}_{t+h}p_x^{00} = {}_t p_x^{00} {}_h p_{x+t}^{00} + {}_t p_x^{01} {}_h p_{x+t}^{10}$ . Use this equation to derive the  
 Kolmogorov forward differential equation for  ${}_t p_x^{00}$ .

(c) You are also given:

(i)  $\mu_{x+t}^{01} = 0.5$ , for all  $t$


(ii)  $\mu_{x+t}^{02} = kt$ , for all  $t$

(iii)  ${}_2 p_x^{00} = 0.165$


Calculate  $k$ .

## Other transition models with actuarial applications


## Joint life model



## Multiple decrement model



## Accidental death model



## A simple retirement model

