
9.1 Modeling with Differential Equations

Differential equations. An equation containing an unknown function and some of its derivatives is a *differential equation*. Examples are $\frac{dy}{dx} = 4x$ and $y''(x) + x^3y'(x) = xy(x)$.

The order of a differential equation is the order of the highest-order derivative in the equation.

For example, $\frac{dy}{dx} = 4x$ is a first-order differential equation while $y''(x) + x^3y'(x) = xy(x)$ is a second-order differential equation.

Example: Which of the functions below satisfy the differential equation $y'' + y = \sin x$?

- (a) $y = \sin x$
- (b) $y = \cos x$
- (c) $y = \frac{1}{2}x \sin x$
- (d) $y = -\frac{1}{2}x \cos x$

Thinking about the problem:

First we will find y' and then y'' for each of the functions and then compute $y'' + y$ in each case to see if we get $\sin x$.

Doing the problem:

- (a) $y = \sin x \implies y'' = -\sin x \implies y'' + y = 0,$
- (b) $y = \cos x \implies y'' = -\cos x \implies y'' + y = 0,$
- (c) $y = \frac{1}{2}x \sin x \implies y'' = \cos x - \frac{1}{2}x \sin x \implies y'' + y = \cos x,$
- (d) $y = -\frac{1}{2}x \cos x \implies y'' = \sin x + \frac{1}{2}x \cos x \implies y'' + y = \sin x.$

The only solution to $y'' + y = \sin x$ among the four functions here is (d) $y = -\frac{1}{2}x \cos x$.

Solutions should show all of your work, not just a single final answer.

1. We consider the differential equation $\frac{dy}{dt} = 1 - 2y$. (Here the independent variable is t .)

(a) Find all constant solutions. That is, if $y = K$ for constant K satisfies the differential equation, what does K need to be?

(b) Show every function of the form $y(t) = \frac{1}{2} + Ce^{-2t}$, where C is a constant, is a solution of the differential equation.

(c) If $y(t)$ is a function described by part (b), what can you say about the long-term behavior $\lim_{t \rightarrow \infty} y(t)$?

2. We consider the differential equation $\frac{dy}{dx} = xy$. (Here the independent variable is x .)

(a) Find all constant solutions.

(b) Show every function of the form $y(x) = Ce^{x^2/2}$, where C is a constant, is a solution.

(c) For a solution as in part (b), describe C as a value of $y(x)$.

3. T/F (with justification)

Every differential equation has a constant solution.