




Recall constructing vectors to denote moving between 2 points:





$$\vec{u} \cdot \vec{v} = |\vec{u}| |\vec{v}| \cos \theta$$
$$\theta = \cos^{-1} \left( \frac{\vec{u} \cdot \vec{v}}{|\vec{u}| |\vec{v}|} \right)$$



$$\text{proj}_{\vec{v}} \vec{u} = \left( \frac{\vec{u} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \right) \vec{v}$$

Ex: Let  $A = (2, -1, 1)$ ,  $B = (3, 1, 1)$ ,  
 $C = (-1, 0, 4)$ . Find  $\angle ABC$ .



Use vectors  $\vec{BA}$ ,  $\vec{BC}$

$$\vec{BA} = \langle 2-3, -1-1, 1-1 \rangle = \langle -1, -2, 0 \rangle$$

$$\vec{BC} = \langle -1-3, 0-1, 4-1 \rangle = \langle -4, -1, 3 \rangle$$

$$\theta = \cos^{-1} \left( \frac{\vec{BA} \cdot \vec{BC}}{|\vec{BA}| |\vec{BC}|} \right) = \cos^{-1} \left( \frac{4+2}{\sqrt{5} \cdot \sqrt{26}} \right) = \cos^{-1} \left( \frac{6}{\sqrt{130}} \right)$$


Ex: If  $\vec{v} = 2\vec{i} - \vec{j} + 6\vec{k}$ ,  $\vec{u} = \vec{j} - \vec{k}$ ,  
what is  $\text{proj}_{\vec{v}} \vec{u}$ ?

$$\text{proj}_{\vec{v}} \vec{u} = \left( \frac{\vec{u} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \right) \vec{v} = \frac{0 - 1 - 6}{4 + 1 + 36} \vec{v} = \frac{-7}{41} \vec{v}$$

in the  
direction of  $\vec{v}$ ,  
or "onto"  $\vec{v}$

$$= \frac{-14}{41} \vec{i} + \frac{7}{41} \vec{j} - \frac{42}{41} \vec{k}.$$

# LINES



Red line can be created by adding multiples of  $\vec{v}$  to  $\vec{r}_0$  ...

$\langle x, y, z \rangle = \vec{r}_0 + t\vec{v}$  } vector equation for the line

$$\vec{r}_0 = \langle x_0, y_0, z_0 \rangle$$

$$\vec{v} = \langle v_1, v_2, v_3 \rangle$$

Then

$$\langle x, y, z \rangle = \langle x_0 + t v_1, y_0 + t v_2, z_0 + t v_3 \rangle$$

"Parametric equations"

$$x = x_0 + t v_1$$

$$y = y_0 + t v_2$$

$$z = z_0 + t v_3.$$

"Symmetric" equations...


$$\text{Solve for } t: x = x_0 + t v_1 \Rightarrow t = \frac{x - x_0}{v_1}$$

$$\text{Also, } t = \frac{y - y_0}{v_2} = \frac{z - z_0}{v_3} = \frac{x - x_0}{v_1}$$

symmetric equations

If, e.g.  $v_3 = 0$ , then  $z = z_0 + t \cancel{v_3} = z_0$

$$\Rightarrow \frac{x - x_0}{v_1} = \frac{y - y_0}{v_2} \quad \& \quad z = z_0$$



$$\vec{n} = \langle n_1, n_2, n_3 \rangle$$

$$\vec{v} = \langle x - x_0, y - y_0, z - z_0 \rangle$$

$$\vec{n} \cdot \vec{v} = n_1(x - x_0) + n_2(y - y_0) + n_3(z - z_0) = 0$$

Scalar equation for plane



Note how we can algebraically  
manipulate the equation of the plane...

$$n_1x + n_2y + n_3z - n_1x_0 - n_2y_0 - n_3z_0 = 0.$$

→ looks like  $ax + by + cz + d = 0$ .

So if you see this, then  
you know  $\vec{n} = \langle a, b, c \rangle$ . 😊

Ex: One line is given by  $\frac{x-1}{4} = \frac{z+2}{3}, y = -2$  and another by  $\vec{r}(t) = \langle t+1, 2t-2, t-2 \rangle$ . They lie in a common plane; find the equation of the plane.

Direction vectors are  $\vec{v}_1 = \langle 4, 0, 3 \rangle$

$$\vec{v}_2 = \langle 1, 2, 1 \rangle$$

$$\Rightarrow \vec{n} = \vec{v}_1 \times \vec{v}_2 = \langle 0 \cdot 1 - 3 \cdot 2, 1 \cdot 3 - 4 \cdot 1, 4 \cdot 2 - 0 \cdot 1 \rangle = \langle -6, -1, 8 \rangle$$

We need a point  $(x_0, y_0, z_0)$  in the plane...  
any point on either line works, e.g.  $(1, -2, -2)$

$$\text{Plane equation: } \vec{n} \cdot \langle x - x_0, y - y_0, z - z_0 \rangle = 0$$

$$\Rightarrow \langle -6, -1, 8 \rangle \cdot \langle x - 1, y + 2, z + 2 \rangle = 0$$

$$\Rightarrow -6(x - 1) - (y + 2) + 8(z + 2) = 0.$$

Ex: Find the equation (vector form) of the line passing through  $(0, -1, 1)$  perpendicular to the plane  $2y - z + 5x - 1 = 0$ .

$\vec{n} = \langle 5, 2, -1 \rangle$ : direction of line

$$\begin{aligned}\vec{r}(t) &= \langle 0, -1, 1 \rangle + t \langle 5, 2, -1 \rangle \\ &= \langle 5t, 2t - 1, 1 - t \rangle.\end{aligned}$$

## Review of other surfaces

(a) one-sheet  
hyperboloid

(b) elliptic cylinder

(c) ellipsoid

(d) hyperbolic  
paraboloid


(e) two-sheet hyperboloid

(f) elliptic paraboloid


(g) cone

(h) non-elliptic cylinder


$$(a) \quad \frac{x^2}{3} - \frac{y^2}{9} + \frac{z^2}{9} = 1$$



(b)  $\frac{y^2}{4} + z^2 = 1$



$$(c) \left(\frac{x}{8}\right)^2 + \left(\frac{y}{4}\right)^2 + \left(\frac{z}{2}\right)^2 = 1$$







(d)

$$x = y^2 - \frac{z^2}{9}$$


Hyperbolic  
paraboloid




$$(e) \left(\frac{z}{1/3}\right)^2 - \left(\frac{y}{1/2}\right)^2 - x^2 = 1$$




$$(f) \quad y = \frac{x^2}{4} + \frac{z^2}{8}$$



$$(9) \left( \frac{y}{1/\sqrt{5}} \right)^2 = x^2 + \left( \frac{z}{1/\sqrt{10}} \right)^2$$



$$(h) \quad z = x^2 + 1$$



parabolic  
(non-elliptic)

cylinder

## Intersections of surfaces


For example,  $z = 1 + y^2 + \frac{1}{2}x^2$

&  $y = 3x + 5$

Look for an easy way to get 2 variables in terms of a third...

$$\left. \begin{array}{l} z = 1 + (3x + 5)^2 + \frac{1}{2}x^2 \\ y = 3x + 5 \end{array} \right\} \begin{array}{l} \text{Let "x"} \\ \text{be the} \\ \text{parameter} \end{array}$$

# Arc length formula



$$L = \int_a^b |\vec{r}'(t)| dt$$


Ex: Let a curve have parameterization  
 $\vec{r}(t) = \left\langle t, \frac{2\sqrt{2}}{3} t^{3/2}, \frac{1}{2} t^2 \right\rangle, 0 \leq t \leq 2.$

Find the length of the arc segment.

$$\begin{aligned} \int_0^2 \vec{r}'(t) &= \left\langle 1, \frac{2\sqrt{2}}{3} \cdot \frac{3}{2} \sqrt{t}, t \right\rangle = \langle 1, \sqrt{2t}, t \rangle. \\ \int_0^2 |\vec{r}'(t)| dt &= \int_0^2 \sqrt{1 + 2t + t^2} dt = \int_0^2 \sqrt{(1+t)^2} dt \\ &= \int_0^2 (1+t) dt = \left. 2 + \frac{1}{2} t^2 \right|_0^2 = 4. \end{aligned}$$



## The normal and binormal vectors



How can we find three orthogonal unit vectors that move with a particle through space?

$\vec{T}$  points in the direction of motion, so that will be the first vector. Note

$$\text{that } |\vec{T}| = 1 \Rightarrow |\vec{T}|^2 = \vec{T} \cdot \vec{T} = 1$$

$$\Rightarrow \frac{d}{dt}(\vec{T} \cdot \vec{T}) = \frac{d}{dt}(1) = 0$$

$$\Rightarrow 2\vec{T} \cdot \vec{T}' = 0$$

$$\Rightarrow \vec{T} \perp \vec{T}'$$

so normalize  
this...  
 $\vec{N} = \frac{\vec{T}'}{|\vec{T}'|}$

$\vec{N} = \frac{\vec{T}'}{|\vec{T}'|}$  : NORMAL vector  
for a curve (not a plane).

Bi-normal :  $\vec{B} = \vec{T} \times \vec{N}$

$$|\vec{B}| = |\vec{T}| |\vec{N}| \sin(90^\circ) = 1 \cdot 1 \cdot 1 = 1.$$

$\{\vec{T}, \vec{N}, \vec{B}\}$  : the "TNB frame".

Ex: Find the TNB-frame vectors  
for  $\vec{r}(t) = \langle \sqrt{2} \cos(t), 4t, \sqrt{2} \sin(t) \rangle$   
at the point  $(0, 2\pi, \sqrt{2})$ .

$$\underbrace{\hspace{10em}}_{\text{so } t = \pi/2}$$

$$\vec{r}'(t) = \langle -\sqrt{2} \sin(t), 4, \sqrt{2} \cos(t) \rangle$$

$$|\vec{r}'(t)| = \sqrt{2(\sin^2(t) + \cos^2(t)) + 16} = \sqrt{18} = 3\sqrt{2}$$

$$\vec{T} = \frac{\vec{r}'}{|\vec{r}'|} = \left\langle -\frac{1}{3} \sin(t), \frac{2\sqrt{2}}{3}, \frac{1}{3} \cos(t) \right\rangle$$

$$\vec{T}' = \left\langle -\frac{1}{3} \cos(t), 0, -\frac{1}{3} \sin(t) \right\rangle$$

$$\vec{N} = \left\langle -\cos(t), 0, -\sin(t) \right\rangle$$

$$\vec{T}\left(\frac{\pi}{2}\right) = \left\langle -\frac{1}{3}, \frac{2\sqrt{2}}{3}, 0 \right\rangle$$

$$\vec{N}\left(\frac{\pi}{2}\right) = \left\langle 0, 0, -1 \right\rangle$$

$$\vec{B} = \left\langle -\frac{2\sqrt{2}}{3}, -\frac{1}{3}, 0 \right\rangle.$$

## Velocity and acceleration

$\vec{r}(t)$  : position at time  $t$ .

$\vec{v}(t) = \vec{r}'(t)$  : velocity at time  $t$ .

$$\vec{v} = |\vec{v}| \hat{v}$$

speed  $\uparrow$   $\uparrow$  direction

$\vec{a}(t) = \vec{v}'(t)$   
 $= \vec{r}''(t)$  : acceleration

Given acceleration, we can also find velocity and position by integrating...

$$\vec{v} = \int \vec{a}(t) dt$$

$$\vec{r} = \int \vec{v}(t) dt$$

We require unknown constants to be added at each step; their values are often determined later from "initial conditions".

## Newton's Second Law

$$\vec{F} = m \vec{a}$$

*force*      *mass*

Often we just want the acceleration:

$$\vec{a} = \frac{1}{m} \vec{F}.$$



Ex: A particle with mass 4 kg is initially moving with velocity  $\langle 1/2, 1, 1/2 \rangle$  m/s and is acted on by a force  $\vec{F}(t) = \langle t, t^2, 1 \rangle$  N, until 2 seconds later. Find the distance the particle has moved during those 2 seconds.

$$\vec{F} = m\vec{a}$$
$$\vec{a} = \frac{1}{m}\vec{F} = \frac{1}{4}\langle t, t^2, 1 \rangle$$

$$\int \vec{a} = \vec{v} = \frac{1}{4} \left\langle \frac{1}{2}t^2, \frac{1}{3}t^3, t \right\rangle + \vec{C}_1$$

$$\vec{v}(0) = \vec{C}_1 = \left\langle \frac{1}{2}, 1, \frac{1}{2} \right\rangle$$

$$\vec{r} = \int \vec{v} = \frac{1}{4} \left\langle \frac{1}{6}t^3, \frac{1}{12}t^4, \frac{1}{2}t^2 \right\rangle + t \left\langle \frac{1}{2}, 1, \frac{1}{2} \right\rangle$$

$$d = \left| \vec{r}(2) - \vec{r}(0) \right|$$

$$d = \left| \frac{1}{4} \left\langle \frac{8}{6}, \frac{16}{12}, \frac{4}{2} \right\rangle + \langle 1, 2, 1 \rangle \right|$$

$$= \left| \left\langle \frac{1}{3} + 1, \frac{1}{3} + 2, \frac{1}{2} + 1 \right\rangle \right|$$

$$= \left| \left\langle \frac{4}{3}, \frac{7}{3}, \frac{3}{2} \right\rangle \right|$$

$$d = \frac{\sqrt{341}}{6}.$$