

Functions of multiple variables


$$f = f(x_1, x_2, x_3, \dots, x_n)$$

with $f(\cdot)$ a real number

We say that f is a real, scalar-valued function of n variables;


$$f : \mathbb{R}^n \rightarrow \mathbb{R}.$$

Consider $f(x, y) = x + y$; $f: \mathbb{R}^2 \rightarrow \mathbb{R}$.
We visualize sometimes via a
"surface plot" ... set $z = f(x, y)$.



$$f : \mathcal{D} \subset \mathbb{R}^2 \rightarrow \mathbb{R}$$


↑
we often need to restrict the domain



Usually, the domain may be determined by inspection.

$$f(x,y) = \ln(x^2 + y^2 - 1)$$


$$D = \{(x,y) \mid x^2 + y^2 > 1\}$$




EX: Find the domain of

$$f(x,y) = \sqrt{2-x^2-y^2}.$$


$$\text{Then } 2-x^2-y^2 \geq 0 \Rightarrow x^2+y^2 \leq 2$$




We have seen before that traces may be used to help sketch surface plots.



Level curves are another visualization tool.




} Project traces $z = k$ down into x - y plane.




} Visualize just in the x - y plane

Now add in more k-values...



Ex: Sketch a contour plot for $f(x, y) = x^2 - y^2$.



Level surfaces

Consider $f(x, y, z) = k$

e.g. $f(x, y, z) = x^2 + y^2 + z^2 = k > 0$

Spheres of radius \sqrt{k}

One could try to visualize these together,
but this is difficult (how do you see one sphere
within the next?) so we will not focus on level
surfaces.

LIMITS If $f(x,y) \rightarrow L$ as $(x,y) \rightarrow (a,b)$
then we say

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = L.$$


More precisely, if for ANY $\epsilon > 0$ there
is some $\delta = \delta(\epsilon) > 0$ such that

$$0 < \sqrt{(x-a)^2 + (y-b)^2} < \delta \Rightarrow |f(x,y) - L| < \epsilon$$


then the limit exists.

Graphically :

domain



$f(x, y)$




• If we choose $\epsilon > 0$ very small there must be a neighborhood of (a, b) that f maps into $(L - \epsilon, L + \epsilon)$.


1D reminder:

$$\lim_{x \rightarrow a} f(x) = ?$$

(Does not exist)



In 2D (or higher) the limit may only exist along specific paths.



EX: Consider $f(x,y) = \frac{\sin(x^2+y^2)}{x^2+y^2}$.

Note $(0,0)$ is NOT in the domain, but we can still talk about $\lim_{(x,y) \rightarrow (0,0)} f(x,y) = 1$.

(Same as $\lim_{r \rightarrow 0^+} \frac{\sin(r)}{r} = 1$)

EX: Find $\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2+y^2}$ or write DNE
if the limit does not exist.

The way you usually show DNE is to find two paths that yield different limits.

1) Set $y=0 \dots \lim_{\substack{(x,0) \rightarrow (0,0) \\ (x \neq 0)}} \frac{x \cdot 0}{x^2} = 0$

2) Set $y=-x \dots \lim_{\substack{(x,-x) \rightarrow (0,0) \\ (x \neq 0)}} \frac{x(-x)}{2x^2} = \underline{\underline{-\frac{1}{2} \neq 0}}$

To show that limits exist we will usually cheat a bit and cite CONTINUITY.

When $\lim_{(x,y) \rightarrow (a,b)} f(x,y) = f(a,b)$ } • limit exists
• equals $f(a,b)$

We say f is continuous at (a,b) .
(CTS)

* If f is CTS. at every point of a set A

We say " f is CTS on A ".

Polynomials are CTS everywhere and rational functions are CTS on their domains.

EX: Find $\lim_{(x,y) \rightarrow (1,1)} \frac{x^2 - y^2}{x^2 + y^2}$ or write

DNE if the limit does not exist.

Since $(1,1)$ is in the domain, we note $\frac{x^2 - y^2}{x^2 + y^2}$ is rational, hence $\lim_{(x,y) \rightarrow (1,1)} \frac{x^2 - y^2}{x^2 + y^2} = \frac{1^2 - 1^2}{1^2 + 1^2} = 0$.

Ex: Define $g(x,y) = \begin{cases} \frac{x^2-y^2}{x^2+y^2}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$.

Show $\lim_{(x,y) \rightarrow (0,0)} g(x,y) = DNE$.

Useful short-cut: try $y = kx$

$$\lim_{(x,kx) \rightarrow (0,0)} g(x,kx) = \lim_{x \rightarrow 0} \frac{x^2 - k^2 x^2}{x^2 + k^2 x^2} = \lim_{x \rightarrow 0} \frac{1 - k^2}{1 + k^2} = \frac{1 - k^2}{1 + k^2}$$

The limit is k -dependent $\Rightarrow DNE$.

Ex: $\lim_{(x,y) \rightarrow (0,0)} \frac{1}{x^2+y^2} = \infty$; we can

conceivably discuss such limits like in Calc I.

Ex: Is $f(x,y) = \begin{cases} 1, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$ CTS at $(0,0)$?

No; $\lim_{(x,y) \rightarrow (0,0)} f(x,y) = 1 \neq f(0,0) = 0$.

Guidelines to identify continuous functions so you know when certain limits automatically exist

- (1) Sums, differences and products of CTS functions are CTS (check domain)
- (2) Ratios of CTS functions (check domain)
- (3) Compositions of CTS functions are CTS**

$$** \text{ If } g(a) = b = \lim_{x \rightarrow a} g(x) \text{ \& } \lim_{x \rightarrow b} f(x) = f(b)$$

$$\text{then } \lim_{x \rightarrow a} f(g(x)) = f(b) = f(g(a)).$$

EX: Find the limit or write DNE...

$$\lim_{(x,y) \rightarrow (1,0)} \ln \left(\frac{1+y^2}{x^2+xy} \right) = ?$$

$\ln(z)$ is CTS at $z=1$ and $\frac{1+y^2}{x^2+xy}$ is rational / CTS at $(1,0)$ so just plug in

$$(x,y) = (1,0) \dots \lim_{(x,y) \rightarrow (1,0)} \ln \left(\frac{1+y^2}{x^2+xy} \right) = \ln(1) = 0.$$

EX: $\lim_{(x,y) \rightarrow (1,-2)} e^{-x^2 y} \sin(x+y^2) = ?$

$\underbrace{\sin(x+y^2)}_{\text{CTS}}$

$\underbrace{\hspace{10em}}_{\text{CTS}}$

$= e^{-1^2(-2)} \sin(1+(-2)^2) = \boxed{e^2 \sin(5)}$

EX: $\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{\sqrt{x^2+y^2}} = ?$

division by zero = red flag...

The limit turns out to be zero, but it takes thought. Note the numerator is effectively "higher-order" than the denominator, so it vanishes faster as $(x,y) \rightarrow (0,0)$.

This requires an " ϵ - δ " proof... we will skip.

EX: $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^4}{x^2+y^8} = ?$

$(0,0)$ is NOT in the domain... need to investigate.

Along paths like $y=kx$ the limit = 0.

However, y^8 is a red flag... numerator is 5th order. Take $y=x^{1/4}$...

$$\lim_{(x, x^{1/4}) \rightarrow (0,0)} \frac{x \cdot x}{x^2 + x^2} = \frac{1}{2} \neq 0. \quad \boxed{DNE}$$

Practice!

#1


$$(a) \lim_{(x,y) \rightarrow (1,1)} \frac{y^2 \sin^2 x}{x^4 + y^4} = ?$$

$$(b) \lim_{(x,y) \rightarrow (0,0)} \frac{y^2 \sin^2 x}{x^4 + y^4} = ?$$

$$(a) = \frac{(1)^2 \sin^2(1)}{1^4 + 1^4} = \frac{1}{2} \sin^2(1). \quad (\text{CTS})$$

$$(b) = \lim_{(x,y) \rightarrow (0,0)} \frac{y^2 x^2}{x^4 + y^4} \left. \begin{array}{l} y = kx? \quad \frac{k^2 x^4}{x^4(1+k^4)} \\ \text{DNE} \end{array} \right\}$$

#2 Sketch the domain of
 $f(x,y) = \sqrt{x^2-1} - \sqrt{1-y^2}$




#3 Sketch level sets of $f(x,y)=x+y^2$.

$$k=0 \Rightarrow x=-y^2$$

$$k=1 \Rightarrow x=1-y^2$$

$$k=-1 \Rightarrow x=-1-y^2$$



$$\textcircled{\#4} \lim_{(x,y) \rightarrow (0,0)} \frac{x + \sin(y)}{x + y + 1} = \frac{0 + 0}{0 + 0 + 1} = \frac{0}{1} = 0 \text{ (CTS)}$$

$$\textcircled{\#5} \lim_{(x,y) \rightarrow (1,0)} \ln\left(\frac{2x^2 + y^2}{x^2 - 2y^2}\right) = \ln\left(\frac{2}{1}\right) = \ln(2) \text{ (CTS)}$$

$$\textcircled{\#6} \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{2x^2 + y^2} = DNE$$

$$\frac{x^2 + k^2 x^2}{2x^2 + k^2 x^2} = \frac{1 + k^2}{2 + k^2}$$