

# DIMENSION-INDEPENDENT HARNACK INEQUALITIES FOR SUBORDINATED SEMIGROUPS

MARIA GORDINA<sup>†</sup>, MICHAEL RÖCKNER<sup>‡</sup>, AND FENG-YU WANG<sup>\*,\*\*</sup>)

ABSTRACT. Dimension-independent Harnack inequalities are derived for a class of subordinate semigroups. In particular, for a diffusion satisfying the Bakry-Emery curvature condition, the subordinate semigroup with power  $\alpha$  satisfies a dimension-free Harnack inequality provided  $\alpha \in (\frac{1}{2}, 1)$ , and it satisfies the log-Harnack inequality for all  $\alpha \in (0, 1)$ . Some infinite-dimensional examples are also presented.

## CONTENTS

1.	Introduction	1
2.	Proofs	5
3.	Some infinite-dimensional examples	8
3.1.	Stochastic porous medium equation	8
3.2.	Singular stochastic semi-linear equations	9
3.3.	The Ornstein–Uhlenbeck type semigroups with jumps	10
3.4.	Infinite-dimensional Heisenberg groups	11
	References	12

## 1. INTRODUCTION

By using the gradient estimate for diffusion semigroups, the following dimension-free Harnack inequality was established in [19] for the diffusion semigroup  $P_t$  generated by  $L = \Delta + Z$  on a complete Riemannian manifold  $M$  with curvature  $\text{Ric} - \nabla Z$  bounded below by  $-K \in \mathbb{R}$

$$(1.1) \quad (P_t f(x))^p \leq \exp\left(\frac{pK\rho(x,y)^2}{2(p-1)(e^{2Kt}-1)}\right) P_t f^p(y), \quad t > 0, x, y \in M, f \in \mathcal{B}_b^+(M),$$

where  $p > 1$ ,  $\rho$  is the Riemannian distance, and  $\mathcal{B}_b^+(M)$  is the class of all bounded positive measurable functions on  $M$ . This inequality has been extended and applied in the study of contractivity properties, heat kernel bounds, strong Feller properties

---

*Date:* May 17, 2010 *File:*12GRWcomplete.tex.

*1991 Mathematics Subject Classification.* Primary; 60G51, 58G32. Secondary; 58J65, 60J60.

*Key words and phrases.* Harnack inequality, log-Harnack inequality, subordination.

<sup>†</sup> Research was supported in part by NSF Grant DMS-0706784.

<sup>‡</sup> Research was supported in part by the German Science Foundation (DFG) through CRC 701.

<sup>\*</sup> Research was supported in part by WIMICS, NNSFC (10721091) and the 973-Project.

and cost-entropy properties for finite- and infinite-dimensional diffusions. In particular, using the coupling method and Girsanov transformations developed in [4], this inequality has been derived for diffusions without using curvature conditions, see e.g. [5, 6, 9, 13–15, 17, 20] and references therein. See also [1–3] for applications to the short time behavior of transition probabilities. On the other hand, however, due to absence of a chain rule for the “gradient estimate” argument and an explicit Girsanov theorem, this technique of proving a dimension independent Harnack inequalities is not applicable to pure jump processes. The main purpose of this paper is to establish such inequalities for a class of  $\alpha$ -stable like jump processes by using subordination.

Let  $(E, \rho)$  be a Polish space with the Borel  $\sigma$ -algebra  $\mathcal{B}(E)$ , and  $P_t$  the semigroup for a time-homogenous Markov process on  $E$ . Let  $\{\mu_t\}_{t \geq 0}$  be a convolution semigroup of probability measures on  $[0, \infty)$ , i.e. one has  $\mu_{t+s} = \mu_t * \mu_s$  for  $s, t \geq 0$  and  $\mu_t \rightarrow \mu_0 := \delta_0$  weakly as  $t \rightarrow 0$ . Thus, the Laplace transform for  $\mu_t$  has the form

$$(1.2) \quad \int_0^\infty e^{-xs} \mu_t(ds) = e^{-tB(x)}, \text{ for any } x \geq 0, t \geq 0$$

for some Bernstein function  $B$ , see e.g. [12]. We shall study the Harnack inequality for the subordinated semigroup

$$(1.3) \quad P_t^B := \int_0^\infty P_s \mu_t(ds), \quad t \geq 0.$$

Obviously, if  $P_t$  is generated by a negatively definite self-adjoint operator  $(L, \mathcal{D}(L))$  on  $L^2(\nu)$  for some  $\sigma$ -finite measure  $\nu$  on  $E$ , then  $P_t^B$  is generated by  $-B(-L)$ . In particular, if  $B(x) = x^\alpha$  for  $\alpha \in (0, 1]$ , we shall denote the corresponding  $\mu_t$  by  $\mu_t^\alpha$ , and  $P_t^B$  by  $P_t^\alpha$  respectively.

We shall use (1.3) and a known dimension independent Harnack inequality for  $P_t$  to establish the corresponding Harnack inequality for  $P_t^B$ . For instance, suppose we know that

$$(P_t f(x))^p \leq \exp(\Phi(p, t, x, y)) P_t f^p(y), \quad x, y \in E, t > 0, p > 1, f \in \mathcal{B}_b^+(E)$$

for some  $\Phi : (1, \infty) \times (0, \infty) \times E^2 \rightarrow [0, \infty)$ . Then (1.3) implies

$$(1.4) \quad \begin{aligned} (P_t^B f(x))^p &= \left( \int_0^\infty P_s f(x) \mu_t(ds) \right)^p \\ &\leq \left( \int_0^\infty (P_s f^p(y))^{1/p} \exp\left(\frac{\Phi(p, s, x, y)}{p}\right) \mu_t(ds) \right)^p \\ &\leq (P_t^B f^p(y)) \left( \int_0^\infty \exp\left(\frac{\Phi(p, s, x, y)}{p-1}\right) \mu_t(ds) \right)^{p-1}. \end{aligned}$$

In general,  $\Phi(p, s, x, y) \rightarrow \infty$  as  $s \rightarrow 0$ , so we have to verify that  $\exp[\Phi(p, s, x, y)/(p-1)]$  is integrable w.r.t.  $\mu_t(ds)$ . Similarly to (1.1), for many specific models the singularity of  $\Phi(p, s, x, y)$  at  $s = 0$  behaves like  $e^{\delta/s^\kappa}$  for some  $\delta = \delta(p, x, y) > 0, \kappa \geq 1$  (see Section 3 below for specific examples). In this case, the following results say that the Harnack inequality provided by (1.4) is valid for  $P_t^\alpha$  with  $\alpha > \kappa/(\kappa + 1)$ .

**Theorem 1.1.** *Let  $p > 1, \kappa > 0$  and  $\alpha \in \left(\frac{\kappa}{\kappa+1}, 1\right)$  be fixed. Suppose that  $P_t$  satisfies the Harnack inequality*

$$(1.5) \quad (P_t f(x))^p \leq \exp(H(x, y)(\varepsilon + t^{-\kappa})) P_t f^p(y), \quad x, y \in E, f \in \mathcal{B}_b^+(E), t > 0,$$

for some positive measurable function  $H$  on  $E \times E$  and a constant  $\varepsilon \geq 0$ . Then there exists a constant  $c > 0$  depending on  $\alpha$  and  $\kappa$  such that

$$\begin{aligned} & (P_t^\alpha f(x))^p \\ & \leq e^{\varepsilon H(x, y)} \left( 1 + \left[ \exp \left( \left( \frac{cH(x, y)}{(p-1)t^{\kappa/\alpha}} \right)^{1/(1-(\alpha^{-1}-1)\kappa)} \right) - 1 \right]^{(1-(\alpha^{-1}-1)\kappa)} \right)^{p-1} P_t^\alpha f^p(y) \\ & \leq 2^{p-1} \exp \left( \varepsilon H(x, y) + C_{p, \kappa, \alpha} \left( \frac{H(x, y)}{t^{\kappa/\alpha}} \right)^{1/(1-(\alpha^{-1}-1)\kappa)} \right) P_t^\alpha f^p(y), \quad t > 0, x, y \in E \end{aligned}$$

holds for all  $f \in \mathcal{B}_b^+(E)$ , where

$$C_{p, \kappa, \alpha} = \frac{(1 - (\alpha^{-1} - 1)\kappa)c^{1/(1-(\alpha^{-1}-1)\kappa)}}{(p-1)^{(\alpha^{-1}-1)\kappa/(1-(\alpha^{-1}-1)\kappa)}}.$$

Consequently, if  $P_t$  has an invariant probability measure  $\mu$ , we have that

(i) for any  $p, q > 1$ ,

$$\frac{\|P_t^\alpha\|_{p \rightarrow q}}{2^{(p-1)/p}} \leq \left( \int_E \frac{\mu(dx)}{\left( \int_E \exp \left[ -\varepsilon H(x, y) - C_{p, \kappa, \alpha} \left( \frac{H(x, y)}{t^{\kappa/\alpha}} \right)^{1/(1-(\alpha^{-1}-1)\kappa)} \right] \mu(dy) \right)^{q/p}} \right)^{1/q};$$

(ii) if  $P_t^\alpha$  has a transition density  $p_t^\alpha(x, y)$  w.r.t.  $\mu$  such that for any  $x \in \text{supp}(\mu)$

$$\begin{aligned} & \int_E p_t^\alpha(x, y)^2 \mu(dy) \\ & \leq 2 \left( \int_E \exp \left( -\varepsilon H(x, y) - C_{p, \kappa, \alpha} \left( \frac{H(x, y)}{t^{\kappa/\alpha}} \right)^{1/(1-(\alpha^{-1}-1)\kappa)} \right) \mu(dy) \right)^{-1}. \end{aligned}$$

As an application of Theorem 1.1 (ii), we have the following explicit heat kernel upper bounds for stable like processes.

**Example 1.2.** *Let  $P_t$  be generated by  $L = \Delta + Z$  on a complete Riemannian manifold such that  $\text{Ric} - \nabla Z \geq -K$ . By (1.1), (1.5) holds for  $H(x, y) = \rho(x, y)^2$  and  $\kappa = 1$ . So, for  $\alpha \in (1/2, 1]$ , Theorem 1.1 (ii) implies*

$$p_{2t}^\alpha(x, x) \leq \frac{c}{\mu(\{y : \rho(x, y) \leq t^{1/2\alpha}\})}, \quad x \in M, t > 0$$

for some constant  $c > 0$ . In particular, for  $L = \Delta$  on  $\mathbb{R}^d$ ,  $\mu(dx) = dx$  and  $K = 0$ , we have

$$\sup_{x, y \in \mathbb{R}^d} p_t^\alpha(x, y) = \sup_{x \in \mathbb{R}^d} p_t^\alpha(x, x) \leq ct^{-d/2\alpha}, t > 0,$$

for some constant  $c > 0$ . This is sharp due to the well known explicit bounds of heat kernels for the classical stable processes on  $\mathbb{R}^d$ .

Theorem 1.1 does not apply to  $\alpha \in (0, \frac{\kappa}{\kappa+1}]$ , since in this case  $\int_0^\infty e^{\delta/s^\kappa} \mu_t^\alpha(ds) = \infty$  for large  $\delta > 0$ . A more careful analysis allows us to treat the case  $\alpha = \frac{\kappa}{\kappa+1}$  under certain restrictions on  $x, y, t$ . Thus results of this type apply also to the Cauchy process.

**Proposition 1.3** (The case  $\alpha = \frac{\kappa}{\kappa+1}$ ). *Suppose that  $P_t$  satisfies the Harnack inequality (1.5) for some positive measurable function  $H$  on  $E \times E$  and a constant  $\varepsilon \geq 0$ . Then there exists a constant  $C > 0$  depending on  $\kappa$  such that*

$$\begin{aligned} & (P_t^{\frac{\kappa}{\kappa+1}} f(x))^p \\ & \leq e^{\varepsilon H(x,y)} \left( 1 + \frac{C}{\frac{e^{(p-1)}(\kappa t)^{\kappa+1}}{H(x,y)^\kappa} - 1} \right)^{p-1} P_t^{\frac{\kappa}{\kappa+1}} f^p(y), \quad f \in \mathcal{B}_b^+(E) \end{aligned}$$

holds for all  $t > 0, x, y \in E$  such that

$$e(p-1)(t\kappa)^{\kappa+1} > \kappa(\kappa+1)^{\kappa+1} H(x,y).$$

In other cases we can still prove the log-Harnack inequality. For diffusion semigroups, the known log-Harnack inequality looks like

$$(1.6) \quad P_t \log f(x) \leq \log P_t f(y) + H(x,y)(\varepsilon + t^{-\kappa}), \quad x, y \in E, t > 0, f \geq 1,$$

for some positive measurable function  $H$  on  $E \times E$  and some constants  $\varepsilon \geq 0, \kappa \geq 1$ . In many cases, one has  $H(x,y) = c\rho(x,y)^2$  for a constant  $c > 0$  and the intrinsic distance  $\rho$  induced by the diffusion (see e.g. [18]).

**Theorem 1.4.** *If (1.6) holds, then for any  $\alpha \in (0, 1]$ ,*

$$P_t^\alpha \log f(x) \leq \log P_t^\alpha f(y) + H(x,y) \left( \varepsilon + \log P_t^\alpha f(y) + H(x,y) \left( \varepsilon + \frac{\Gamma(\frac{\kappa}{\alpha})}{\alpha t^{\frac{\kappa}{\alpha}} \Gamma(\kappa)} \right) \right),$$

$t > 0, x, y \in E, f \geq 1$ .

As observed in [6] and [18], the log-Harnack inequality implies an entropy-cost inequality for the semigroup and an entropy inequality for the corresponding transition density. Let  $W_H$  be the Wasserstein distance induced by  $H$ , i.e.

$$W_H(\mu_1, \mu_2) = \inf_{\pi \in \mathcal{C}(\mu_1, \mu_2)} \int_{E \times E} H(x,y) \pi(dx, dy),$$

where  $\mu_1, \mu_2$  are probability measures on  $E$  and  $\mathcal{C}(\mu_1, \mu_2)$  is the set of all couplings for  $\mu_1$  and  $\mu_2$ .

**Corollary 1.5.** *Assume that (1.6) holds and let  $P_t$  have an invariant probability measure  $\mu$ . Then for any  $\alpha \in (0, 1]$ :*

(1) *The entropy-cost inequality*

$$\mu(((P_t^\alpha)^* f) \log((P_t^\alpha)^* f)) \leq W_H(f\mu, \mu) \left( \varepsilon + \log P_t^\alpha f(y) + H(x,y) \left( \varepsilon + \frac{\Gamma(\frac{\kappa}{\alpha})}{\alpha t^{\frac{\kappa}{\alpha}} \Gamma(\kappa)} \right) \right),$$

$t > 0, f \geq 0, \mu(f) = 1$

- holds for all  $\alpha \in (0, 1]$ , where  $(P_t^\alpha)^*$  is the adjoint of  $P_t^\alpha$  in  $L^2(E; \mu)$ .
- (2) If  $H(x, y) \rightarrow 0$  as  $y \rightarrow x$  holds for any  $x \in E$ , then  $P_t^\alpha$  is strong Feller and thus has a transition density  $p_t(x, y)$  w.r.t.  $\mu$  on  $\text{supp } \mu$ , which satisfies the entropy inequality

$$\int_E p_t(x, z) \log \frac{p_t(x, z)}{p_t(y, z)} \mu(dz) \leq H(x, y) \left( \varepsilon + \frac{\Gamma\left(\frac{\kappa}{\alpha}\right)}{\alpha t^{\frac{\kappa}{\alpha}} \Gamma(\kappa)} \right), \quad t > 0, x, y \in \text{supp } \mu.$$

## 2. PROOFS

*Proof of Theorem 1.1.* The consequences of the desired Harnack inequality are straightforward. Indeed, (i) follows by noting that the claimed Harnack inequality implies

$$\begin{aligned} & (P_t^\alpha f(x))^p \int_E \exp \left[ -\varepsilon H(x, y) - C_{p, \kappa, \alpha} \left( \frac{H(x, y)}{t^{\kappa/\alpha}} \right)^{1/(1-(\alpha^{-1}-1)\kappa)} \right] \mu(dy) \\ & \leq \mu(P_t^\alpha f^p) = \mu^\alpha(f^p), \end{aligned}$$

which also implies (ii) by taking  $p = 2$  and  $f(z) = p_t^\alpha(x, z)$ ,  $z \in E$ . Indeed, with  $f = 1_A$  for a  $\mu$ -null set  $A$ , this inequality implies that the associated transition probability  $P_t^\alpha(x, \cdot)$  is absolutely continuous w.r.t.  $\mu$  and hence, has a density  $p_t^\alpha(x, \cdot)$  for every  $x \in E$ . Then the desired upper bound for  $\int_E p_t^\alpha(x, y)^2 \mu(dy)$  follows by first applying the above inequality with  $p = 2$  and  $f(z) = p_t^\alpha(x, z) \wedge n$  then letting  $n \rightarrow \infty$ . So, it remains to prove the first assertion.

By (1.5), (1.4) holds for  $\Phi(p, s, x, y) = H(x, y)(\varepsilon + s^{-\kappa})$ , i.e.

$$(2.1) \quad (P_t^\alpha f(x))^p \leq e^{\varepsilon H(x, y)} (P_t^\alpha f^p(y)) \left( \int_0^\infty \exp \left[ \frac{H(x, y)}{(p-1)s^\kappa} \right] \mu_t(ds) \right)^{p-1}.$$

So it suffices to estimate the integral  $\int_0^\infty e^{\delta/s^\kappa} \mu_t(ds)$  for  $\delta := \frac{H(x, y)}{(p-1)} > 0$ .

We use the formula

$$s^{-r} = \frac{1}{\Gamma(r)} \int_0^\infty x^{r-1} e^{-xs} dx, \quad r > 0.$$

to obtain

$$\begin{aligned} \int_0^\infty \frac{\mu_t^\alpha(ds)}{s^r} &= \int_0^\infty \frac{1}{\Gamma(r)} \int_0^\infty x^{r-1} e^{-xs} dx \mu_t(ds) = \\ & \frac{1}{\Gamma(r)} \int_0^\infty x^{r-1} \int_0^\infty e^{-xs} \mu_t(ds) dx = \frac{1}{\Gamma(r)} \int_0^\infty x^{r-1} e^{-tB(x)} dx. \end{aligned}$$

In particular, for  $B(x) = x^\alpha$  we have

$$(2.2) \quad \int_0^\infty \frac{\mu_t^\alpha(ds)}{s^r} = \frac{1}{\Gamma(r)} \int_0^\infty x^{r-1} e^{-tx^\alpha} dx = \frac{1}{\alpha \Gamma(r)} \int_0^\infty y^{\frac{r}{\alpha}-1} e^{-ty} dy = \frac{\Gamma\left(\frac{r}{\alpha}\right)}{\alpha \Gamma(r)} t^{-\frac{r}{\alpha}}.$$

We can use the generalization of Stirling's formula giving the asymptotic behavior of the Gamma function for large  $r$

$$\Gamma(r) = \sqrt{2\pi} r^{r-\frac{1}{2}} e^{-r+\eta(r)},$$

where

$$\eta(r) = \sum_{n=0}^{\infty} \left( r + n + \frac{1}{2} \right) \ln \left( 1 + \frac{1}{r+n} \right) - 1 = \frac{\theta}{12r}, 0 < \theta < 1.$$

We apply this estimate to  $\Gamma(\kappa n)$ ,  $\Gamma\left(\frac{\kappa n}{\alpha}\right)$  and  $n!$ . Thus

$$\begin{aligned} \int_0^{\infty} e^{\frac{\delta}{s^{\kappa}}} \mu_t^{\alpha}(ds) &= 1 + \sum_{n=1}^{\infty} \frac{\delta^n}{n!} \frac{\Gamma\left(\frac{\kappa n}{\alpha}\right)}{\alpha \Gamma(\kappa n)} t^{-\frac{\kappa n}{\alpha}} = \\ &1 + \frac{1}{\alpha} \sum_{n=1}^{\infty} \frac{\delta^n}{n!} (\kappa n)^{\kappa n \left(\frac{1}{\alpha}-1\right)} e^{-\kappa n \left(\frac{1}{\alpha}-1\right)} \alpha^{\frac{1}{2}-\frac{\kappa n}{\alpha}} e^{\frac{\theta_1 \alpha - \theta_2}{12 \kappa n}} t^{-\frac{\kappa n}{\alpha}} \leq \\ (2.3) \quad &1 + \frac{1}{\sqrt{\alpha}} \sum_{n=1}^{\infty} \frac{\delta^n}{n!} (\kappa n)^{\kappa n \left(\frac{1}{\alpha}-1\right)} e^{-\kappa n \left(\frac{1}{\alpha}-1\right)} \alpha^{-\frac{\kappa n}{\alpha}} e^{\frac{\alpha}{12 \kappa n}} t^{-\frac{\kappa n}{\alpha}} = \\ &1 + \frac{1}{\sqrt{\alpha}} \sum_{n=1}^{\infty} \frac{n^{\kappa n \left(\frac{1}{\alpha}-1\right)}}{n!} \left( \delta \left( \frac{\kappa}{e} \right)^{\kappa \left(\frac{1}{\alpha}-1\right)} \alpha^{-\frac{\kappa}{\alpha}} t^{-\frac{\kappa}{\alpha}} \right)^n e^{\frac{\alpha}{12 \kappa n}} \leq \\ &1 + \frac{1}{\sqrt{2\pi\alpha}} \sum_{n=1}^{\infty} n^{\kappa n \left(\frac{1}{\alpha}-1\right) - n - \frac{1}{2}} \left( \delta \left( \frac{\kappa}{e} \right)^{\kappa \left(\frac{1}{\alpha}-1\right)} \alpha^{-\frac{\kappa}{\alpha}} t^{-\frac{\kappa}{\alpha}} \right)^n e^{\frac{\alpha}{12 \kappa n}}. \end{aligned}$$

This series converges for  $\alpha > \frac{\kappa}{\kappa+1}$ , moreover, there is a constant  $c$  depending only on  $\kappa$  such that

$$\frac{1}{\sqrt{2\pi\alpha n}} \left( \left( \frac{\kappa}{e} \right)^{\kappa \left(\frac{1}{\alpha}-1\right)} \alpha^{-\frac{\kappa}{\alpha}} t^{-\frac{\kappa}{\alpha}} \right)^n e^{\frac{\alpha}{12 \kappa n}} \leq c^n.$$

Denote

$$c(\delta, \alpha, \kappa) := 1 + \sum_{n=1}^{\infty} n^{\kappa \left(\frac{1}{\alpha}-1\right) - 1} (c\delta t^{-\frac{\kappa}{\alpha}})^n,$$

then

$$(P_t^{\alpha} f(x))^p \leq e^{\varepsilon H(x,y)} \left( c \left( \frac{H(x,y)}{p-1}, \alpha, \kappa \right) \right)^{p-1} P_t^{\alpha} f^p(y).$$

Note that for  $a > 0, 1 \geq b > 0$  we have the following estimate

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{a^n}{n^{bn}} &= \sum_{n=1}^{\infty} \frac{(2a)^n}{n^{bn}} \frac{1}{2^n} \leq \left( \sum_{n=1}^{\infty} \frac{(2a)^{\frac{n}{b}}}{n^n} \frac{1}{2^n} \right)^b \leq \\ &\left( \sum_{n=1}^{\infty} \frac{(2a)^{\frac{n}{b}}}{n!} \frac{1}{2^n} \right)^b = \left( e^{\frac{(2a)^{1/b}}{2}} - 1 \right)^b, \end{aligned}$$

where we used Jensen's inequality. Thus for any  $\alpha \in \left( \frac{\kappa}{\kappa+1}, 1 \right)$  we use the above estimate with  $b := \kappa \left( 1 - \frac{1}{\alpha} \right) + 1 \leq 1$  to see that

$$c(\delta, \alpha, \kappa) = 1 + \sum_{n=1}^{\infty} n^{\kappa(\frac{1}{\alpha}-1)-1} (c\delta t^{-\frac{\kappa}{\alpha}})^n \leq 1 + \left( \exp \left( \frac{(2c\delta t^{-\frac{\kappa}{\alpha}})^{\kappa(\frac{1}{\alpha}-1)+1}}{2} \right) - 1 \right)^{\kappa(1-\frac{1}{\alpha})+1}.$$

Thus we can say that there is  $c > 0$  depending on  $\alpha$  and  $\kappa$  such that

$$\int_0^{\infty} e^{\frac{H(x,y)}{(p-1)s^{\kappa}}} \mu_t^{\alpha}(ds) \leq 1 + \left( \exp \left( \left( \frac{cH(x,y)}{(p-1)t^{\frac{\kappa}{\alpha}}} \right)^{\kappa(\frac{1}{\alpha}-1)+1} \right) - 1 \right)^{\kappa(1-\frac{1}{\alpha})+1}$$

Using the inequality

$$1 + (x-1)^a \leq 2x^a$$

for any  $x \geq 1$  and  $0 \leq a \leq 1$  we see that

$$\int_0^{\infty} e^{\frac{\delta}{s^{\kappa}}} \mu_t^{\alpha}(ds) \leq 2 \exp \left( \left( \kappa \left( 1 - \frac{1}{\alpha} \right) + 1 \right) \left( \frac{cH(x,y)}{(p-1)t^{\frac{\kappa}{\alpha}}} \right)^{\kappa(\frac{1}{\alpha}-1)+1} \right)$$

which completes the proof. □

*Proof of Proposition 1.3.* In the case  $\alpha = \frac{\kappa}{\kappa+1}$  the series in (2.3) converges for  $t > 0$  and  $x, y \in E$  such that

$$(2.4) \quad e(p-1)(t\kappa)^{\kappa+1} > \kappa(\kappa+1)^{\kappa+1}H(x,y).$$

Note that for  $\delta := \frac{H(x,y)}{p-1}$  the last line of (2.3) reduces to

$$\begin{aligned} & 1 + \sqrt{\frac{\kappa+1}{2\pi\kappa}} \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} \left( \frac{\delta\kappa}{e} \left( \frac{\kappa+1}{\kappa t} \right)^{\kappa+1} \right)^n e^{\frac{1}{12(\kappa+1)n}} \\ & \leq 1 + C \sum_{n=1}^{\infty} \left( \frac{\delta\kappa}{e} \left( \frac{\kappa+1}{\kappa t} \right)^{\kappa+1} \right)^n \\ & = 1 + \frac{C}{\frac{e}{\delta\kappa} \left( \frac{\kappa t}{\kappa+1} \right)^{\kappa+1} - 1}. \end{aligned}$$

This completes the proof. □

*Proof of Theorem 1.4.* By (2.2) with  $r = \kappa$ , we have

$$\int_0^{\infty} \frac{\mu_t^{\alpha}(ds)}{s^{\kappa}} = \frac{\Gamma\left(\frac{\kappa}{\alpha}\right)}{\alpha t^{\frac{\kappa}{\alpha}} \Gamma(\kappa)}.$$

Using (1.2), (1.6) we obtain

$$\begin{aligned}
P_t^\alpha \log f(x) &= \int_0^\infty P_s \log f(x) \mu_t^\alpha(ds) \leq \int_0^\infty (\log P_s f(y) + H(x, y)(\varepsilon + s^{-\kappa})) \mu_t^\alpha(ds) \\
&= \log P_t^\alpha f(y) + H(x, y) \left( \varepsilon + \frac{\Gamma\left(\frac{\kappa}{\alpha}\right)}{\alpha t^{\frac{\kappa}{\alpha}} \Gamma(\kappa)} \right).
\end{aligned}$$

This completes the proof.  $\square$

*Proof of Corollary 1.5.* (1) It suffices to prove for  $f \in \mathcal{B}_b^+(E)$  such that  $\inf f > 0$  and  $\mu(f) = 1$ . In this case, there exists a constant  $c > 0$  such that  $cf \geq 1$ . By Theorem 1.4 for  $cP_t^\alpha f$  in place of  $f$ , we obtain

$$P_t^\alpha \log(P_t^\alpha)^* f(x) \leq \log P_t^\alpha (P_t^\alpha)^* f(y) + H(x, y) \left( \varepsilon + \frac{\Gamma\left(\frac{\kappa}{\alpha}\right)}{\alpha t^{\frac{\kappa}{\alpha}} \Gamma(\kappa)} \right).$$

Since  $\mu$  is invariant for  $P_t^\alpha$  and  $(P_t^\alpha)^*$ , taking the integral for both sides w.r.t.  $\pi \in (f\mu, \mu)$  and minimizing in  $\pi$ , we prove the first assertion.

(2) The strong Feller property follows from Theorem 1.4 according to [18, Proposition 2.3], while by [18, Proposition 2.4] the desired entropy inequality for the transition density is equivalent to the log-Harnack inequality for  $P_t^\alpha$  provided by Theorem 1.4.  $\square$

### 3. SOME INFINITE-DIMENSIONAL EXAMPLES

As explained in Section 1, Theorems 1.1 and 1.4 hold for  $\kappa = 1$  if  $P_t$  is a diffusion semigroup on a Riemannian manifold with the Ricci curvature bounded below. In this section we present some infinite dimensional examples where these theorems can be used.

**3.1. Stochastic porous medium equation.** Let  $\Delta$  be the Dirichlet Laplace operator on a bounded interval  $(a, b)$  and  $W_t$  the cylindrical Brownian motion on  $L^2((a, b); dx)$ . Since the eigenvalues  $\{\lambda_i\}$  of  $-\Delta$  satisfies  $\sum_{i=1}^\infty \lambda_i^{-1} < \infty$ ,  $W_t$  is a continuous process on  $\mathbb{H}$ , the completion of  $L^2((a, b); dx)$  under the inner product

$$\langle x, y \rangle := \sum_{i=1}^\infty \frac{1}{\lambda_i} \langle x, e_i \rangle \langle y, e_i \rangle,$$

where  $e_i$  is the unit eigenfunction corresponding to  $\lambda_i$  for each  $i \geq 1$ . Let  $\|\cdot\|$  denote the norm on  $\mathbb{H}$ , and suppose  $r > 1$ . Then the following stochastic porous medium equation has a unique strong solution on  $\mathbb{H}$  for any  $X_0 \in \mathbb{H}$  (see e.g. [7]):

$$dX_t = \Delta X_t^r dt + dW_t.$$

Let  $P_t$  be the corresponding Markov semigroup. According to [20, Remark 1.1 and Theorem 1.2], Theorem 1.1 in [20] holds for  $\theta = r - 1$  and some constant  $\gamma, \delta, \xi > 0$ . Thus, there exist two constants  $c_1, c_2 > 0$  depending on  $r$  such that

$$(P_t f)^p(x) \leq (P_t f^p(y)) \exp \left[ \frac{c_1 p \|x - y\|^{4/(1+r)}}{(p-1)(1 - e^{-c_2 t})^{(3+r)/(1+r)}} \right], \quad p > 1, t > 0, x, y \in \mathbb{H}$$



holds for all  $f \in \mathcal{B}_b^+(\mathbb{H})$ . By [18, Proposition 2.2] for  $\rho(x, y)^2 = \|x - y\|^{2/(1+r)}$ , this implies the log-Harnack inequality

$$P_t \log f(x) \leq \log P_t f(x) + \frac{c_1 \|x - y\|^{4/(1+r)}}{(1 - e^{-c_2 t})^{(3+r)/(1+r)}}, \quad x, y \in \mathbb{H}, f \geq 1.$$

Therefore, Theorems 1.1 and 1.4 apply to  $P_t^\alpha$  for

$$\kappa = \frac{r + r}{1 + r}$$

and some constant  $\varepsilon$  depending on  $r$ .

**3.2. Singular stochastic semi-linear equations.** Let  $\mathbb{H}$  be a separable Hilbert space with inner product  $\langle \cdot, \cdot \rangle$ , and  $W_t$  the cylindrical Brownian motion on  $\mathbb{H}$ . Consider the stochastic equation

$$(3.1) \quad dX_t = (AX_t + F(X_t))dt + \sigma dW_t, \quad X_0 \in H.$$

Let  $A, F$  and  $\sigma$  satisfy the following hypotheses:

(H1)  $(A, \mathcal{D}(A))$  is the generator of a  $C_0$ -semigroup,  $T_t = e^{tA}$ ,  $t \geq 0$ , on  $\mathbb{H}$  and for some  $\omega \in \mathbb{R}$

$$(3.2) \quad \langle Ax, x \rangle \leq \omega \|x\|^2, \quad \forall x \in \mathcal{D}(A).$$

(H2)  $\sigma$  is a bounded positively definite, self-adjoint operator on  $\mathbb{H}$  such that  $\sigma^{-1}$  is bounded and  $\int_0^\infty \|T_t \sigma\|_{HS}^2 dt < \infty$ , where  $\|\cdot\|_{HS}$  denotes the norm on the space of all Hilbert-Schmidt operators on  $\mathbb{H}$ .

(H3)  $F : \mathcal{D}(F) \subset \mathbb{H} \rightarrow \mathbb{H}$  is an  $m$ -dissipative map, i.e.,

$$\langle F(x) - F(y), x - y \rangle \leq 0, \quad x, y \in \mathcal{D}(F), u \in F(x), v \in F(y),$$

(“dissipativity”) and

$$\text{Range}(I - F) := \bigcup_{x \in \mathcal{D}(F)} (x - F(x)) = \mathbb{H}.$$

Furthermore,  $F_0(x) \in F(x)$ ,  $x \in \mathcal{D}(F)$ , is such that

$$\|F_0(x)\| = \min_{y \in F(x)} \|y\|.$$

Here we recall that for  $F$  as in (H3) we have that  $F(x)$  is closed, non empty and convex.

The corresponding Kolmogorov operator is then given as follows: Let  $\mathcal{E}_A(H)$  denote the linear span of all real parts of functions of the form  $\varphi = e^{i\langle h, \cdot \rangle}$ ,  $h \in D(A^*)$ , where  $A^*$  denotes the adjoint operator of  $A$ , and define for any  $x \in \mathcal{D}(F)$ ,

$$L_0 \varphi(x) = \frac{1}{2} \text{Tr}(\sigma^2 D^2 \varphi(x)) + \langle x, A^* D \varphi(x) \rangle + \langle F_0(x), D \varphi(x) \rangle, \quad \varphi \in \mathcal{E}_A(H).$$

Additionally, we assume:

(H4) There exists a probability measure  $\mu$  on  $H$  (equipped with its Borel  $\sigma$ -algebra  $\mathcal{B}(H)$ ) such that

- (i)  $\mu(\mathcal{D}(F)) = 1$ ,
- (ii)  $\int_H (1 + \|x\|^2)(1 + \|F_0(x)\|) \mu(dx) < \infty$ ,
- (iii)  $\int_H L_0 \varphi d\mu = 0$  for all  $\varphi \in \mathcal{E}_A(H)$ .

By [8], the closure of  $(L_0, \mathcal{E}_A(\mathbb{H}))$  in  $L^1(\mathbb{H}; \mu)$  generates a Markov semigroup  $P_t$  with  $\mu$  as an invariant probability measure, which is point-wisely determined on  $\mathbb{H}_0 := \text{supp}\mu$ . If moreover the following hypotheses holds:

(H5) (i)  $(1 + \omega - A, \mathcal{D}(A))$  satisfies the weak sector condition: there exists a constant  $K > 0$  such that

$$(3.3) \quad \langle (1 + \omega - A)x, y \rangle \leq K \langle (1 + \omega - A)x, x \rangle^{1/2} \langle (1 + \omega - A)y, y \rangle^{1/2}, \quad \forall x, y \in \mathcal{D}(A).$$

(ii) There exists a sequence of  $A$ -invariant finite dimensional subspaces  $\mathbb{H}_n \subset \mathcal{D}(A)$  such that  $\bigcup_{n=1}^{\infty} \mathbb{H}_n$  is dense in  $\mathbb{H}$ .

Then (see [9, Theorem 1.6])

$$(P_t f(x))^p \leq P_t f^p(y) \exp \left[ \|\sigma^{-1}\|^2 \frac{p\omega \|x - y\|^2}{(p-1)(1 - e^{-2\omega t})} \right], \quad t > 0, \quad x, y \in \mathbb{H}_0.$$

As mentioned above, according to [18, Proposition 2.2] this implies the corresponding log-Harnack inequality. Therefore, our Theorems 1.1 and 1.4 apply to  $P_t^p$  for  $\kappa = 1$ .

**3.3. The Ornstein–Uhlenbeck type semigroups with jumps.** Consider the following stochastic differential equation driven by a Lévy process

$$(3.4) \quad dX_t = AX_t dt + dZ_t, \quad X_0 = x \in \mathbb{H},$$

where  $A$  is the infinitesimal generator of a strongly continuous semigroup  $(T_t)_{t \geq 0}$  on  $\mathbb{H}$ ,  $Z_t := \{Z_t^u, u \in \mathbb{H}\}$  is a cylindrical Lévy process with characteristic triplet  $(a, R, M)$  on some filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ , that is, for every  $u \in \mathbb{H}$  and  $t \geq 0$

$$\begin{aligned} \mathbb{E} \exp(i \langle Z_t, u \rangle) &= \exp(it \langle a, u \rangle - \frac{t}{2} \langle Ru, u \rangle \\ &\quad - \int_{\mathbb{H}} [1 - \exp(i \langle x, u \rangle) + i \langle x, u \rangle 1_{\{\|x\| \leq 1\}}(x)] M(dx)), \end{aligned}$$

where  $a \in \mathbb{H}$ ,  $R$  is a symmetric linear operator on  $\mathbb{H}$  such that

$$R_t := \int_0^t T_s R T_s^* ds$$

is a trace class operator for each  $t > 0$ , and  $M$  is a Lévy measure on  $\mathbb{H}$ . (For simplicity, we shall write  $Z_t^u = \langle Z_t, u \rangle$  for every  $u \in \mathbb{H}$ .) In this case, (3.4) has a unique mild solution

$$X_t = T_t x + \int_0^t T_{t-s} dZ_s, \quad t \geq 0.$$

Let

$$P_t f(x) = \mathbb{E} f(X_t), \quad x \in \mathbb{H}, \quad f \in \mathbb{B}_b(\mathbb{H}).$$

If

$$\|R^{-1/2} T_t R x\| \leq \sqrt{h(t)} \|R^{1/2} x\|, \quad x \in \mathbb{H}, \quad t \geq 0$$

holds for some positive function  $h \in C([0, \infty))$ . Then by [16, Theorem 1.2] (see also [17] for the diffusion case),

$$(P_t f)^\alpha(x) \leq \exp \left[ \frac{\alpha \|R^{-1/2}(x-y)\|^2}{2(\alpha-1) \int_0^t h(s)^{-1} ds} \right] P_t f^\alpha(y), \quad t > 0, x-y \in R^{1/2}\mathbb{H}$$

holds for all  $f \in \mathcal{B}_b^+(\mathbb{H})$ . By this and [18, Proposition 2.2] which implies the corresponding log-Harnack inequality, Theorems 1.1 and 1.4 apply to some  $\varepsilon \geq 0$  and  $\kappa \geq 1$  if

$$\limsup_{t \rightarrow 0} \frac{1}{t^\kappa} \int_0^t \frac{ds}{h(s)} > 0.$$

**3.4. Infinite-dimensional Heisenberg groups.** In [10] an integrated Harnack inequality similar to (1.1) has been established for a Brownian motion on infinite-dimensional Heisenberg groups modeled on an abstract Wiener space. The inequality is the consequence of the Ricci curvature bounds for both finite-dimensional approximations to these groups and the group itself, and the results established for inductive limits of finite-dimensional Lie groups in [11]. Even though the methods described in that paper are applicable to inductive and projective limits of finite-dimensional Lie groups, the infinite-dimensional Heisenberg groups provide a very concrete setting. We follow the exposition in [10].

Let  $(W, H, \mu)$  be an abstract Wiener space over  $\mathbb{R}(\mathbb{C})$ ,  $\mathbf{C}$  be a real(complex) finite dimensional inner product space, and  $\omega : W \times W \rightarrow \mathbf{C}$  be a continuous skew symmetric bilinear quadratic form on  $W$ . Further, let

$$(3.5) \quad \|\omega\|_0 := \sup \{ \|\omega(w_1, w_2)\|_{\mathbf{C}} : w_1, w_2 \in W \text{ with } \|w_1\|_W = \|w_2\|_W = 1 \}$$

be the uniform norm on  $\omega$  which is finite since  $\omega$  is assumed to be continuous. We will need the Hilbert-Schmidt norm of  $\omega$  which is defined as

$$\|\omega\|_2^2 = \|\omega\|_{H^* \otimes H^* \otimes \mathbf{C}} := \sum_{i,j=1}^{\infty} \|\omega(e_i, e_j)\|_{\mathbf{C}}^2,$$

which is finite by Proposition 3.14 in [10].

**Definition 3.1.** Let  $\mathfrak{g}$  denote  $W \times \mathbf{C}$  when thought of as a Lie algebra with the Lie bracket operation given by

$$(3.6) \quad [(A, a), (B, b)] := (0, \omega(A, B)).$$

Let  $G := G(\omega)$  denote  $W \times \mathbf{C}$  when thought of as a group with the multiplication law given by

$$(3.7) \quad g_1 g_2 = g_1 + g_2 + \frac{1}{2} [g_1, g_2] \text{ for any } g_1, g_2 \in G.$$

It is easily verified that  $\mathfrak{g}$  is a Lie algebra and  $G$  is a group. The identity of  $G$  is the zero element,  $\mathbf{e} := (0, 0)$ .

**Notation 3.2.** Let  $\mathfrak{g}_{CM}$  denote  $H \times \mathbf{C}$  when viewed as a Lie subalgebra of  $\mathfrak{g}$  and  $G_{CM}$  denote  $H \times \mathbf{C}$  when viewed as a subgroup of  $G = G(\omega)$ . We will refer to  $\mathfrak{g}_{CM}$  ( $G_{CM}$ ) as the **Cameron–Martin subalgebra (subgroup)** of  $\mathfrak{g}$  ( $G$ ). (For explicit examples of such  $(W, H, \mathbf{C}, \omega)$ , see [10].)

We equip  $G = \mathfrak{g} = W \times \mathbf{C}$  with the Banach space norm

$$(3.8) \quad \|(w, c)\|_{\mathfrak{g}} := \|w\|_W + \|c\|_{\mathbf{C}}$$

and  $G_{CM} = \mathfrak{g}_{CM} = H \times \mathbf{C}$  with the Hilbert space inner product,

$$(3.9) \quad \langle (A, a), (B, b) \rangle_{\mathfrak{g}_{CM}} := \langle A, B \rangle_H + \langle a, b \rangle_{\mathbf{C}}.$$

The associate Hilbertian norm is given by

$$(3.10) \quad \|(A, \delta)\|_{\mathfrak{g}_{CM}} := \sqrt{\|A\|_H^2 + \|\delta\|_{\mathbf{C}}^2}.$$

As was shown in [10, Lemma 3.3], these Banach space topologies on  $W \times \mathbf{C}$  and  $H \times \mathbf{C}$  make  $G$  and  $G_{CM}$  into topological groups.

Then we can define a Brownian motion on  $G$  starting at  $\mathbf{e} = (0, 0) \in G$  to be the process

$$(3.11) \quad g(t) = \left( B(t), B_0(t) + \frac{1}{2} \int_0^t \omega(B(\tau), dB(\tau)) \right).$$

We denote by  $\nu_t$  the corresponding heat kernel measure on  $G$ . The following estimate was used in the proof of Theorem 8.1 in [10]. For any  $h \in G_{CM}$ ,  $1 < p < \infty$

$$(3.12) \quad \int_G |f(xh)| d\nu_t(x) \leq \|f\|_{L^p(G, \nu_t)} \exp\left(\frac{c(-k(\omega)t)(p-1)}{2t} d_{G_{CM}}^2(e, h)\right).$$

where

$$c(t) = \frac{t}{e^t - 1} \quad \text{for all } t \in \mathbb{R}$$

with the convention that  $c(0) = 1$  and

$$k(\omega) := \frac{1}{2} \sup_{\|A\|_H=1} \|\omega(\cdot, A)\|_{H^* \otimes \mathbf{C}}^2 \leq \frac{1}{2} \|\omega\|_2^2 < \infty.$$

Equation (3.12) implies the corresponding  $L^p$ -estimates of Radon-Nikodym derivatives of  $\nu_t$  relative to the left and right multiplication by elements in  $G_{CM}$ . This in turn is equivalent to the Harnack inequality (1.1) following an argument similar to Lemma D.1 in [11]

$$[(P_t f)(x)]^p \leq C^p (P_t f^p)(y) \quad \text{for all } f \geq 0.$$

Thus we are in position to apply our results to the heat kernel measure  $\nu_t$  subordinated as described in Section 1.

## REFERENCES

- [1] Shigeki Aida, *Uniform positivity improving property, Sobolev inequalities, and spectral gaps*, J. Funct. Anal. **158** (1998), no. 1, 152–185. MR MR1641566 (2000d:60125)
- [2] Shigeki Aida and Hiroshi Kawabi, *Short time asymptotics of a certain infinite dimensional diffusion process*, Stochastic analysis and related topics, VII (Kusadasi, 1998), Progr. Probab., vol. 48, Birkhäuser Boston, Boston, MA, 2001, pp. 77–124. MR MR1915450 (2003m:60219)
- [3] Shigeki Aida and Tusheng Zhang, *On the small time asymptotics of diffusion processes on path groups*, Potential Anal. **16** (2002), no. 1, 67–78. MR MR1880348 (2003e:58052)
- [4] Marc Arnaudon, Anton Thalmaier, and Feng-Yu Wang, *Harnack inequality and heat kernel estimates on manifolds with curvature unbounded below*, Bull. Sci. Math. **130** (2006), no. 3, 223–233. MR MR2215664 (2007i:58032)
- [5] ———, *Gradient estimates and harnack inequalities on non-compact riemannian manifolds*, to appear **x** (2009), x.

- [6] Sergey G. Bobkov, Ivan Gentil, and Michel Ledoux, *Hypercontractivity of Hamilton-Jacobi equations*, J. Math. Pures Appl. (9) **80** (2001), no. 7, 669–696. MR MR1846020 (2003b:47073)
- [7] G. Da Prato, Michael Röckner, B. L. Rozovskii, and Feng-Yu Wang, *Strong solutions of stochastic generalized porous media equations: existence, uniqueness, and ergodicity*, Comm. Partial Differential Equations **31** (2006), no. 1-3, 277–291. MR MR2209754 (2007b:60153)
- [8] Giuseppe Da Prato and Michael Röckner, *Singular dissipative stochastic equations in Hilbert spaces*, Probab. Theory Related Fields **124** (2002), no. 2, 261–303. MR MR1936019 (2003k:60151)
- [9] Giuseppe Da Prato, Michael Röckner, and Feng-Yu Wang, *Singular stochastic equations on Hilbert spaces: Harnack inequalities for their transition semigroups*, J. Funct. Anal. **257** (2009), no. 4, 992–1017. MR MR2535460
- [10] Bruce K. Driver and Maria Gordina, *Heat kernel analysis on infinite-dimensional Heisenberg groups*, J. Funct. Anal. **255** (2008), no. 9, 2395–2461. MR MR2473262
- [11] ———, *Integrated Harnack inequalities on Lie groups*, Journal of Differential Geometry **3** (2009), 501–550.
- [12] N. Jacob, *Pseudo differential operators and Markov processes. Vol. I*, Imperial College Press, London, 2001, Fourier analysis and semigroups. MR MR1873235 (2003a:47104)
- [13] Wei Liu, *Fine properties of stochastic evolution equations and their applications*, Ph.D. thesis, Doctor-Thesis, Bielefeld University, 2009.
- [14] Wei Liu and Feng-Yu Wang, *Harnack inequality and strong Feller property for stochastic fast-diffusion equations*, J. Math. Anal. Appl. **342** (2008), no. 1, 651–662. MR MR2440828 (2009k:60137)
- [15] Shun-Xiang Ouyang, *Harnack inequalities and applications for stochastic equations*, Ph.D. thesis, Ph.D. thesis, Bielefeld University, 2009.
- [16] Shun-Xiang Ouyang, Michael Röckner, and Feng-Yu Wang, *Harnack inequalities and applications for Ornstein-Uhlenbeck semigroups with jumps*, preprint.
- [17] Michael Röckner and Feng-Yu Wang, *Harnack and functional inequalities for generalized Mehler semigroups*, J. Funct. Anal. **203** (2003), no. 1, 237–261. MR MR1996872 (2005d:47077)
- [18] Feng-Yu Wang, *Heat kernel inequalities for curvature and second fundamental form*, preprint.
- [19] ———, *Logarithmic Sobolev inequalities on noncompact Riemannian manifolds*, Probab. Theory Related Fields **109** (1997), no. 3, 417–424. MR MR1481127 (98i:58253)
- [20] ———, *Harnack inequality and applications for stochastic generalized porous media equations*, Ann. Probab. **35** (2007), no. 4, 1333–1350. MR MR2330974 (2008e:60192)

† DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269, U.S.A.  
*E-mail address:* `gordina@math.uconn.edu`

‡ DEPARTMENT OF MATHEMATICS, BIELEFELD UNIVERSITY, D-33501 BIELEFELD, GERMANY

‡ DEPARTMENTS OF MATHEMATICS AND STATISTICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY ST WEST LAFAYETTE, IN 47907-2067 USA  
*E-mail address:* `roeckner@math.uni-bielefeld.de`

\*DEPARTMENT OF MATHEMATICS, SWANSEA UNIVERSITY, SINGLETON PARK, SA2 8PP, UK  
*E-mail address:* `wangfy@bnu.edu.cn`

\*\* SCHOOL OF MATH. SCI. & LAB. MATH. COM. SYS., BEIJING NORMAL UNIVERSITY, BEIJING 100875, CHINA  
*E-mail address:* `wangfy@bnu.edu.cn`