

# HEAT KERNEL ANALYSIS AND CAMERON-MARTIN SUBGROUP FOR INFINITE DIMENSIONAL GROUPS

MARIA GORDINA

*Department of Mathematics and Statistics,  
McMaster University,  
Hamilton, ON L8S 4K1  
Canada*

ABSTRACT. The heat kernel measure  $\mu_t$  is constructed on  $GL(H)$ , the group of invertible operators on a complex Hilbert space  $H$ . This measure is determined by an infinite dimensional Lie algebra  $\mathfrak{g}$  and a Hermitian inner product on it. The Cameron-Martin subgroup  $G_{CM}$  is defined and its properties are discussed. In particular, there is an isometry from the  $L^2_{\mu_t}$ -closure of holomorphic polynomials into a space  $\mathcal{H}^t(G_{CM})$  of functions holomorphic on  $G_{CM}$ . This means that any element from this  $L^2_{\mu_t}$ -closure of holomorphic polynomials has a version holomorphic on  $G_{CM}$ . In addition, there is an isometry from  $\mathcal{H}^t(G_{CM})$  into a Hilbert space associated with the tensor algebra over  $\mathfrak{g}$ . The latter isometry is an infinite dimensional analog of the Taylor expansion. As examples we discuss a complex orthogonal group and a complex symplectic group.

## TABLE OF CONTENTS

1. Introduction	2
2. Notation and main results	2
3. Estimates of derivatives of holomorphic functions	5
4. Isometries	7
5. The heat kernel measure	10
6. Approximation of the process	16
7. Cameron-Martin subgroup	19
8. Holomorphic polynomials and skeletons	24
9. Examples	28
References	31

---

*Key words and phrases.* Heat kernel measure, holomorphic function, infinite dimensional group, infinite dimensional Lie algebra, stochastic differential equation.

## 1. INTRODUCTION

Our goal is to study Hilbert spaces of holomorphic functions on a group associated with an infinite dimensional Lie algebra  $\mathfrak{g}$  which is itself equipped with a Hermitian inner product  $(\cdot, \cdot)$  and corresponding norm  $|\cdot|$ . We assume that  $\mathfrak{g}$  is a Lie subalgebra of  $B(H)$ , the space of bounded linear operators on a complex separable Hilbert space  $H$ . The group under consideration is a Lie subgroup of  $GL(H)$ , the group of invertible elements of  $B(H)$ . Note that  $B(H)$  is the natural (infinite dimensional) Lie algebra of  $GL(H)$  with the commutator as a Lie bracket.

One of the main ingredients in this work is the construction of the heat kernel measure on  $GL(H)$  which is determined by  $\mathfrak{g}$  and the norm  $|\cdot|$ . In some cases it is possible to show that the heat kernel measure is supported in a subgroup of  $GL(H)$  (see Section 9). The construction of the heat kernel measure requires the use of stochastic differential equations in a Hilbert space. We will assume that  $\mathfrak{g}$  is a subspace of the Hilbert-Schmidt operators on  $H$ .

It is well known that the Cameron-Martin subspace plays an important role for an abstract Wiener space. Analogously, we define the Cameron-Martin subgroup,  $G_{CM}$ , and discuss its properties. One of these properties is that functions in the  $L^2$ -closure of holomorphic polynomials have holomorphic versions on  $G_{CM}$ . Following [15, 16] we call these versions skeletons. The map taking an  $L^2$ -function to its skeleton is an isometry to  $\mathcal{H}^t(G_{CM})$ , a space of functions holomorphic on  $G_{CM}$  with a direct limit-type norm derived from finite dimensional approximations to  $G_{CM}$ . We also show that the Taylor map, from holomorphic functions on  $G_{CM}$  into a dual of the universal enveloping algebra of  $\mathfrak{g}$ , is isometric on  $\mathcal{H}^t(G_{CM})$ . This isometry is a noncommutative version of one of the isomorphisms between different representations of a bosonic Fock space.

An outline of the history of the subject has been given in [7, 8]. We should mention here works by Sugita [15, 16] for an abstract complex Wiener space, in particular, his results on skeletons of  $L^2$ -functions on the Cameron-Martin subspace.

**Acknowledgement.** I thank Professor B. Driver, Professor L. Gross and Professor B. Hall for their help throughout the process of preparation of this work.

## 2. NOTATION AND MAIN RESULTS

To describe results of this paper in more detail we need to use finite dimensional approximations to  $\mathfrak{g}$ . Let  $G_1 \subseteq G_2 \subseteq \dots \subseteq G_n \subseteq \dots \subseteq B(H)$  be a sequence of complex connected (finite dimensional) Lie subgroups of  $GL(H)$ . We assume that their Lie algebras  $\mathfrak{g}_n \subseteq B(H)$  are equipped with consistent Hermitian inner products, that is,  $(\cdot, \cdot)_{n+m}|_{\mathfrak{g}_n} = (\cdot, \cdot)_n$ , where  $(\cdot, \cdot)_n$  is the inner product on  $\mathfrak{g}_n$ . The corresponding norm is denoted by  $|\cdot|_n$ .

Let  $\mathfrak{g} = \bigcup_{n=1}^{\infty} \mathfrak{g}_n$  with the Hermitian inner product  $(\xi, \eta) = (\xi, \eta)_n$  for any  $\xi, \eta \in \mathfrak{g}_n$ .

We assume that  $|x| \geq \|x\|$  for any  $x \in \mathfrak{g}$ , where  $\|\cdot\|$  is the operator norm. Denote by  $\mathfrak{g}_\infty$  the closure of  $\mathfrak{g}$  in the norm  $|\cdot|$ , that is, all elements of finite norm. We also assume that the closure coincides with the completion of  $\mathfrak{g}$  with respect to the norm  $|\cdot|$ . Note that  $\mathfrak{g}_\infty$  is a subset of  $B(H)$ .

Let  $d$  denote the Riemannian metric  $d(y, z) = \inf_h \left\{ \int_0^1 |h^{-1}\dot{h}| ds \right\}$ , where  $h : [0, 1] \rightarrow GL(H)$  is a piecewise differentiable path,  $h(0) = y, h(1) = z, \dot{h} = \frac{dh}{ds}$ ,

$h' = h^{-1}\dot{h} \in \mathfrak{g}_\infty$ . Let  $G_\infty$  be the closure of  $\bigcup_{n=1}^{\infty} G_n$  in the Riemannian metric  $d_\infty = \inf_n d_n$ , where  $d_n$  is the Riemannian metric on  $G_n$ . Again we assume that the closure coincides with the completion. By  $\mathcal{H}^t(G_\infty)$  we denote a space of holomorphic functions on  $G_\infty$  with a certain direct limit-type norm  $\|\cdot\|_{t,\infty}$ . The precise definition is given in Section 4.

Let  $(1-D)_X^{-1}f = \sum_{k=0}^{\infty} (D^k f)(X)$  be the series of all derivatives of a function  $f$  on  $G_\infty$ . Then the Taylor map  $(1-D)_I^{-1}$  is an isometry from  $\mathcal{H}^t(G_\infty)$  into a subspace,  $J_t^0$ , of the dual of the tensor algebra of  $\mathfrak{g}_\infty$  equipped with the norm

$$|\alpha|_t^2 = \sum_{k=0}^{\infty} \frac{t^k}{k!} |\alpha_k|^2, \alpha = \sum_{k=0}^{\infty} \alpha_k, \alpha_k \in (\mathfrak{g}^{\otimes k})^*, k = 0, 1, 2, \dots, t > 0$$

See Notation 3.2 and Notation 4.2 for more detailed definitions. The following theorem will be proved in Section 4.

**Theorem.**  $\mathcal{H}^t(G_\infty)$  is a Hilbert space and  $(1-D)_I^{-1}$  is an isometry from  $\mathcal{H}^t(G_\infty)$  into  $J_t^0$ .

Moreover, in case the  $G_n$  are simply connected Theorem 4.5 gives the image of  $(1-D)_I^{-1}$ , which can be informally described as a completion of the universal enveloping algebra.

The heat kernel measure is constructed in Section 5 using stochastic differential equations on Hilbert spaces, in this case on the space of Hilbert-Schmidt operators. Denote by  $HS$  the space of Hilbert-Schmidt operators on  $H$  with the Hilbert-Schmidt (Hermitian) inner product  $\langle \cdot, \cdot \rangle_{HS}$  and corresponding real inner product  $\langle \cdot, \cdot \rangle_{HS} = \text{Re}(\cdot, \cdot)_{HS}$ . In most of the results of this paper we assume that  $G_n \subset I + HS$ ,  $\mathfrak{g}_n \subset HS$ . The following is a summary of results contained in Theorem 5.1 and Theorem 5.4

**Theorem.** Let  $W_t$  be the Wiener process in  $HS$  with covariance determined by the norm on  $\mathfrak{g}_\infty$ . Then the stochastic differential equation

$$\begin{aligned} dX_t &= X_t dW_t, \\ X_0 &= I \end{aligned}$$

has a unique solution in  $(I + HS) \cap GL(H)$ .

The transition probability of the process  $X_t$  determines the fundamental solution of the heat equation with the following informal Laplacian

$$(\Delta v)(X) = \frac{1}{2} \sum_{n=1}^{\infty} (\tilde{\xi}_n \tilde{\xi}_n v)(X),$$

where  $\{\xi_n\}_{n=1}^{\infty}$  is an orthonormal basis of  $\mathfrak{g}_\infty$  as a real space with the real inner product  $\langle \xi, \eta \rangle = \text{Re}(\xi, \eta)$  and  $(\tilde{\xi}_n v)(X) = \frac{d}{dt} \Big|_{t=0} v(Xe^{t\xi_n})$  for a function  $v : I + HS \rightarrow \mathbb{R}$ . The corresponding measure is called the heat kernel measure and the space of functions square integrable with respect to this measure is denoted by  $L^2(I + HS, \mu_t)$  and the norm by  $\|\cdot\|_t$ .

In addition to  $G_\infty$  we consider the Cameron-Martin subgroup.

**Definition 2.1.**  $G_{CM} = \{x \in B(H) : d(x, I) < \infty\}$  is called the Cameron-Martin subgroup.

Proposition 7.1 proves that  $G_{CM}$  is a group. Note that  $G_\infty \subset G_{CM}$  and the following theorem shows that under a condition on the Lie bracket they are actually equal. The next theorem will be proved in Section 7.

**Theorem.** *If  $\| [x, y] \| \leq C |x| |y|$  for all  $x, y \in \mathfrak{g}_\infty$ , then*

1.  $G_{CM} = G_\infty$ .
2. *The exponential map is a diffeomorphism from a neighborhood of 0 onto a neighborhood of  $I$  in  $G_{CM}$ .*

Note that under the condition of this theorem  $\mathfrak{g}_\infty$  is a Lie algebra.

Naturally defined holomorphic polynomials on  $I + HS$  play an important role in several results. One of them is that there is a natural isometry from the space of holomorphic polynomials  $\mathcal{HP}$  to  $\mathcal{H}^t(G_\infty)$ . To prove this we use approximations to the process  $Y_t + I$  discussed in Section 6. In addition, this isometry defines holomorphic skeletons on  $G_{CM}$  for the elements of the closure of  $\mathcal{HP}$  in  $L^2(I + HS, \mu_t)$ . This closure is denoted by  $\mathcal{HL}^2(I + HS, \mu_t)$ . The following results are contained in Theorem 8.7 and Theorem 8.5.

**Theorem.**

1.  $\mathcal{HP} \subset L^2(I + HS, \mu_t)$ .
2. *The identity  $\|f\|_{t,\infty} = \|f\|_t$  for any  $f \in \mathcal{HP}$  extends to an isometry  $\mathfrak{J}_{G_\infty}$  from  $\mathcal{HL}^2(I + HS, \mu_t)$  into  $\mathcal{H}^t(G_\infty)$ .*
3. *Suppose  $p_n \xrightarrow[n \rightarrow \infty]{L^2(I+HS, \mu_t)} f$ ,  $p_n \in \mathcal{HP}$ . Then there is a holomorphic function  $\tilde{f}$ , a skeleton of  $f$ , on  $G_{CM}$  such that  $p_n(x) \rightarrow \tilde{f}(x)$  for any  $x \in G_{CM}$ .*

We will denote the skeleton map by  $\mathfrak{J}_{G_{CM}}$ . Note that elements of  $\mathcal{HL}^2(I + HS, \mu_t)$  are defined up to a set of  $\mu_t$ -measure zero. Still the map  $\mathfrak{J}_{G_{CM}}$  gives a holomorphic version on  $G_{CM}$  of any element from  $\mathcal{HL}^2(I + HS, \mu_t)$ , even though  $G_{CM}$  itself might be of  $\mu_t$ -measure zero. In addition,  $\mathfrak{J}_{G_{CM}}f|_{G_\infty}$  is actually the isometry  $\mathfrak{J}_{G_\infty}$  from part 2 of the last theorem. This means that for holomorphic polynomials the skeleton map is the restriction map (to  $G_{CM}$ ).

Finally, Section 9 provides several examples to this abstract setting. One of the examples is the Hilbert-Schmidt complex orthogonal group which has been discussed in [7]. In addition we consider the Hilbert-Schmidt complex symplectic group and a group of infinite diagonal matrices.

For some  $\mathfrak{g}$  and natural norms  $|\cdot|$  on it, the Taylor map is an isometry between trivial spaces (see Section 9). But we show that for the natural condition on the norm  $|\cdot|$  considered in this paper, there are non-constant functions in  $\mathcal{H}^t(G_\infty)$ . Namely, this space contains all holomorphic polynomials. Indeed, Theorem 8.7 proves that the holomorphic polynomials are square integrable with respect to the heat kernel measure. Then in the same theorem we show that the  $L^2$ -norm of a polynomial is equal to the  $\mathcal{H}^t(G_\infty)$ -norm.

The following commutative diagram illustrates all the isometries described in this paper:

$$\begin{array}{ccccc}
 & & \mathcal{HL}^2(I + HS, \mu_t) & & \\
 & \swarrow \text{skeleton map} & & \searrow \mathfrak{J}_{G_\infty} & \\
 & \mathfrak{J}_{G_{CM}} & & & \\
 \mathcal{H}^t(G_{CM}) & \xrightarrow{\text{restriction map}} & \mathcal{H}^t(G_\infty) & \xrightarrow{\text{Taylor map}} & J_t^0
 \end{array}$$

## 3. ESTIMATES OF DERIVATIVES OF HOLOMORPHIC FUNCTIONS

Let  $\mu_t^n$  be the heat kernel measure on  $G_n$ ; let  $\mathcal{H}L^2(G_n, \mu_t^n)$  be the space of holomorphic functions square integrable with respect to  $\mu_t^n$ .

**Notation 3.1.**  $\|f\|_{L^2(G_n, \mu_t^n)} = \|f\|_{t,n}$ .

**Notation 3.2.** Suppose  $f$  is a function from  $G_{CM}$  to  $\mathbb{C}$ . Let  $(Df)(X)$  denote the unique element of  $\mathfrak{g}_\infty^*$  (as a complex space) such that

$$(Df)(X)(\xi) = (\tilde{\xi}f)(X) = \left. \frac{d}{dt} \right|_{t=0} f(Xe^{t\xi}), \quad \xi \in \mathfrak{g}_\infty, X \in G_\infty(G_{CM}),$$

if the derivative exists. Similarly  $(D^k f)(X)$  denotes the unique element of  $(\mathfrak{g}_\infty^{\otimes k})^*$  such that

$$(D^k f)(X)(\beta) = (\tilde{\beta}f)(X), \quad \beta \in \mathfrak{g}_\infty^{\otimes k}, X \in G_{CM}$$

and  $(D_n^k f)(X)$  denotes a unique element of  $(\mathfrak{g}_n^*)^{\otimes k} = (\mathfrak{g}_n^{\otimes k})^*$  such that

$$(D_n^k f)(X)(\beta) = (\tilde{\beta}f)(X), \quad \beta \in \mathfrak{g}_n^{\otimes k}, X \in G_n.$$

We will use the following notation:

$$(1 - D)^{-1}_X f = \sum_{k=0}^{\infty} (D^k f)(X) \quad \text{and} \quad (1 - D_n)^{-1}_X f = \sum_{k=0}^{\infty} (D_n^k f)(X).$$

The following estimate was proved by Driver and Gross in [5] for  $f \in \mathcal{H}L^2(G_n, \mu_t^n)$ :

$$|(\tilde{\beta}f)(g)|_{(\mathfrak{g}_n^{\otimes k})^*}^2 \leq \|f\|_{t,n}^2 \frac{k!|\beta|_n^2}{r^k} e^{|g|_n^2/s}, \text{ for } g \in G_n, r > 0, s + r \leq t, \beta \in \mathfrak{g}_n^{\otimes k},$$

where  $|(\tilde{\beta}f)(g)|_{(\mathfrak{g}_n^{\otimes k})^*}$  is  $((T_g G_n)^*)^{\otimes k}$ -norm (which can be identified with  $(\mathfrak{g}_n^*)^{\otimes k}$ ) and  $|g|_n = d_n(I, g)$ . We will need a slight modification of this estimate. Taking supremum over all  $\beta \in \mathfrak{g}_n^{\otimes k}, |\beta|_n = 1$ , we get

$$(3.1) \quad |(D_n^k f)(g)|_{(\mathfrak{g}_n^*)^{\otimes k}}^2 \leq \|f\|_{t,n}^2 \frac{k!}{r^k} e^{|g|_n^2/s},$$

where  $D_n^k$  is defined for  $G_n$  and  $\mathfrak{g}_n$  by Notation 3.2. Note that if  $\|f\|_{t,n}$  are uniformly bounded, then (3.1) gives a uniform bound, i.e. a bound independent of  $n$ . The following estimates can be proved for  $D_n^k$ :

**Lemma 3.3.** Let  $r > 0, q + r \leq t, X, Y \in G_n, f \in \mathcal{H}L^2(G_n, \mu_t^n)$ . Then

$$|(D_n^k f)(X) - (D_n^k f)(Y)|_{(\mathfrak{g}_n^{\otimes k})^*} \leq \|f\|_{t,n} K_{k+1,n} d_n(X, Y),$$

where  $K_{k,n} = K_{k,n}(X, Y) = \left(\frac{k!}{r^k}\right)^{1/2} e^{(|X|_n^2 + d_n(X, Y)^2)/q}$ .

*Proof.* Take  $h : [0, 1] \rightarrow G_n$  such that  $h(0) = X, h(1) = Y$ . Then by (3.1)

$$\begin{aligned} |(D_n^k f)(X) - (D_n^k f)(Y)|_{(\mathfrak{g}_n^{\otimes k})^*} &= \left| \int_0^1 \frac{d}{ds} (D_n^k f)(h(s)) ds \right|_{(\mathfrak{g}_n^{\otimes k})^*} \\ &\leq \int_0^1 \left| \frac{d}{ds} (D_n^k f)(h(s)) \right|_{(\mathfrak{g}_n^{\otimes k})^*} ds \leq \int_0^1 |D(D_n^k f)(h(s))(h^{-1}\dot{h})|_{(\mathfrak{g}_n^{\otimes k})^*} ds \end{aligned}$$

$$\begin{aligned}
& \leq \int_0^1 |D((D_n^k f)(h(s)))|_{(\mathfrak{g}_n^{\otimes k+1})^*} |h^{-1}\dot{h}|_n ds \\
& \leq \|f\|_{t,n} \left( \frac{(k+1)!}{r^{k+1}} \sup_{u \in [0,1]} e^{|h(u)|_n^2/q} \right)^{1/2} \int_0^1 |h^{-1}\dot{h}|_n ds \\
& \leq \|f\|_{t,n} \left( \frac{(k+1)!}{r^{k+1}} \right)^{1/2} \sup_{u \in [0,1]} e^{(|h(0)|_n^2 + d_n(h(0), h(u))^2)/q} \int_0^1 |h^{-1}\dot{h}|_n ds
\end{aligned}$$

Taking infimum over all such  $h$  we see that

$$|D_n^k f(X) - D_n^k f(Y)|_{(\mathfrak{g}_n^{\otimes k})^*} \leq \|f\|_{t,n} \left( \frac{(k+1)!}{r^{k+1}} \right)^{1/2} e^{(|X|_n^2 + d_n(X, Y)^2)/q} d_n(X, Y).$$

□

**Lemma 3.4.** *Let  $X \in G_n, f, g \in \mathcal{H}L^2(G_n, \mu_t^n), t > 0$ . Then*

$$|D_n^k f(X) - D_n^k g(X)|_{(\mathfrak{g}_n^{\otimes k})^*} \leq M_{k,n} \|f - g\|_{t,n},$$

where  $M_{k,n} = M_{k,n}(X, t) = \left( \frac{k!}{(t/2)^k} \right)^{1/2} e^{|X|_n^2/t}$ .

*Proof.* From (3.1) we have that for  $r > 0, q + r \leq t$

$$|D_n^k f(X) - D_n^k g(X)|_{(\mathfrak{g}_n^{\otimes k})^*}^2 \leq \|f - g\|_{t,n}^2 \frac{k!}{r^k} e^{|X|_n^2/q}.$$

Now take  $q = r = t/2$  to get what we claimed. □

**Lemma 3.5.** *Let  $X \in G_n, \xi \in \mathfrak{g}_n, f \in \mathcal{H}L^2(G_n, \mu_t^n)$ . Then there is a constant  $C = C(X, \xi, t, n) > 0$  such that*

$$\left| \frac{f(Xe^{u\xi}) - f(X)}{u} - (Df)(X)(\xi) \right| \leq \|f\|_{t,n} C u$$

for small enough  $u > 0$ .

*Proof.* Let  $h(s) = Xe^{s\xi}, 0 \leq s \leq u$ .

$$\begin{aligned}
\frac{f(Xe^{u\xi}) - f(X)}{u} - (Df)(X)(\xi) &= \frac{1}{u} \int_0^u \frac{d}{ds} f(h(s)) ds - (Df)(X)(\xi) \\
&= \frac{1}{u} \int_0^u \left( \frac{d}{ds} f(h(s)) - (Df)(X)(\xi) \right) ds = \frac{1}{u} \int_0^u ((Df)(h(s))(\xi) - (D_X f)(\xi)) ds.
\end{aligned}$$

Thus by Lemma 3.3 for any  $r > 0, q + r \leq t$

$$\begin{aligned}
\left| \frac{f(Xe^{u\xi}) - f(X)}{u} - (D_X f)(\xi) \right| &\leq \frac{1}{u} \int_0^u |((Df)(h(s))(\xi) - (D_X f)(\xi))| ds \\
&\leq \frac{1}{u} \int_0^u \|f\|_{t,n} K_{2,n}(h(s), X) d_n(h(s), X) |\xi|_n ds \\
&\leq \frac{1}{u} \int_0^u \|f\|_{t,n} \frac{\sqrt{2}}{r} e^{(|X|_n^2 + |\xi|_n^2 s^2)/q} s |\xi|_n^2 ds \\
&= \|f\|_{t,n} \frac{\sqrt{2}}{r} e^{|X|_n^2/q} |\xi|_n^2 \frac{1}{u} \int_0^u e^{(|\xi|_n^2/q) s^2} s ds \\
&= \|f\|_{t,n} \frac{q}{\sqrt{2}r} e^{|X|_n^2/q} \frac{e^{(|\xi|_n^2/q)u^2} - 1}{u} \leq \|f\|_{t,n} C u
\end{aligned}$$

for small  $u$ . □

**Theorem 3.6.** *Let  $f$  be a function on  $\cup_n G_n$ . Suppose that  $f|_{G_n}$  is holomorphic for any  $n$  and  $\sup \|f\|_{t,n} < \infty$ . Then  $f$  and all its derivatives have unique continuous extensions from  $\cup_n G_n$  to  $G_\infty$ .*

*Proof.* Take  $X \in G_\infty$ . We would like to define the extensions by

$$D^k f(X) = \lim_{n \rightarrow \infty} D^k f(X_n), \quad X_n \in G_n, \quad X_n \xrightarrow[n \rightarrow \infty]{d_\infty} X.$$

Let us check that the limit exists. Assume that  $l \leq n$ . By Lemma 3.3 for  $X_n, X_l \in G_n$

$$|D^k f(X_n) - D^k f(X_l)|_{(\mathfrak{g}_m^*)^{\otimes k}} \leq \|f\|_{t,n} K_{k+1,n}(X_n, X_l) d_n(X_n, X_l)$$

Note that  $d_m(X, Y)$  converges to  $d(X, Y)$  as  $m \rightarrow \infty$ . Now take the limit as  $m \rightarrow \infty$ , and then as  $n, l \rightarrow \infty$ . Thus  $D^k f(X_n)$  is a Cauchy sequence and therefore the limit exists. The uniqueness of the extension is easy to verify. □

**Remark 3.7.** The estimates in this section hold for the continuous extensions of  $D_n^k$  from  $\cup_n G_n$  to  $G_\infty$  if one takes the limit as  $n \rightarrow \infty$ . In general, the extensions of derivatives might not be derivatives of the extensions.

#### 4. ISOMETRIES

**Lemma 4.1.** *Suppose  $f|_{G_n} \in \mathcal{HL}^2(G_n, \mu_t^n)$  for all  $n$ . Then  $\|f\|_{t,n} \leq \|f\|_{t,n+1}$  for any  $n$ .*

*Proof.* First of all,  $\|f\|_{t,n}^2 = \|(1 - D_n)_I^{-1} f\|_{t,n}^2$  by the Driver-Gross isomorphism (see [5]), where  $\|(1 - D_n)_I^{-1} f\|_{t,n}^2 = \sum_{k=0}^{\infty} |(D_n^k f)(I)|_{t,n}^2$ . Take an orthonormal basis

$\{\eta_l\}_{l=1}^{d_{n+1}}$  of the complex inner product space  $\mathfrak{g}_{n+1}$  such that  $\{\eta_l\}_{l=1}^{d_n}$  is an orthonormal basis of  $\mathfrak{g}_n$ . Then

$$\begin{aligned} |(D_{n+1}^k f)(I)|_{t,n+1}^2 &= \sum_{1 \leq i_m \leq d_{n+1}} |\tilde{\eta}_{i_1} \dots \tilde{\eta}_{i_k} f(I)|^2 \\ &\geq \sum_{1 \leq i_m \leq d_n} |\tilde{\eta}_{i_1} \dots \tilde{\eta}_{i_k} f(I)|^2 = |(D_n^k f)(I)|_{t,n}^2. \end{aligned}$$

Therefore  $\|(1 - D_n)_I^{-1} f\|_{t,n}^2 \leq \|(1 - D_{n+1})_I^{-1} f\|_{t,n+1}^2$ , so the claim holds.  $\square$

**Notation 4.2.** Let  $\mathfrak{h}$  be a complex Lie algebra with a Hermitian inner product on it. Then  $T(\mathfrak{h})$  will denote the algebraic tensor algebra over  $\mathfrak{h}$  as a complex vector space and  $T'(\mathfrak{h})$  will denote the algebraic dual of  $T(\mathfrak{h})$ . Define a norm on  $T(\mathfrak{h})$  by

$$(4.1) \quad |\beta|_t^2 = \sum_{k=0}^n \frac{k!}{t^k} |\beta_k|^2, \beta = \sum_{k=0}^n \beta_k, \beta_k \in \mathfrak{h}^{\otimes k}, k = 0, 1, 2, \dots, t > 0.$$

Here  $|\beta_k|$  is the cross norm on  $\mathfrak{h}^{\otimes k}$  arising from the inner product on  $\mathfrak{h}^{\otimes k}$  determined by the norm  $|\cdot|$  on  $\mathfrak{h}$ .  $T_t(\mathfrak{h})$  will denote the completion of  $T(\mathfrak{h})$  in this norm. The topological dual of  $T_t(\mathfrak{h})$  may be identified with the subspace  $T_t^*(\mathfrak{h})$  of  $T'(\mathfrak{h})$  consisting of such  $\alpha \in T'(\mathfrak{h})$  that the norm

$$(4.2) \quad |\alpha|_t^2 = \sum_{k=0}^{\infty} \frac{t^k}{k!} |\alpha_k|^2, \alpha = \sum_{k=0}^{\infty} \alpha_k, \alpha_k \in (\mathfrak{h}^{\otimes k})^*, k = 0, 1, 2, \dots, t > 0$$

is finite. Here  $|\alpha_k|$  is the norm on  $(\mathfrak{h}^{\otimes k})^*$  dual to the cross norm on  $\mathfrak{h}^{\otimes k}$ .

There is a natural pairing for any  $\alpha \in T'(\mathfrak{h})$  and  $\beta \in T(\mathfrak{h})$  denoted by

$$\langle \alpha, \beta \rangle = \sum_{k=0}^{\infty} \langle \alpha_k, \beta_k \rangle, \alpha = \sum_{k=0}^{\infty} \alpha_k, \beta = \sum_{k=0}^n \beta_k, \alpha_k \in (\mathfrak{h}^{\otimes k})^*, \beta_k \in \mathfrak{h}^{\otimes k}, k = 0, 1, 2, \dots$$

Denote by  $J(\mathfrak{h})$  the two-sided ideal in  $T(\mathfrak{h})$  generated by  $\{\xi \otimes \eta - \eta \otimes \xi - [\xi, \eta], \xi, \eta \in \mathfrak{h}\}$ . Let  $J^0(\mathfrak{h}) = \{\alpha \in T'(\mathfrak{h}) : \alpha(J) = 0\}$ . Finally, let  $J_t^0(\mathfrak{h}) = T_t^*(\mathfrak{h}) \cap J^0(\mathfrak{h})$ . We will denote  $J_t^0(\mathfrak{g})$  by  $J_t^0$ ,  $J_t^0(\mathfrak{g}_n)$  by  $J_{t,n}^0$ .

The coefficients  $\frac{k!}{t^k}$  in the norm on the tensor algebra  $T(\mathfrak{h})$  are related to the heat kernel.

By Theorem 3.6 all  $D_n^k$  have continuous extensions from  $\cup_n G_n$  to  $G_\infty$ , which allows us to introduce the following definition.

**Definition 4.3.**  $\mathcal{H}^t(G_\infty)$  is a space of continuous functions on  $G_\infty$  such that their restrictions to  $G_n$  are holomorphic for every  $n$  and  $\|f\|_{t,\infty} = \sup_n \{\|f\|_{t,n}\} =$

$$\lim_{n \rightarrow \infty} \|f\|_{t,n} < \infty.$$

**Theorem 4.4.**  $\mathcal{H}^t(G_\infty)$  is a Hilbert space and  $(1 - D)_I^{-1}$  is an isometry from  $\mathcal{H}^t(G_\infty)$  into  $J_t^0$ .

*Proof.* First let us show that  $(1 - D)_I^{-1}$  is an isometry.  $T_n$  is a subalgebra of  $T$ . Note that  $T'_n$  can be easily identified with a subspace of  $T'$ . Namely, for any  $\alpha_n \in T'_n$  we can define  $\alpha$  as follows:

$$\alpha = \begin{cases} \alpha_n & \text{on } T_n \\ 0 & \text{on } T_n^\perp. \end{cases}$$



Therefore  $T'_n = (T_n^\perp)^0$ . Define  $\Pi_n$  to be an orthogonal projection from  $T$  to  $T_n$ . Let  $\Pi'_n$  denote the following map from  $T'$  to  $T'_n$ :  $(\Pi'_n \alpha)(x) = \alpha(\Pi_n x)$ ,  $\alpha \in T'$ ,  $x \in T$ . Then  $\Pi'_n \circ (1 - D)_I^{-1} : \mathcal{H}L^2(G_\infty) \rightarrow T'_n$  is equal to  $(1 - D_n)_I^{-1}$ . Indeed, note that if we choose an orthonormal basis of the complex inner product space  $\mathfrak{g}$  such that  $\{\eta_m\}_{m=1}^{d_n}$  is an orthonormal basis of  $\mathfrak{g}_n$ , then  $\Pi_n$  can be described explicitly

$$\Pi_n(\eta_{k_1} \otimes \eta_{k_2} \otimes \dots \otimes \eta_{k_l}) = \begin{cases} 0 & \text{if } k_s > d_n \text{ for some } 1 \leq s \leq l \\ \eta_{k_1} \otimes \eta_{k_2} \otimes \dots \otimes \eta_{k_l} & \text{otherwise} \end{cases}$$

$$\begin{aligned} \text{and } \langle \Pi'_n \circ (1 - D)_I^{-1} f, \eta_{k_1} \otimes \eta_{k_2} \otimes \dots \otimes \eta_{k_l} \rangle &= \langle (1 - D)_I^{-1} f, \Pi_n(\eta_{k_1} \otimes \eta_{k_2} \otimes \dots \otimes \eta_{k_l}) \rangle \\ &= \begin{cases} 0 & \text{if } k_s > d_n \text{ for some } 1 \leq s \leq l \\ \langle (1 - D)_I^{-1} f, \eta_{k_1} \otimes \eta_{k_2} \otimes \dots \otimes \eta_{k_l} \rangle & \text{otherwise} \end{cases} \end{aligned}$$

So  $\Pi'_n \circ (1 - D)_I^{-1} = (1 - D_n)_I^{-1}$ . Driver and Gross proved in [5] that  $\Pi'_n \circ (1 - D)_I^{-1}$  is an isometry from  $\mathcal{H}L^2(G_n, \mu_t^n)$  into  $J_{t,n}^0$ . Let us define a restriction map  $R_n : \mathcal{H}L^2(G_\infty) \rightarrow \mathcal{H}L^2(G_n, \mu_t^n)$  by  $f \mapsto f|_{G_n}$ . Thus we have a commutative diagram:

$$(4.3) \quad \begin{array}{ccc} \mathcal{H}L^2(G_\infty) & \xrightarrow{(1-D)_I^{-1}} & J_t^0 \\ R_n \downarrow & & \downarrow \Pi'_n \\ \mathcal{H}L^2(G_n, \mu_t^n) & \xrightarrow[\text{Driver-Gross}]{(1-D_n)_I^{-1}} & J_{t,n}^0 \end{array}$$

Now we can prove that  $(1 - D)_I^{-1}$  is an isometry. By Lemma 4.1

$$\|f\|_{t,\infty} = \lim_{n \rightarrow \infty} \|f\|_{t,n} = \lim_{n \rightarrow \infty} \|R_n f\|_{t,n}.$$

It is clear that  $|\Pi'_n \alpha|_t = |\Pi'_n \alpha|_{t,n} \xrightarrow{n \rightarrow \infty} |\alpha|_t$  for any  $\alpha \in T'$ . In particular,  $|\Pi'_n \circ (1 - D)_I^{-1} f|_{t,n} \xrightarrow{n \rightarrow \infty} |(1 - D)_I^{-1} f|_t$ . At the same time  $|\Pi'_n \circ (1 - D)_I^{-1} f|_{t,n} = |(1 - D_n)_I^{-1} \circ R_n f|_{t,n} = \|R_n f\|_{t,n} \xrightarrow{n \rightarrow \infty} \|f\|_{t,\infty}$  by the Driver-Gross isomorphism.

Now let us check that  $\mathcal{H}^t(G_\infty)$  is a Hilbert space. It is clear that  $\|\cdot\|_{t,\infty}$  is a seminorm. Suppose that  $\|f\|_{t,\infty} = 0$ . Then  $\|f\|_{t,n} = 0$  for any  $n, t > 0$ . We know that  $\mathcal{H}L^2(G_n, \mu_t^n)$  is a Hilbert space, therefore  $f|_{G_n} = 0$  for all  $n$ . By Lemma 3.6 we have that  $f|_{G_\infty} = 0$ . Thus  $\|\cdot\|_{t,\infty}$  is a norm.

Let us now show that  $\mathcal{H}^t(G_\infty)$  is a complete space. Suppose  $\{f_m\}_{m=1}^\infty$  is a Cauchy sequence in  $\mathcal{H}^t(G_\infty)$ . Then  $\{f_m|_{G_n}\}_{m=1}^\infty$  is a Cauchy sequence in  $\mathcal{H}L^2(G_n, \mu_t^n)$  for all  $n$ . Therefore there exists  $g_n$  from  $\mathcal{H}L^2(G_n, \mu_t^n)$  such that  $f_m|_{G_n} \xrightarrow{m \rightarrow \infty} g_n$ . Note that  $g_{n+m}|_{G_n} = g_n$ . In addition,

$$\|g_n\|_{t,n} \leq \|f_m|_{G_n}\|_{t,n} + \|f_m|_{G_n} - g_n\|_{t,n} \leq \|f_m\|_{t,\infty} + \|f_m|_{G_n} - g_n\|_{t,n}.$$

Note that  $\{\|f_m\|_{t,\infty}\}_{m=1}^\infty$  is again a Cauchy sequence, so it has a (finite) limit as  $m \rightarrow \infty$ . Taking a limit in (4) as  $m \rightarrow \infty$  we get that  $\{\|g_n\|_{t,n}\}_{n=1}^\infty$  are uniformly bounded.

By Lemma 3.6 there exists a continuous function  $g$  on  $G_\infty$  such that  $g|_{G_n} = g_n$ . Thus  $g \in \mathcal{H}^t(G_\infty)$ . Now we need to prove that  $f_m \xrightarrow{m \rightarrow \infty} g$  in  $\mathcal{H}^t(G_\infty)$ . Let  $\alpha = (1 - D)_I^{-1} g$ ,  $\alpha_m = (1 - D)_I^{-1} f_m$ . As we have shown  $(1 - D)_I^{-1}$  is an isometry, so  $\alpha_m$  is a Cauchy sequence in  $J_t^0$ . Thus there exists  $\alpha' \in J_t^0$  such that  $\alpha_m \xrightarrow{m \rightarrow \infty} \alpha'$

(in  $J_t^0$ ). The question is whether  $\alpha = \alpha'$ . Note that

$$\Pi'_n \alpha = (1 - D_n)_I^{-1} g \quad \text{and} \quad \Pi'_n \alpha_m = (1 - D_n)_I^{-1} f_m.$$

We know that  $\mathcal{H}L^2(G_n, \mu_t^n)$  is a Hilbert space, therefore  $\Pi'_n \alpha = \Pi'_n \alpha'$  for any  $n$ . Thus  $\alpha = \alpha'$ , which completes the proof.  $\square$

**Theorem 4.5.** *Suppose connected (finite dimensional) Lie groups  $G_n$  are simply connected. Then the map  $(1 - D)_I^{-1}$  is surjective onto  $J_t^0$ .*

*Proof.* Indeed, for any  $\alpha \in J_t^0$ ,  $\Pi'_n \alpha \in J_{t,n}^0$  by the Driver-Gross isomorphism there exists a unique  $f_n \in \mathcal{H}L^2(G_n, \mu_t^n)$  such that  $(1 - D)_{I,n}^{-1} f_n = \Pi'_n \alpha$ . Moreover,  $R_n f_{n+1} = f_n$  by the commutativity of diagram 4.3 and uniqueness of functions  $f_n$ . Indeed, from the diagram we have  $(1 - D_n)_I^{-1} (R_n f_{n+1}) = \Pi'_n \circ (1 - D)_I^{-1} f_{n+1} = (\Pi'_n \circ \Pi'_{n+1}) \circ (1 - D)_I^{-1} f_{n+1} = \Pi'_n \circ (\Pi'_{n+1} \circ (1 - D)_I^{-1}) f_{n+1} = \Pi'_n \alpha$  by the definition of  $f_{n+1}$ . Thus we can define  $f$  on  $\cup_n G_n$  as follows:  $f|_{G_n} = f_n$ . By Theorem 3.6 there is a unique continuous function  $g$  on  $G_\infty$  such that  $g|_{G_\infty} = f$ . Note that  $\|f\|_{t,n} = \|f_n\|_{t,n} = |\alpha|_t$ , therefore by Proposition 4.1  $\|g\|_{t,\infty} < \infty$  and so  $g \in \mathcal{H}L^2(G_\infty)$ .  $\square$

Here we should note that sometimes  $(1 - D)_I^{-1}$  is an isometry onto a trivial space. Indeed, in [7] we proved the following theorem

**Theorem 4.6.** *Suppose  $\mathfrak{g}$  is a Lie algebra with an inner product  $\langle \cdot, \cdot \rangle$ . Assume that there is an orthonormal basis  $\{\xi_k\}_{k=1}^\infty$  of  $\mathfrak{g}$  such that for any  $k$  there are nonzero  $\alpha_k \in \mathbb{C}$  and an infinite set of distinct pairs  $(i_m, j_m)$  satisfying  $\xi_k = \alpha_k [\xi_{i_m}, \xi_{j_m}]$ . Then  $J_t^0$  is isomorphic to  $\mathbb{C}$ .*

In particular, the conclusion of this theorem holds for a Lie algebra of the Hilbert-Schmidt complex orthogonal group  $SO_{HS}$  and a Lie algebra of the Hilbert-Schmidt complex symplectic group  $Sp_{HS}$  with invariant inner product (see Section 9). One of the ways to deal with this problem is to show that there are non constant functions in  $\mathcal{H}^t(G_\infty)$ . It will be done in Section 8, but before we can manage this we need to construct the heat kernel measure.

## 5. THE HEAT KERNEL MEASURE

Suppose  $G_n \subset I + HS$ ,  $\mathfrak{g}_n \subset HS$  and  $|\cdot| \geq |\cdot|_{HS}$ . Then  $\mathfrak{g}_\infty$  is embedded in  $HS$ . There is a quadratic functional  $\tilde{Q}$  on  $\mathfrak{g}_\infty$  defined by  $\tilde{Q}(x) = \|x\|_{HS}^2$ . Thus there exists a positive operator  $Q : \mathfrak{g}_\infty \rightarrow \mathfrak{g}_\infty$  such that

$$(x, Qx)_{\mathfrak{g}_\infty} = \tilde{Q}(x) = \|x\|_{HS}^2 = (x, x)_{HS}.$$

Note that  $Q$  is a bounded complex linear operator. The operator  $Q$  will be identified with its nonnegative extension to  $HS$  by  $Q|_{(\mathfrak{g}_\infty)^\perp_{HS}} = 0$ . We assume that  $Q$  is a trace class operator on  $HS$  and that all  $\mathfrak{g}_n$  are its invariant subspaces.

We begin with the definition of the process  $Y_t$ . Let  $W_t$  be a  $\mathfrak{g}_\infty$ -valued Wiener process with a covariance operator  $Q : HS \rightarrow \mathfrak{g}_\infty$ .

In what follows let  $\{\xi_n\}_{n=1}^\infty$  denote an orthonormal basis of  $\mathfrak{g}_\infty$  as a real space such that  $\{\xi_n\}_{n=1}^{2d_n}$  is an orthonormal basis of  $\mathfrak{g}_n$ . Here  $d_n = \dim_{\mathbb{C}} \mathfrak{g}_n$ , the complex dimension of  $\mathfrak{g}_n$ . Then  $\{e_n\} = \{Q^{-1/2} \xi_n\}_{n=1}^\infty$  is an orthonormal basis of the  $Span_{HS}(\mathfrak{g}_\infty)$ .

Denote by  $L_2^0 = L_2(\mathfrak{g}_\infty, HS)$  the space of the Hilbert-Schmidt operators from  $\mathfrak{g}_\infty$  to  $HS$  with the (Hilbert-Schmidt) norm  $\|\Psi\|_{L_2^0}^2$ . Let  $B : HS \rightarrow L_2^0$ ,  $B(Y)U = (Y + I)U$  for  $U \in \mathfrak{g}_\infty$ ; then the following theorem holds.

**Theorem 5.1.** 1. *The stochastic differential equation*

$$(5.1) \quad \begin{aligned} dY_t &= B(Y_t)dW_t, \\ Y_0 &= 0 \end{aligned}$$

*has a unique solution, up to equivalence, among the processes satisfying*

$$\mathbf{P} \left( \int_0^T \|Y_s\|_{HS}^2 ds < \infty \right) = 1.$$

2. *For any  $p \geq 2$  there exists a constant  $C_{p,T} > 0$  such that*

$$\sup_{t \in [0, T]} \mathbf{E} \|Y_t\|_{HS}^p \leq C_{p,T}.$$

*Proof of Theorem 5.1.* To prove this theorem we will use Theorem 7.4 from the book by DaPrato and Zabczyk [3], p.186. It is enough to check that

1.  $B(Y)$  is a measurable mapping from  $HS$  to  $L_2^0$ .
2.  $\|B(Y_1) - B(Y_2)\|_{L_2^0} \leq C\|Y_1 - Y_2\|_{HS}$  for any  $Y_1, Y_2 \in HS$ .
3.  $\|B(Y)\|_{L_2^0}^2 \leq K(1 + \|Y\|_{HS}^2)$  for any  $Y \in HS$ .

*Proof of 1.* We want to check that  $B(Y)$  is in  $L_2^0$  for any  $Y$  from  $HS$ . First of all,  $B(Y)U \in HS$ , for any  $U \in \mathfrak{g}_\infty$ . Indeed,  $B(Y)U = (Y + I)U = YU + U \in HS$ , since  $U$  and  $Y$  are in  $HS$ .

Now let us verify that  $B(Y) \in L_2^0$ . Consider the Hilbert-Schmidt norm of  $B$  as an operator from  $\mathfrak{g}_\infty$  to  $HS$ . Then the Hilbert-Schmidt norm of  $B$  can be found as follows

$$\begin{aligned} \|B(Y)\|_{L_2^0}^2 &= \sum_{n=1}^{\infty} \langle B(Y)\xi_n, B(Y)\xi_n \rangle_{HS} = \sum_{n=1}^{\infty} \langle (Y + I)\xi_n, (Y + I)\xi_n \rangle_{HS} \\ &\leq \|Y + I\|^2 \sum_{n=1}^{\infty} \langle \xi_n, \xi_n \rangle_{HS} = \|Y + I\|^2 \sum_{n=1}^{\infty} \langle e_n, Qe_n \rangle_{HS} = \|Y + I\|^2 \text{Tr}Q < \infty, \end{aligned}$$

since the operator norm  $\|Y + I\|$  is finite.

*Proof of 2.* Similarly to the previous proof we have

$$\|B(Y_1) - B(Y_2)\|_{L_2^0} \leq \|Y_1 - Y_2\|_{HS} (\text{Tr}Q)^{1/2}.$$

*Proof of 3.* Use the estimate in the proof of 1

$$\begin{aligned} \|B(Y)\|_{L_2^0} &\leq (\text{Tr}Q)^{1/2} \|Y + I\| \leq (\text{Tr}Q)^{1/2} (\|Y\|_{HS} + 1), \text{ so} \\ \|B(Y)\|_{L_2^0}^2 &\leq 2(\text{Tr}Q)(1 + \|Y\|_{HS}^2). \end{aligned} \quad \square$$

**Lemma 5.2.**  $\sum_{n=1}^{\infty} \xi_n^2 = 0$ .

*Proof.* First let us check that  $\sum_{n=1}^{\infty} \xi_n^2$  does not depend on the choice of the orthonormal basis  $\{\xi_n\}_1^\infty$  in  $\mathfrak{g}_\infty$ . Define a bilinear real form on  $HS \times HS$  by  $L(f, g) = \Lambda(Q^{1/2}fQ^{1/2}g)$ , where  $\Lambda$  is a real bounded linear functional on  $HS$ . Then  $f \mapsto L(f, g)$  is a bounded linear functional on  $HS$  and so  $L(f, g) = \langle f, \tilde{g} \rangle_{HS}$  for some  $\tilde{g} \in HS$ . There exists a linear operator  $B$  on  $HS$  such that  $L(f, g) = \langle f, Bg \rangle_{HS}$ .

We can actually compute  $B$ . Indeed, there exists an element  $h \in HS$  such that  $\Lambda(x) = \langle x, h \rangle_{HS}$ . Then

$$L(f, g) = \langle Q^{1/2} f Q^{1/2} g, h \rangle_{HS}$$

and so

$$L(f, g) = Tr \left( Q^{1/2} f Q^{1/2} g h^* \right) = \langle Q^{1/2} f, h(Q^{1/2} g)^* \rangle_{HS} = \langle f, Q^{1/2} (h(Q^{1/2} g)^*) \rangle_{HS}.$$

Thus  $Bg = Q^{1/2} (h(Q^{1/2} g)^*)$ . for some  $h \in HS$ . Therefore  $B$  is trace class and since trace is independent of a basis

$$Tr B = \sum_{n=1}^{\infty} \langle e_n, B e_n \rangle_{HS} = \sum_{n: e_n \in \mathfrak{g}_{\infty}} \Lambda(Q^{1/2} e_n Q^{1/2} e_n) = \Lambda \left( \sum_{n=1}^{\infty} \xi_n^2 \right)$$

does not depend on the choice of  $\{\xi_n\}_{n=1}^{\infty}$  for any  $\Lambda$ . Now choose  $\{\xi_n\}_{n=1}^{\infty}$  so that  $\xi_{2n} = i\xi_{2n-1}$  for  $n = 1, 2, \dots$ . Here  $i = \sqrt{-1}$ . Then  $(\xi_{2n-1})^2 + (\xi_{2n})^2 = 0$ .  $\square$

**Remark 5.3.** In case when  $\mathfrak{g}_{\infty}$  is a real space without complex structure, the process is a solution of the equation

$$dY_t = B(Y_t) dW_t + \sum_{n=1}^{\infty} \xi_n^2 (Y_t + I) dt,$$

$$Y_0 = 0$$

The  $dt$  term is necessary to ensure that the generator of  $Y_t$  is the Laplacian.

**Theorem 5.4.**  $Y_t + I \in GL(H)$  for any  $t > 0$  with probability 1. The inverse of  $Y_t + I$  is  $\tilde{Y}_t + I$ , where  $\tilde{Y}_t$  is a solution to the stochastic differential equation

$$(5.2) \quad \begin{aligned} d\tilde{Y}_t &= \tilde{B}(\tilde{Y}_t) dW_t, \\ \tilde{Y}_0 &= 0, \end{aligned}$$

where  $\tilde{B}(Y)(U) = -U(Y + I), U \in HS$ .

*Proof of Theorem 5.4.* First we will check that  $(\tilde{Y}_t + I)(Y_t + I) = I$  with probability 1 for any  $t > 0$ . To do this we will apply Itô's formula to  $G(Y_t, \tilde{Y}_t)$ , where  $G$  is defined as follows:  $G(\mathbf{Y}) = \Lambda((Y_1 + I)(Y_2 + I))$ , where  $\mathbf{Y} = (Y_1, Y_2) \in HS \times HS$  and  $\Lambda$  is a nonzero linear real bounded functional from  $HS \times HS$  to  $\mathbb{R}$ . Here we view  $G$  as a function on a Hilbert space  $HS \times HS$  and consider  $G(\mathbf{Y}_t) = G(Y_t, \tilde{Y}_t)$ . Then  $(Y_t + I)(\tilde{Y}_t + I) = I$  if and only if  $\Lambda((Y_t + I)(\tilde{Y}_t + I) - I) = 0$  for any  $\Lambda$ . In order to use Itô's formula we must verify several properties of the process  $\tilde{Y}_t$  and the mapping  $G$ :

1.  $\tilde{B}(\tilde{Y}_s)$  is an  $L_2^0$ -valued process stochastically integrable on  $[0, T]$
2.  $G$  and the derivatives  $G_t, G_{\mathbf{Y}}, G_{\mathbf{Y}\mathbf{Y}}$  are uniformly continuous on bounded subsets of  $[0, T] \times HS \times HS$ .

*Proof of 1.* See 1 in the proof of Theorem 5.1.

*Proof of 2.* Let us calculate  $G_t, G_{\mathbf{Y}}, G_{\mathbf{Y}\mathbf{Y}}$ . First,  $G_t = 0$ . For any  $\mathbf{S} \in HS \times HS$

$$G_{\mathbf{Y}}(\mathbf{Y})(\mathbf{S}) = \Lambda(S_1(Y_2 + I) + (Y_1 + I)S_2),$$

For any  $\mathbf{S}, \mathbf{T} \in HS \times HS$

$$G_{\mathbf{Y}\mathbf{Y}}(\mathbf{Y})(\mathbf{S} \otimes \mathbf{T}) = \Lambda(S_1 T_2 + T_1 S_2).$$

Thus condition 2 is satisfied.

We will use the notation:

$$\begin{aligned} G_{\mathbf{Y}}(\mathbf{Y})(\mathbf{S}) &= \langle \bar{G}_{\mathbf{Y}}(\mathbf{Y}), \mathbf{S} \rangle_{HS}, \\ G_{\mathbf{Y}\mathbf{Y}}(\mathbf{Y})(\mathbf{S} \otimes \mathbf{T}) &= \langle \bar{G}_{\mathbf{Y}\mathbf{Y}}(\mathbf{Y})\mathbf{S}, \mathbf{T} \rangle_{HS}, \end{aligned}$$

where  $\bar{G}_{\mathbf{Y}}$  is an element of  $HS \times HS$  and  $\bar{G}_{\mathbf{Y}\mathbf{Y}}$  is an operator on  $HS \times HS$  corresponding to the functionals  $G_{\mathbf{Y}} \in (HS \times HS)^*$  and  $G_{\mathbf{Y}\mathbf{Y}} \in ((HS \times HS) \otimes (HS \times HS))^*$ .

Now we can apply Itô's formula to  $G(\mathbf{Y}_t)$ :

$$(5.3) \quad \begin{aligned} G(\mathbf{Y}_t) &= \int_0^t \langle \bar{G}_{\mathbf{Y}}(\mathbf{Y}_s), (B(Y_s)dW_s, \tilde{B}(\tilde{Y}_s)dW_s) \rangle_{HS} \\ &\quad + \int_0^t \frac{1}{2} Tr_{HS}[\bar{G}_{\mathbf{Y}\mathbf{Y}}(\mathbf{Y}_s)(B(Y_s)Q^{1/2}, (\tilde{B}(\tilde{Y}_s)Q^{1/2})^*)] ds. \end{aligned}$$

Let us calculate the two integrands in (5.3) separately. The first integrand is

$$\begin{aligned} &\langle \bar{G}_{\mathbf{Y}}(\mathbf{Y}_s), (B(Y_s)dW_s, \tilde{B}(\tilde{Y}_s)dW_s) \rangle_{HS} \\ &= G_{\mathbf{Y}}(\mathbf{Y}_s)((Y_s + I)dW_s, -dW_s(\tilde{Y}_s + I)) \\ &= \Lambda((Y_s + I)dW_s(\tilde{Y}_s + I) - (Y_s + I)dW_s(\tilde{Y}_s + I)) = 0. \end{aligned}$$

The second integrand is

$$\begin{aligned} &\frac{1}{2} Tr_{HS}[\bar{G}_{\mathbf{Y}\mathbf{Y}}(\mathbf{Y}_s) \left( (B(Y_s)Q^{1/2}, \tilde{B}(\tilde{Y}_s)Q^{1/2})(B(Y_s)Q^{1/2}, \tilde{B}(\tilde{Y}_s)Q^{1/2})^* \right)] \\ &= \frac{1}{2} \sum_{n=1}^{\infty} G_{\mathbf{Y}\mathbf{Y}}(\mathbf{Y}_s) \left( (B(Y_s)Q^{1/2}e_n, \tilde{B}(\tilde{Y}_s)Q^{1/2}e_n) \otimes (B(Y_s)Q^{1/2}e_n, \tilde{B}(\tilde{Y}_s)Q^{1/2}e_n) \right) \\ &= \frac{1}{2} \sum_{n=1}^{\infty} \Lambda \left( -(Y_s + I)\xi_n^2(\tilde{Y}_s + I) - (Y_s + I)\xi_n^2(\tilde{Y}_s + I) \right) \\ &= \sum_{n=1}^{\infty} \Lambda \left( -(Y_s + I)\xi_n^2(\tilde{Y}_s + I) \right) = 0 \end{aligned}$$

by Lemma 5.2. This shows that the stochastic differential of  $G$  is zero, so  $G(\mathbf{Y}_t) = I$  for any  $t > 0$ . By the Fredholm alternative  $Y_t + I$  has an inverse; therefore it has to be  $(\tilde{Y}_t + I)$ .  $\square$

In some cases we can prove that the process  $Y_t + I$  lives in a smaller group. For example, it can be shown if the group is defined by a relatively simple relation (see Section 9).

Let us define  $\mu_t$  as

$$\int_{I+HS} f(X)\mu_t(dX) = Ef(X_t(I)) = P_{t,0}f(I)$$

for any bounded Borel function  $f$  on  $I + HS$ .

**Definition 5.5.**  $\mu_t$  is called *the heat kernel measure on  $G_\infty$* . The space of all square integrable functions on  $I + HS$  is denoted by  $L^2(I + HS, \mu_t)$  and the corresponding norm by  $\|f\|_{L^2(I + HS, \mu_t)} = \|f\|_t$ .

A motivation for such a name for  $\mu_t$  is the fact that Kolmogorov's backward equation corresponding to the process  $Y_t + I$  is the heat equation in a sense. First of all, the coefficient  $B$  depends only on  $Y \in HS$ ; therefore  $P_{s,t}f(Y) = Ef(Y(t, s; Y)) = P_{t-s}f(Y)$ . According to Theorem 9.16 from [3] for any  $\varphi \in C_b^2(HS)$  and  $Y \in HS$  the function  $v(t, Y) = P_t\varphi(Y)$  is a unique strict solution from  $C_b^{1,2}(HS)$  for the parabolic type equation (Kolmogorov's backward equation)

$$(5.4) \quad \begin{aligned} \frac{\partial}{\partial t}v(t, Y) &= \frac{1}{2}Tr[v_{YY}(t, Y)(B(Y)Q^{1/2})(B(Y)Q^{1/2})^*] \\ v(0, Y) &= \varphi(Y), t > 0, Y \in HS. \end{aligned}$$

Here  $C_b^n(HS)$  denotes the space of all functions from  $HS$  to  $\mathbb{R}$  that are  $n$ -times continuously Frechet differentiable with all derivatives up to order  $n$  bounded and  $C_b^{k,n}(HS)$  denotes the space of all functions from  $[0, T] \times HS$  to  $\mathbb{R}$  that are  $k$ -times continuously Frechet differentiable with respect to  $t$  and  $n$ -times continuously Frechet differentiable with respect to  $Y$  with all partial derivatives continuous in  $[0, T] \times HS$  and bounded.

Let us rewrite Equation (5.4) as the heat equation. First

$$\begin{aligned} &Tr[v_{YY}(t, Y)(B(Y)Q^{1/2})(B(Y)Q^{1/2})^*] \\ &= \sum_{n=1}^{\infty} v_{YY}(t, Y)(B(Y)Q^{1/2}e_n \otimes B(Y)Q^{1/2}e_n) \\ &= \sum_{n=1}^{\infty} v_{YY}(t, Y)((Y + I)Q^{1/2}e_n \otimes (Y + I)Q^{1/2}e_n) \\ &= \sum_{n=1}^{\infty} v_{YY}(t, Y)((Y + I)\xi_n \otimes (Y + I)\xi_n), \end{aligned}$$

where  $v_{YY}(t, Y)$  is viewed as a functional on  $HS \otimes HS$ . Now change  $Y$  to  $X - I$ . Then for any smooth bounded function  $\varphi(X) : I + HS \rightarrow \mathbb{R}$ , the function  $v(t, X) = P_t\varphi(X)$  satisfies this equation, which can be considered as the heat equation

$$(5.5) \quad \begin{aligned} \frac{\partial}{\partial t}v(t, X) &= L_1v(t, X) \\ v(0, X) &= \varphi(X), t > 0, X \in I + HS, \end{aligned}$$

where the differential operator  $L_1$  on the space  $C_b^{1,2}(I + HS)$  is defined by

$$L_1v \stackrel{\text{def}}{=} \frac{1}{2} \sum_{n=1}^{\infty} v_{YY}(t, Y)((Y + I)\xi_n \otimes (Y + I)\xi_n).$$

Our goal is to show that  $L_1$  is a Laplacian in a sense. More precisely,  $L_1$  is a half of the sum of second derivatives in the directions of an orthonormal basis of  $\mathfrak{g}_\infty$ . Define the Laplacian by

$$(5.6) \quad (\Delta v)(X) = \frac{1}{2} \sum_{n=1}^{\infty} (\tilde{\xi}_n \tilde{\xi}_n v)(X),$$

where  $(\tilde{\xi}_n v)(X) = \frac{d}{dt} \Big|_{t=0} v(Xe^{t\xi_n})$  for a function  $v : I + HS \rightarrow \mathbb{R}$  and so  $\tilde{\xi}_n$  is the left-invariant vector field on  $GL(H)$  corresponding to  $\xi_n$ .

Let us calculate derivatives of  $v : I + HS \rightarrow \mathbb{R}$  in the direction of  $\xi_n$

$$(\tilde{\xi}_n v)(X) = v_X(X) \frac{d}{dt} \Big|_{t=0} (Xe^{t\xi_n}) = v_X(X)(X\xi_n)$$

and therefore

$$(\tilde{\xi}_n \tilde{\xi}_n v)(X) = v_{XX}(X)(X\xi_n \otimes X\xi_n) + v_X(X\xi_n^2).$$

Thus the Laplacian is

$$(\Delta v)(X) = \frac{1}{2} \sum_{n=1}^{\infty} [v_{XX}(X)(X\xi_n \otimes X\xi_n) + v_X(X\xi_n^2)] = \frac{1}{2} \sum_{n=1}^{\infty} v_{XX}(X)(X\xi_n \otimes X\xi_n)$$

by Lemma 5.2.

Since  $\tilde{\xi}_n$  is a left-invariant vector field, the Laplacian  $\Delta$  is a left-invariant differential operator such that  $L_1 v = \Delta v$  for any  $v \in C_b^2(I + HS)$ .

**Proposition 5.6.** *For any  $p \geq 2, t > 0$*

$$E \|Y_t\|_{HS}^p < \frac{1}{C_{p,t}} (e^t C_{p,t} - 1),$$

where  $C_{p,t} = C_{\frac{p}{2}} 2^{p-1} (TrQ)^{\frac{p}{2}} t^{\frac{p}{2}-1}$ ,  $C_p = (p(2p-1))^p (\frac{2p}{2p-1})^{2p^2}$ .

*Proof.* First of all, let us estimate  $E \|\int_0^t B(Y_s) dW_s\|_{HS}^p$ . From part 3 of the proof of Theorem 5.1 we know that  $\|B(Y)\|_{L_2^0}^2 \leq 2TrQ(\|Y\|_{HS}^2 + 1)$ . In addition we will use Lemma 7.2 from the book by DaPrato and Zabczyk [3], p.182: for any  $r \geq 1$  and for an arbitrary  $L_2^0$ -valued predictable process  $\Phi(t)$ ,

$$(5.7) \quad E \left( \sup_{s \in [0,t]} \left\| \int_0^s \Phi(u) dW(u) \right\|_{HS}^{2r} \right) \leq C_r E \left( \int_0^t \|\Phi(s)\|_{L_2^0}^2 ds \right)^r, \quad t \in [0, T],$$

where  $C_r = (r(2r-1))^r (\frac{2r}{2r-1})^{2r^2}$ . Thus

$$(5.8) \quad \begin{aligned} E \left\| \int_0^t B(Y_s) dW_s \right\|_{HS}^p &\leq C_{\frac{p}{2}} E \left( \int_0^t \|B(Y_s)\|_{L_2^0}^2 ds \right)^{\frac{p}{2}} \\ &\leq C_{\frac{p}{2}} (2TrQ)^{\frac{p}{2}} E \left( \int_0^t (\|Y\|_{HS}^2 + 1) ds \right)^{\frac{p}{2}} \leq C_{\frac{p}{2}} 2^{\frac{p}{2}} (TrQ)^{\frac{p}{2}} t^{\frac{p}{2}-1} E \int_0^t (\|Y\|_{HS}^2 + 1)^{\frac{p}{2}} ds \end{aligned}$$

Now we can use inequality  $(x+1)^q \leq 2^{q-1}(x^q+1)$  for any  $x \geq 0$  for the estimate (5.8)

$$\begin{aligned} E \left\| \int_0^t B(Y_s) dW_s \right\|_{HS}^p &\leq C_{\frac{p}{2}} 2^{\frac{p}{2}} (TrQ)^{\frac{p}{2}} t^{\frac{p}{2}-1} 2^{\frac{p}{2}-1} E \int_0^t (1 + \|Y_s\|_{HS}^p) ds \\ &= C_{\frac{p}{2}} (TrQ)^{\frac{p}{2}} 2^{p-1} t^{\frac{p}{2}-1} (t + E \int_0^t \|Y\|_{HS}^p ds). \end{aligned}$$

Finally,

$$E \|Y_t\|_{HS}^p \leq E \left\| \int_0^t B(Y_s) dW_s \right\|_{HS}^p \leq C_{p,t} (t + E \int_0^t \|Y_s\|_{HS}^p ds),$$

where  $C_{p,t} = C_{\frac{p}{2}} 2^{p-1} (TrQ)^{\frac{p}{2}} t^{\frac{p}{2}-1}$ .

Thus,  $E \|Y_t\|_{HS}^p < \frac{1}{C_{p,t}} (e^t C_{p,t} - 1)$  by Gronwall's lemma.  $\square$

**Lemma 5.7.** *Let  $f : I + HS \rightarrow [0, \infty]$  be a continuous function in  $L^2(I + HS, \mu_t)$ . If  $\sup_n \|f\|_{t,n} < \infty$  for all  $n$ , then  $Ef(X_n) \xrightarrow[n \rightarrow \infty]{} Ef(X)$ .*

*Proof.* Note that there exists a subsequence  $\{X_{n_k}\}$  such that  $X_{n_k} \xrightarrow[k \rightarrow \infty]{} X$  a.s.

We will prove first that if  $\sup_k \|f\|_{t,n_k} < \infty$ , then  $Ef(X_{n_k}) \xrightarrow[k \rightarrow \infty]{} Ef(X)$ . Denote

$$g_k(\omega) = f(X_{n_k}(\omega)), g(\omega) = f(X(\omega)), \omega \in \Omega.$$

Our goal is to prove that  $\int_{\Omega} g_k(\omega) dP \rightarrow \int_{\Omega} g(\omega) dP$  as  $k \rightarrow \infty$ . Define  $f_l(X) = \min\{f(X), l\}$  for  $l > 0$  and  $g_{k,l}(\omega) = f_l(X_{n_k}(\omega)), g_l(\omega) = f_l(X(\omega))$ . Then  $g_{k,l} \leq l, g_l \leq l$  for any  $\omega \in \Omega$ , so  $\int_{\Omega} g_{k,l}(\omega) dP \xrightarrow[k \rightarrow \infty]{} \int_{\Omega} g_l(\omega) dP$  by the Dominated Convergence Theorem since  $f$  is continuous. By Chebyshev's inequality

$$\begin{aligned} \int_{\Omega} (g_k(\omega) - g_{k,l}(\omega)) dP &= \int_{\{\omega: f(X_{n_k}) \geq l\}} (f(X_{n_k}) - l) dP \\ &\leq \int_{\Omega} f(X_{n_k}) \mathbb{1}_{\{\omega: f(X_{n_k}) \geq l\}} dP \leq \|f\|_{t,n_k} (P\{\omega : f(X_{n_k}) \geq l\})^{\frac{1}{2}} \\ &\leq \|f\|_{t,n_k} \frac{E|f(X_{n_k})|}{l}. \end{aligned}$$

Thus

$$0 \leq \int_{\Omega} (g_k(\omega) - g_{k,l}(\omega)) dP \leq \frac{\sup_n \|f\|_{t,n}^2}{l}.$$

Similarly

$$0 \leq \int_{\Omega} (g(\omega) - g_l(\omega)) dP \leq \frac{1}{l} \|f\|_t E|f(X)|.$$

Therefore  $Ef(X_{n_k}) \rightarrow Ef(X)$  as  $k \rightarrow \infty$ .

To complete the proof suppose that the conclusion is not true. Then there is a subsequence  $X_{n_k}$  such that  $|Ef(X_{n_k}) - Ef(X)| > \varepsilon$  for any  $k$ . However, we always can choose a subsequence  $X_{n_{k_m}}$  such that  $X_{n_{k_m}} \xrightarrow[m \rightarrow \infty]{} X$  a.s. and therefore  $Ef(X_{n_{k_m}}) \xrightarrow[m \rightarrow \infty]{} Ef(X)$ . This is a contradiction.  $\square$

## 6. APPROXIMATION OF THE PROCESS

Let  $P_n$  be the projection onto  $\mathfrak{g}_n$ . Note that since we assume that  $\mathfrak{g}_n$  are invariant subspaces of  $Q$ , the projection from  $\mathfrak{g}_{\infty}$  onto  $\mathfrak{g}_n$  (defined in terms of the norm  $|\cdot|$ ) is the restriction of the projection from  $HS$  onto  $\mathfrak{g}_n$  (defined in terms of the norm  $|\cdot|_{HS}$ ). Note also that since  $\mathfrak{g}_n \cap \text{Ker} Q = \{0\}$ ,  $P_n Q P_n$  is positive and invertible on  $\mathfrak{g}_n$  and  $(P_n Q P_n)^{-1} = P_n Q^{-1} P_n$ .

Choose a real orthonormal basis  $\{\xi_m\}_{m=1}^{\infty}$  of  $\mathfrak{g}_{\infty}$  in the same way as it was done in Section 5. Consider an equation

$$dY_{n,t} = B_n(Y_{n,t}) dW_t, \quad Y_{n,t}(0) = 0,$$

where

$$B_n(Y)U = (Y + I)(P_n U).$$



This equation has a unique solution by the same arguments as in Section 5. Denote  $Q_n = P_n Q P_n$ .

**Lemma 6.1.**  $Y_{n,t}$  is a solution of the equation

$$(6.1) \quad \begin{aligned} dY_{n,t} &= B_n(Y_{n,t})dW_{n,t}, \\ Y_{n,t}(0) &= 0, \end{aligned}$$

where  $W_{n,t} = P_n W_t$ .

*Proof.* Check that  $P_n Q P_n$  is the covariance operator of  $W_{n,t}$ . By the definition of a covariance operator we know that  $\langle f, Qg \rangle_{HS} = E \langle f, W_1 \rangle_{HS} \langle g, W_1 \rangle_{HS}$ , therefore

$$\begin{aligned} \langle f, Q_n g \rangle_{HS} &= \langle P_n f, Q P_n g \rangle_{HS} = E \langle P_n f, W_1 \rangle_{HS} \langle P_n g, W_1 \rangle_{HS} = \\ &= E \langle f, P_n W_1 \rangle_{HS} \langle g, P_n W_1 \rangle_{HS}. \end{aligned}$$

Thus Equation 6.1 is actually the same as Equation 5.1 with the Wiener process that has the covariance  $Q_n$  instead of  $Q$ .  $\square$

We will use the following lemma from [7].

**Lemma 6.2.**

1.  $Tr Q_n \leq Tr Q$ .
2.  $P_n Q \xrightarrow[n \rightarrow \infty]{} Q$ ,
3.  $Q P_n \xrightarrow[n \rightarrow \infty]{} Q$ ,
4.  $P_n Q P_n \xrightarrow[n \rightarrow \infty]{} Q$ ,

where convergence is in the trace class norm.

**Theorem 6.3.** Denote by  $\mathcal{H}_2$  the space of equivalence classes of  $HS$ -valued predictable processes with the norm:

$$\|Y\|_2 = \left( \sup_{t \in [0, T]} E \|Y(t)\|_{HS}^2 \right)^{1/2}.$$

Then

$$\|Y_n - Y\|_2 \xrightarrow[n \rightarrow \infty]{} 0.$$

*Proof of Theorem 6.3.* We want to apply the local inversion theorem (see, for example, Lemma 9.2, from the book by DaPrato and Zabczyk [3], p. 244) to  $K(y, Y) = y + \int_0^t B(Y)dW_t$ , where  $y$  is the initial value of  $Y$  and  $Y = Y(y, t)$  is an  $HS$ -valued predictable process. Analogously we define  $K_n(y, Y) = y + \int_0^t B_n(Y)dW_t$ . To apply this lemma we need to check that  $K$  and  $K_n$  satisfy the following conditions

1. For any  $Y_1(t)$  and  $Y_2(t)$  from  $\mathcal{H}_2$

$$\sup_{t \in [0, T]} E \|K(y, Y_1) - K(y, Y_2)\|_{HS}^2 \leq \alpha \sup_{t \in [0, T]} E \|Y_1(t) - Y_2(t)\|_{HS}^2,$$

where  $0 \leq \alpha < 1$

2. For any  $Y_1(t)$  and  $Y_2(t)$  from  $\mathcal{H}_2$

$$\sup_{t \in [0, T]} E \|K_n(y, Y_1) - K_n(y, Y_2)\|_{HS}^2 \leq \alpha \sup_{t \in [0, T]} E \|Y_1(t) - Y_2(t)\|_{HS}^2,$$

where  $0 \leq \alpha < 1$

3.  $\lim_{n \rightarrow \infty} K_n(y, Y) = K(y, Y)$  in  $\mathcal{H}_2$ .

*Proof of 1.*

To estimate the part with the stochastic differential we will use Lemma 7.2 from [3], p.182: for any  $r \geq 1$  and for arbitrary  $L_2^0$ -valued predictable process  $\Phi(t)$ ,

$$(6.2) \quad E\left(\sup_{s \in [0, t]} \left\| \int_0^s \Phi(u) dW(u) \right\|_{HS}^{2r}\right) \leq C_r E\left(\int_0^t \|\Phi(s)\|_{L_2^0}^2 ds\right)^r, \quad t \in [0, T],$$

where  $C_r = (r(2r-1))^r \left(\frac{2r}{2r-1}\right)^{2r^2}$ . Thus

$$\begin{aligned} E\|K(y, Y_1) - K(y, Y_2)\|_{HS}^2 &= E\left\| \int_0^t B(Y_1) - B(Y_2) dW_s \right\|_{HS}^2 \\ &\leq 4E \int_0^t \|B(Y_1) - B(Y_2)\|_{L_2^0}^2 ds \leq 4TrQE \int_0^t \|Y_1 - Y_2\|_{HS}^2 ds \\ &\leq 4tTrQ \sup_{t \in [0, T]} E\|Y_1 - Y_2\|_{HS}^2. \end{aligned}$$

Note that for small  $t$  we can make  $(2(TrQ)^2t + 8TrQ)t$  as small as we wish, therefore 1 holds.

*Proof of 2.*

Similarly to the proof of condition 2 in the proof of Theorem 5.1 and by Lemma 6.2 we have that

$$\begin{aligned} \|B_n(Y_1) - B_n(Y_2)\|_{L_2^0}^2 &= \|P_n(\cdot)(Y_1 - Y_2)\|_{\mathfrak{g}_\infty \rightarrow HS}^2 \\ &\leq Tr(Q^{1/2}P_nQ^{1/2})\|Y_1 - Y_2\|_{HS}^2 \\ &= Tr(QP_n)\|Y_1 - Y_2\|_{HS}^2 \leq TrQ\|Y_1 - Y_2\|_{HS}^2. \end{aligned}$$

Now use the same estimates as in 1 to see that 2 holds.

*Proof of 3.* Here again we will use (6.2) to estimate the part with the stochastic differential

$$\begin{aligned} &\|K_n(y, Y) - K(y, Y)\|_2^2 = \left\| \int_0^t (B_n(Y) - B(Y)) dW_t \right\|_2^2 \\ &= \\ &\sup_{t \in [0, T]} E\left\| \int_0^t (B_n(Y) - B(Y)) dW_s \right\|_{HS}^2 \leq 4E \int_0^t \|B_n(Y) - B(Y)\|_{L_2^0}^2 ds \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Indeed, estimate  $\|B(Y) - B_n(Y)\|_{L_2^0}^2$ , as

$$\begin{aligned} \|B(Y) - B_n(Y)\|_{L_2^0}^2 &= \sum_{m=1}^{\infty} \|(B(Y) - B_n(Y))\xi_m\|_{HS}^2 \\ &= \sum_{m=1}^{\infty} \|(I - P_n)\xi_m(Y + I)\|_{HS}^2 \leq \|Y + I\|^2 \sum_{m=1}^{\infty} \|(I - P_n)\xi_m\|_{HS}^2 \\ &\leq \|Y + I\|^2 Tr[Q^{1/2}(I - P_n)Q^{1/2}] = \|Y + I\|^2 Tr[(I - P_n)Q^{1/2}Q^{1/2}] \\ &= \|Y + I\|^2 Tr[(I - P_n)Q] \xrightarrow{n \rightarrow \infty} 0 \end{aligned}$$

by Lemma 6.2.

We know that there are unique elements  $Y, Y_n$  in the space  $\mathcal{H}_2$  such that  $Y = K(y, Y)$ ,  $Y_n = K_n(y, Y_n)$  and therefore  $\lim_{n \rightarrow \infty} Y_n = Y$  for any  $y$  by the local inversion lemma.  $\square$

Denote  $P_t^n f(Y) = Ef(Y_n(t, Y))$ ,  $Y \in \mathfrak{g}_n$  and  $v^n(t, Y) = P_t^n f(Y)$ . Then similarly to 5.4 the following holds for  $v^n(t, Y)$ .

For any  $f \in C_b^2(\mathfrak{g}_n)$  the function  $v^n(t, Y)$  is a unique strict solution from  $C_b^{1,2}(\mathfrak{g}_n)$  for the parabolic type equation

$$\begin{aligned} \frac{\partial}{\partial t} v^n(t, Y) &= \frac{1}{2} Tr[v_{YY}^n(B_n(Y)Q^{1/2})(B_n(Y)Q^{1/2})^*] \\ v^n(0, Y) &= f(Y), t > 0, Y \in \mathfrak{g}_n. \end{aligned}$$

We want to show that  $P_t^n$  corresponds to the heat kernel measure defined on  $G_n$  as on a Lie group (i. e. as in the finite dimensional case).

Note that for any  $Y \in \mathfrak{g}_n$

$$\begin{aligned} Tr[v_{YY}^n(t, Y)(B_n(Y)Q^{1/2})(B_n(Y)Q^{1/2})^*] \\ = \sum_{m=1}^{2d_n} v_{YY}^n(t, Y)((Y + I)\xi_m \otimes (Y + I)\xi_m). \end{aligned}$$

Thus

$$L_1 v = \frac{1}{2} Tr[v_{YY}^n(t, Y)(B_n(Y)Q^{1/2})(B_n(Y)Q^{1/2})^*]$$

is equal to the Laplacian  $\Delta_n$  on  $G_n$  defined similarly to (5.6). Thus, the transition probability  $P_t^n$  is equal to  $\mu_t^n(dX)$ , where the latter is the heat kernel measure on  $G_n$  defined in the usual way.

## 7. CAMERON-MARTIN SUBGROUP

The definition of the Cameron-Martin subgroup  $G_{CM}$  was given in Section 2. Let  $C_{CM}^1$  denote the space of piecewise differentiable paths  $h : [0, 1] \rightarrow GL(H)$  such that  $h' = h^{-1}\dot{h}$  is in  $\mathfrak{g}_\infty$ . The main purpose of this section is to prove that  $G_{CM} = G_\infty$  under a condition on the Lie bracket. In doing so we also show that under this condition the exponential map is a local diffeomorphism. This implies, in particular, that the group  $GL(H)$  is closed in the Riemannian metric induced by the operator norm.

**Proposition 7.1.**  *$G_{CM}$  is a group.*

*Proof.* Let  $x, y \in G_{CM}$ . Take  $f : [0, 1] \rightarrow G_{CM}$ ,  $f(0) = x$ ,  $f(1) = y$ . Define  $h(s) = y^{-1}f(s)$ . Then  $|h^{-1}\dot{h}| = |f^{-1}y y^{-1}\dot{f}| = |f^{-1}\dot{f}|$ . Therefore  $d(y^{-1}x, I) \leq d(x, y) \leq d(x, I) + d(y, I) < \infty$ , thus  $xy^{-1} \in G_{CM}$ .  $\square$

**Theorem 7.2.** *If  $|[x, y]| \leq C|x||y|$  for all  $x, y \in \mathfrak{g}_\infty$ , then*

1.  $G_{CM} = G_\infty$ .
2. *The exponential map is a diffeomorphism from a neighborhood of 0 in  $\mathfrak{g}_\infty$  onto a neighborhood of I in  $G_{CM}$ .*

We will need several lemmas before we can prove Theorem 7.2.

**Lemma 7.3.** *Take  $g(t) : [0, 1] \rightarrow G_{CM}$ ,  $g \in C_{CM}^1$ . Suppose that  $\|g(t) - I\| < 1$  for any  $t \in [0, 1]$ .*

*Define  $h(t) = \log g(t) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} (g(t) - I)^n$ . Denote  $A(t) = g(t)^{-1} \dot{g}(t)$ , then  $h$  is a unique solution of the ordinary differential equation in  $\mathfrak{g}_{\infty}$*

$$(7.1) \quad \dot{h}(t) = F(h, t), \quad h(0) = 0,$$

where

$$(7.2) \quad F(x, t) = A(t) + \frac{1}{2}[x, A(t)] - \sum_{p=1}^{\infty} \frac{1}{2(p+2)p!} [\dots [x, A(t)], \overbrace{x, \dots, x}^p].$$

*Proof.* Indeed,  $h(t+s) - h(t) = \log g(t+s) - h(t) = \log(g(t)g(t)^{-1}g(t+s)) - h(t)$ . Denote  $f(t, s) = \log(g(t)^{-1}g(t+s))$ . Then  $h(t+s) - h(t) = \log(e^{h(t)}e^{f(t,s)}) - h(t) = DCH(h(t), f(t, s)) - h(t)$ , where  $DCH(x, y)$  is given by the Dynkin-Campbell-Hausdorff formula for  $x, y \in \mathfrak{g}_{\infty}$

$$(7.3) \quad DCH(x, y) = \log(\exp x \exp y)$$

$$= \sum_m \sum_{p_i, q_i} \frac{(-1)^{m-1}}{m(\sum_i p_i + q_i)} \frac{[\dots [\overbrace{x, x, \dots, x}^{p_1}, \overbrace{y, \dots, y}^{q_1}], \dots, \overbrace{y, \dots, y}^{q_m}], \dots, y]}{p_1! q_1! \dots p_m! q_m!}.$$

Thus

$$\begin{aligned} DCH(h(t), f(t, s)) &= h(t) + f(t, s) + \frac{1}{2}[h(t), f(t, s)] \\ &+ \frac{1}{12}[[h(t), f(t, s)], f(t, s)] - \frac{1}{12}[[h(t), f(t, s)], h(t)] + \dots \end{aligned}$$

Let us find  $\frac{df(t,s)}{ds}|_{s=0}$ .

$$\begin{aligned} &\frac{f(t, s + \varepsilon) - f(t, s)}{\varepsilon} \\ &= \frac{\log(g(t)^{-1}g(t+s+\varepsilon)) - \log(g(t)^{-1}g(t+s))}{\varepsilon} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \frac{(g(t)^{-1}g(t+s+\varepsilon) - I)^n - (g(t)^{-1}g(t+s) - I)^n}{\varepsilon} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \frac{1}{\varepsilon} \sum_{k=1}^n (g(t)^{-1}g(t+s+\varepsilon) - I)^{k-1} \\ &g(t)^{-1}(g(t+s+\varepsilon) - g(t+s))(g(t)^{-1}g(t+s) - I)^{n-k} \xrightarrow{\varepsilon \rightarrow 0} \\ &\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \sum_{k=1}^n (g(t)^{-1}g(t+s) - I)^{k-1} g(t)^{-1} \dot{g}(t) (g(t)^{-1}g(t+s) - I)^{n-k}. \end{aligned}$$

Thus

$$\begin{aligned} \frac{df(t, s)}{ds}|_{s=0} &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \sum_{k=1}^n (g(t)^{-1}g(t) - I)^{k-1} g(t)^{-1} \dot{g}(t) (g(t)^{-1}g(t) - I)^{n-k} \\ &= g(t)^{-1} \dot{g} = A(t). \end{aligned}$$

$$\begin{aligned}
\frac{h(t+s) - h(t)}{s} &= \frac{DCH(h(t), f(t, s)) - h(t)}{s} \\
&= \frac{f(t, s)}{s} + \frac{1}{2}[h(t), \frac{f(t, s)}{s}] + \frac{1}{12}[[h(t), \frac{f(t, s)}{s}], f(t, s)] \\
&\quad + \frac{1}{12}[[h(t), f(t, s)], \frac{f(t, s)}{s}] - \frac{1}{12}[[h(t), \frac{f(t, s)}{s}], h(t)] + \dots
\end{aligned}$$

Note that  $f(t, 0) = 0$ , therefore

$$\begin{aligned}
\dot{h}(t) &= A(t) + \frac{1}{2}[h(t), A(t)] + \frac{d}{ds} \sum_p \frac{(-1)^1}{2(p+2)} \frac{[\dots[[h(t), f(s, t)], \overbrace{h(t), \dots, h(t)}^p]]_{s=0}}{1!1!p!0!} \\
&= A(t) + \frac{1}{2}[h(t), A(t)] - \sum_{p=1}^{\infty} \frac{1}{2(p+2)p!} [\dots[[h(t), A(t)], \overbrace{h(t), \dots, h(t)}^p]].
\end{aligned}$$

Note that 7.1 has a unique solution. Indeed, we show later that  $F$  satisfies local Lipschitz conditions

$$|F(x, t) - F(y, t)| \leq C|x - y|$$

for  $|x| < R, |y| < R$  and according to Theorem 3.1 of [13] equation 7.1 has a unique solution. Indeed, the local Lipschitz conditions are also satisfied for the operator norm. To prove the existence and uniqueness of the solutions in  $\mathfrak{g}_{\infty}$ , we first apply Theorem 3.1 of [13] in  $B(H)$ , then in  $\mathfrak{g}_{\infty}$ .

Now let us show that  $F$  satisfies the local Lipschitz conditions. Denote

$$G(p, x) = [\dots[[x, A(t)], \overbrace{x, \dots, x}^p]].$$

Then

$$|F(x, t) - F(y, t)| \leq \frac{1}{2}|x - y| |A_t| + \sum_{p=1}^{\infty} \frac{1}{2(p+2)p!} |G(p, x) - G(p, y)|.$$

Let us estimate  $|G(p, x) - G(p, y)|$  using induction on  $p$ .

When  $p = 1$ ,

$$\begin{aligned}
|G(1, x) - G(1, y)| &= |[x, A_t], x] - [y, A_t], y]| = \\
&|[[x, A_t], x - y] + [[x, A_t] - [y, A_t], y]| \leq |x - y| |A_t| C(C|x| + C|y|) = \\
&C^2 |x - y| |A_t| (|x| + |y|)
\end{aligned}$$

Assume that for  $p = k$  we have the estimate

$$|G(k, x) - G(k, y)| \leq C^{k+1} |x - y| |A_t| \sum_{i=0}^k |x|^i |y|^{k-i}$$

Then for  $p = k + 1$  we have

$$\begin{aligned}
|G(k+1, x) - G(k+1, y)| &= |[G(k, x), x] - [G(k, y), y]| \\
&= |[G(k, x), x] - [G(k, x), y] + [G(k, x), y] - [G(k, y), y]| \\
&\leq C |G(k, x)| |x - y| + C |G(k, x) - G(k, y)| |y| \\
&\leq C^{k+2} |x|^{k+1} |A_t| |x - y| + C^{k+2} |x - y| |A_t| \left( \sum_{i=0}^k |x|^i |y|^{k-i} \right) |y| \\
&= C^{k+2} |x - y| |A_t| \sum_{i=0}^{k+1} |x|^i |y|^{k+1-i}.
\end{aligned}$$

Thus if  $|x| \leq R, |y| \leq R$

$$|G(p, x) - G(p, y)| \leq C^{p+1} |x - y| |A_t| R^p (p + 1)$$

and therefore

$$\begin{aligned}
|F(x, t) - F(y, t)| &\leq \frac{1}{2} |x - y| |A_t| + \sum_{p=1}^{\infty} \frac{1}{2(p+2)p!} C^{p+1} |x - y| |A_t| R^p (p + 1) \\
&\leq \frac{1}{2} |x - y| |A_t| + \frac{1}{2} |x - y| |A_t| C \sum_{p=1}^{\infty} \frac{C^p R^p}{p!} = \frac{1}{2} |x - y| |A_t| (1 + C(e^{CR} - 1)).
\end{aligned}$$

□

**Lemma 7.4.** *There are constants  $C_1, C_2$  and  $0 < \varepsilon < \ln 2/2C$  such that*

$$C_1 |x - y| \leq d(\exp x, \exp y) \leq C_2 |x - y|$$

for any  $x, y \in \mathfrak{g}_{\infty}$  provided  $|x| < \varepsilon, |y| < \varepsilon$ .

*Proof.* Take a path  $g : [0, 1] \rightarrow \exp(\mathfrak{g}_{\infty})$  such that  $g(0) = e^x, g(1) = e^y$ . Let  $h = \log g$ . Then

$$g^{-1}\dot{g} = \sum_{k=0}^{\infty} \frac{(-adh)^k (\dot{h})}{(k+1)!} = \dot{h} - [h, \dot{h}] + \frac{[h, [h, \dot{h}]]}{2!} - \frac{[h, [h, [h, \dot{h}]]]}{3!} + \dots$$

1. First prove that  $d(\exp x, \exp y) \leq C_2 |x - y|$ :

$$|g^{-1}\dot{g}| \leq |\dot{h}| \sum_{k=0}^{\infty} \frac{C^k |h|^k}{k!} = |\dot{h}| e^{C|h|}$$

$$\implies d(\exp x, \exp y) = \inf_g \int_0^1 |g^{-1}\dot{g}| ds \leq \inf_h \int_0^1 |\dot{h}| e^{C|h|} ds.$$

Take  $h(s) = xs + (1-s)y$

$$\begin{aligned}
d(\exp x, \exp y) &\leq \int_0^1 |x - y| e^{C(|x|s + (1-s)|y|)} ds = |x - y| \int_0^1 e^{C(|x| - |y|)s} e^{C|y|} ds \\
&= |x - y| e^{C|y|} \frac{e^{C(|x| - |y|)} - 1}{C(|x| - |y|)} = |x - y| \frac{e^{C|x|} - e^{C|y|}}{C(|x| - |y|)} \leq |x - y| e^{C \max\{|x|, |y|\}},
\end{aligned}$$

since  $\frac{e^a - e^b}{a - b} \leq e^{\max\{|a|, |b|\}}$ . Thus

$$d(\exp x, \exp y) \leq |x - y| e^{C \max\{|x|, |y|\}} \leq |x - y| e^{C\varepsilon}.$$

2. Now let us prove that  $C_1 |x - y| \leq d(\exp x, \exp y)$ :

$$\begin{aligned} |g^{-1}\dot{g}| &= \left| \sum_{k=0}^{\infty} \frac{(-adh)^k(\dot{h})}{(k+1)!} \right| \geq |\dot{h}| \left( 1 - \sum_{k=1}^{\infty} \frac{|h|^k C^k}{k!} \right) \\ &= |\dot{h}| \left( 1 - \sum_{k=0}^{\infty} \frac{|h|^k C^k}{k!} + 1 \right) = |\dot{h}| (2 - e^{C|h|}). \end{aligned}$$

Consider two cases.

2a. Suppose  $\max_s |h(s)| < 2\varepsilon$ . Then

$$\begin{aligned} \int_0^1 |g^{-1}\dot{g}| ds &\geq \int_0^1 |\dot{h}| (2 - e^{C|h|}) ds \\ &\geq (2 - e^{2\varepsilon C}) \int_0^1 |\dot{h}| ds \geq (2 - e^{2\varepsilon C}) \left| \int_0^1 \dot{h} ds \right| = (2 - e^{2\varepsilon C}) |x - y|. \end{aligned}$$

2b. Take a path  $h$  such that  $\max_s |h(s)| \geq 2\varepsilon$ . Denote  $t_1 = \min\{t : |h(t)| = \varepsilon\}$ ,  $t_2 = \min\{t : |h(t)| = 2\varepsilon\}$ . Note that  $t_1 < t_2$  since  $h(0) = x$  and  $|x| < \varepsilon$ . Then

$$\begin{aligned} \int_0^1 |g^{-1}\dot{g}| ds &\geq \int_{t_1}^{t_2} |g^{-1}\dot{g}| ds \\ &\geq \int_{t_1}^{t_2} |\dot{h}| (2 - e^{C|h|}) ds \geq (2 - e^{2\varepsilon C}) \int_{t_1}^{t_2} |\dot{h}| ds \geq (2 - e^{2\varepsilon C}) |h(t_2) - h(t_1)| \\ &\geq \varepsilon (2 - e^{2\varepsilon C}) \geq \frac{|x - y|}{2} (2 - e^{2\varepsilon C}). \end{aligned}$$

2a and 2b imply that

$$d(\exp x, \exp y) = \inf_g \int_0^1 |g^{-1}\dot{g}| ds \geq \frac{|x - y|}{2} (2 - e^{2\varepsilon C}).$$

□

**Lemma 7.5.** *Take  $g(t) : [0, 1] \rightarrow G_{CM}$ ,  $g \in C_{CM}^1$ . Suppose that  $g(0) = e$ ,  $\|g(t) - e\| < 1$  for any  $t \in [0, 1]$  and  $|\log g(1)| < \varepsilon$ , where  $\varepsilon$  is the same as in Lemma 7.4. Then  $g(1) \in G_\infty$ .*

*Proof.* Let  $h = \log g = \sum_{n=1}^{\infty} ((-1)^{n-1}/n)(g(t) - I)^n$ . Then by Lemma 7.3  $h$  is a unique solution in  $\mathfrak{g}_\infty$  of the ordinary differential equation 7.1. In particular, it implies that  $g(t) \in \mathfrak{g}_\infty$ . Therefore, there are  $H_n \in \mathfrak{g}_n$  (projections of  $h(1)$  onto  $\mathfrak{g}_n$ ) such that  $|h(1) - H_n| \xrightarrow{n \rightarrow \infty} 0$ .

By Lemma 7.4

$$d(e^{h(1)}, e^{H_n}) \leq C_2 |h(1) - H_n|.$$

A similar estimate holds for  $d_n$  instead of  $d$ . By a direct calculation  $e^{h(t)} = g(t)$ . Thus  $g(1) \in G_\infty$ , since  $e^{H_n} \in G_n$ . □

**Proof of Theorem 7.2.** 1.  $G_\infty \subseteq G_{CM}$  since  $G_n \subseteq G_{CM}$  for all  $n$  and  $G_\infty$  is closed in the metric  $d_\infty$ .

Now we will show that  $G_{CM} \subseteq G_\infty$ .

Take  $A \in G_\infty$ ,  $B \in G_{CM}$ , a path  $g : [0, 1] \rightarrow G_{CM}$ ,  $g \in C_{CM}^1$ ,  $g(0) = A$ ,  $g(1) = B$ . We want to show that  $B \in G_\infty$ .

Lemma 7.5 proves this in a neighborhood of  $I$ . Prove it for a neighborhood of any element of  $G_\infty$ . Suppose  $\|g(t) - A\| < 1$ ,  $d(\log B, \log A) < \varepsilon$ .

For any  $\delta > 0$  there is  $C_\delta \in \bigcup_n G_n$  such that  $d(A, C_\delta) < \delta$ . Thus

$$d(B, C_\delta) \leq d(B, A) + d(A, C_\delta) < \varepsilon + \delta.$$

Therefore  $d(C_\delta^{-1}B, e) < \varepsilon + \delta$ , so by part 1  $C_\delta^{-1}B \in G_\infty$ . This means that there is  $D_\delta \in \bigcup_n G_n$  such that

$$d(C_\delta^{-1}B, D_\delta) < \delta.$$

Now  $d(C_\delta^{-1}B, D_\delta) = d(B, C_\delta D_\delta) < \delta$ ,  $C_\delta D_\delta \in \bigcup_n G_n$ . This proves that  $B \in G_\infty$ .

Now take any  $A, B$  and a path  $g$  joining them. Divide  $\log g$  into subpaths satisfying the conditions in the first part of the proof.

2. By Lemma 7.3 and Lemma 7.4  $\exp$  and  $\log$  are well defined and differentiable in neighborhoods of the identity and zero respectively.  $\square$

## 8. HOLOMORPHIC POLYNOMIALS AND SKELETONS

**Definition 8.1.** A function  $p : I + HS \rightarrow \mathbb{C}$  is called a *holomorphic polynomial*, if  $p$  is a complex linear combination of finite products of monomials  $p_k^{m,l}(X) = (\langle X f_m, f_l \rangle)^k$ , where  $\{f_m\}_{m=1}^\infty$  is an orthonormal basis of  $H$ . We will denote the space of all such polynomials by  $\mathcal{HP}$ .

These polynomials are holomorphic because the  $k$ th derivative  $(D_X^k p)(\beta)$  exists and is complex linear for any  $p \in \mathcal{P}$ ,  $X \in I + HS$ ,  $\beta \in \mathfrak{g}^{\otimes k}$ . Indeed, let  $\xi$  be any element of  $\mathfrak{g}_\infty$ , then the derivative of  $p_k^{m,l}$  in the direction of  $\xi$  is  $(D p_k^{m,l})(X)(\xi) = k p_1^{m,l}(X \xi) p_{k-1}^{m,l}(X)$ . From this formula we can see that  $D p_k^{m,l}(X)(\xi)$  is complex linear in  $\xi$ .

**Remark 8.2.** Any polynomial  $p \in \mathcal{HP}$  can be written in the form  $p(X) = \sum_{k=1}^m \prod_{l=1}^{n_m} Tr(A_{kl} X)$  for some  $A_{kl} \in HS$ . The converse is not true in general, but the closure in  $L^2(I + HS, \mu_t)$  of all functions of the form  $\sum_{k=1}^m \prod_{l=1}^{n_m} Tr(A_{kl} X)$  coincides with the closure of holomorphic polynomials. Therefore the next definition is basis-independent, though the definition of  $\mathcal{HP}$  depends on the choice of  $\{f_k\}_{k=1}^\infty$ .

**Definition 8.3.** The closure of all holomorphic polynomials in  $L^2(I + HS, \mu_t)$  is called  $\mathcal{HL}^2(I + HS, \mu_t)$ .

**Lemma 8.4.** (*B. Driver's formula*).

Let  $g$  be a smooth path in  $G_{CM}$  such that  $g(0) = I$  and  $g \in C_{CM}^1$ . Then for any  $p \in \mathcal{HP}$

$$p(g(s)) = \sum_{n=0}^{\infty} \int_{\Delta_n(s)} (D^n p)(I)(c(s_1) \otimes \dots \otimes c(s_n)) d\vec{s},$$

where

$$\Delta_n(s) = \{(s_1, \dots, s_n) \in \mathbb{R}^n : 0 \leq s_1 \leq s_2 \leq \dots \leq s_n \leq s\}, c(s) = g^{-1}(s) \frac{dg}{ds}.$$



*Proof.* Take a monomial  $p(x) = (\langle Xf_m, f_l \rangle)^k$ ,  $\deg p = k$ . Similarly to Lemma 5.2 of [4]

$$p(g(s)) = \sum_{n=0}^{N-1} \int_{\Delta_n(s)} (D^n p)(I)(c(s_1) \otimes \dots \otimes c(s_n)) d\vec{s} \\ + \int_{\Delta_N(s)} (D^N p)(g(s_1))(c(s_1) \otimes \dots \otimes c(s_N)) d\vec{s}.$$

We want to estimate the remainder

$$\left| \int_{\Delta_N(s)} (D^N p)(g(s_1))(c(s_1) \otimes \dots \otimes c(s_N)) d\vec{s} \right| \\ \leq \int_{\Delta_N(s)} |(D^N p)(g(s_1))(c(s_1) \otimes \dots \otimes c(s_N))| d\vec{s}.$$

Note that if  $k = \deg p \leq N$  then

$$(D^N p)(X)(\xi_1 \otimes \dots \otimes \xi_N) = \sum p_1(X\vec{\xi}^{\alpha_1}) p_1(X\vec{\xi}^{\alpha_2}) \cdot \dots \cdot p_1(X\vec{\xi}^{\alpha_i}) \cdot \dots \cdot p_1(X\vec{\xi}^{\alpha_k}),$$

where  $\alpha_i = \langle a_1^i, \dots, a_N^i \rangle$ ,  $a_j^i = 0$  or  $1$ ,  $\vec{\xi}^{\alpha_i} = \xi_1^{a_1^i} \dots \xi_N^{a_N^i}$ ,  $\sum_{i=1}^k \alpha_i = \langle 1, \dots, 1 \rangle$ ,  $p_1(Y) = \langle Xf_m, f_l \rangle$  and the sum is taken over all possible  $\alpha = (\alpha_1, \dots, \alpha_k)$ .

For  $\xi \in HS$ ,  $X \in I + HS$

$$|p_1(X\xi)| \leq (\|X - I\|_{HS} + \|\xi\|_{HS}) \leq \|\xi\|_{HS} (\|X - I\|_{HS} + 1).$$

The number of all such  $\alpha$  is  $k^N$  therefore

$$|(D^N p)(X)(\xi_1 \otimes \dots \otimes \xi_N)| \leq \|\xi_1\|_{HS} \dots \|\xi_N\|_{HS} (\|X - I\|_{HS} + 1) k^N.$$

Denote

$$\|c\|_1 = \int_0^1 |c(t)| dt.$$

Thus

$$\int_{\Delta_N(s)} |(D^N p)(g(s_1))(c(s_1) \otimes \dots \otimes c(s_N))| d\vec{s} \\ \leq \|c\|_1^N \sup_{0 \leq t \leq s} (\|g(t) - I\|_{HS} + 1)^k k^N \int_{\Delta_N(s)} d\vec{s} \\ = \|c\|_1^N \sup_{0 \leq t \leq s} (\|g(t) - I\|_{HS} + 1)^k k^N \frac{s^N}{N!} \xrightarrow{N \rightarrow \infty} 0.$$

Similarly we can prove this formula for any holomorphic polynomial.  $\square$

**Theorem 8.5.** *Suppose*

$$p_n \xrightarrow[n \rightarrow \infty]{L^2(I+HS, \mu_t)} f, \quad p_n \in \mathcal{HP}.$$

*Then there is a holomorphic function  $\tilde{f}$ , a skeleton of  $f$ , on  $G_{CM}$  such that*

$$p_n(x) \rightarrow \tilde{f}(x) \text{ for any } x \in G_{CM}.$$

*Proof.* As we know from Theorem 4.4 the map  $(1 - D)_I^{-1}$  is an isometry from  $\mathcal{HL}^2(I + HS, \mu_t)$  to  $J_t^0$ . Thus  $(1 - D)_I^{-1} p_n$  converges in  $J_t^0$  to some  $\alpha = \sum \alpha_n$ . Define

$$\tilde{f}(x) = \sum_{n=0}^{\infty} \int_{\Delta_n(1)} \alpha_n (c(s_1) \otimes \dots \otimes c(s_n)) d\vec{s}.$$

Take a path  $g : [0, 1] \rightarrow G_{CM}$ ,  $g(0) = e, g(1) = X$ . By Lemma 8.4

$$p_m(X) = \sum_{n=0}^{\infty} \int_{\Delta_n(1)} (D^n p_m)(I) (c(s_1) \otimes \dots \otimes c(s_n)) d\vec{s},$$

so

$$\begin{aligned} |p_m(X) - \tilde{f}(x)| &\leq \sum_{n=0}^{\infty} \int_{\Delta_n(1)} |((D^n p_m)(I) - \alpha_n)(c(s_1) \otimes \dots \otimes c(s_n))| d\vec{s} \\ &\leq \sum_{n=0}^{\infty} \int_{\Delta_n(1)} |(D^n p_m)(I) - \alpha_n| |c(s_1) \otimes \dots \otimes c(s_n)| d\vec{s} \\ &\leq \sum_{n=0}^{\infty} |(D^n p_m)(I) - \alpha_n| \int_{\Delta_n(1)} |c(s_1) \otimes \dots \otimes c(s_n)| d\vec{s}. \end{aligned}$$

Thus

$$\begin{aligned} |p_m(X) - \tilde{f}(x)| &\leq \sum_{n=0}^{\infty} |(D^n p_m)(I) - \alpha_n| \frac{\|c\|_1^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{|(D^n p_m)(I) - \alpha_n| t^{n/2}}{\sqrt{n!}} \frac{\|c\|_1^n}{\sqrt{n!} t^{n/2}} \leq \left( \sum_{n=0}^{\infty} \frac{|(D^n p_m)(I) - \alpha_n|^2 t^n}{n!} \right) \left( \sum_{n=0}^{\infty} \frac{\|c\|_1^{2n}}{n! t^n} \right) \\ &= \|(Dp_m)(I) - \alpha\|_t^2 e^{\|c\|_1^2/t} \xrightarrow{m \rightarrow \infty} 0. \end{aligned}$$

Now we will show that  $\tilde{f}$  is holomorphic. Take  $\xi \in \mathfrak{g}$ ,  $g \in C_{CM}^1$ ,  $g(0) = e, g(1) = X$ . Define two paths in  $G_{CM}$

$$g_1(s) = \begin{cases} g(s), & 0 \leq s \leq 1 \\ g(1), & 1 \leq s \leq 1+t \end{cases}$$

$$g_2(s) = \begin{cases} g(s), & 0 \leq s \leq 1 \\ g(1)e^{(s-1)\xi}, & 1 \leq s \leq 1+t \end{cases}$$

Let  $c(s) = g^{-1}(s) \frac{dg}{ds}$ ; then

$$c_1(s) = g_1^{-1}(s) \frac{dg_1}{ds} = \begin{cases} c(s), & 0 \leq s \leq 1 \\ 0, & 1 \leq s \leq 1+t \end{cases}$$

$$c_2(s) = g_2^{-1}(s) \frac{dg_2}{ds} = \begin{cases} c(s), & 0 \leq s \leq 1 \\ \xi, & 1 \leq s \leq 1+t \end{cases}$$

Then by the definition of  $\tilde{f}$

$$\begin{aligned} \tilde{f}(Xe^{t\xi}) - \tilde{f}(X) &= \sum_{n=0}^{\infty} \int_{\Delta_n(1+t)} \alpha_n(c_1(s_1) \otimes \dots \otimes c_1(s_n) - c_2(s_1) \otimes \dots \otimes c_2(s_n)) d\vec{s}_n \\ &= \sum_{n=0}^{\infty} \int_1^{1+t} \int_{\Delta_{n-1}(s_n)} \alpha_n(c_1(s_1) \otimes \dots \otimes c_1(s_n) - c_2(s_1) \otimes \dots \otimes c_2(s_n)) d\vec{s}_{n-1} ds_n \\ &= \sum_{n=0}^{\infty} \int_{\Delta_{n-1}(1)} \alpha_n(c(s_1) \otimes \dots \otimes c(s_{n-1}) \otimes t\xi) d\vec{s}_{n-1} \\ &+ \sum_{n=0}^{\infty} \int_1^{1+t} \int_1^{s_n} \int_{\Delta_{n-2}(s_{n-1})} \alpha_n(c_1(s_1) \otimes \dots \otimes c_1(s_n) - c_2(s_1) \otimes \dots \otimes c_2(s_n)) d\vec{s}_{n-2} ds_{n-1} ds_n \\ &= \sum_{n=0}^{\infty} t \int_{\Delta_{n-1}(1)} \alpha_n(c(s_1) \otimes \dots \otimes c(s_{n-1}) \otimes \xi) d\vec{s}_{n-1} \\ &+ \sum_{n=0}^{\infty} \int_1^{1+t} \int_1^{s_n} \int_{\Delta_{n-2}(s_{n-1})} \alpha_n(c(s_1) \otimes \dots \otimes c(s_{n-2}) \otimes \xi \otimes \xi) d\vec{s}_{n-2} ds_{n-1} ds_n. \end{aligned}$$

Thus

$$\left| \frac{\tilde{f}(Xe^{t\xi}) - \tilde{f}(X)}{t} - \int_{\Delta_{n-1}(1)} \alpha_n(c(s_1) \otimes \dots \otimes c(s_{n-1}) \otimes \xi) d\vec{s}_{n-1} \right| \leq$$

$$\frac{1}{t} \int_1^{1+t} (s_n - 1) \int_{\Delta_{n-2}(s_{n-1})} \alpha_n(c(s_1) \otimes \dots \otimes c(s_{n-2}) \otimes \xi \otimes \xi) d\vec{s}_{n-2} ds_{n-1} ds_n \xrightarrow{n \rightarrow 0} 0$$

since  $0 \leq s_n - 1 \leq t$ .

This means that the derivative of  $\tilde{f}$  at  $X$  in the direction  $\xi$  is

$$(\xi \tilde{f})(X) = \int_{\Delta_{n-1}(1)} \alpha_n(c(s_1) \otimes \dots \otimes c(s_{n-1}) \otimes \xi) d\vec{s}_{n-1}$$

and it is complex linear.  $\square$

**Remark 8.6.** The convergence is uniform on bounded sets. In fact, one can show that the derivatives also converge uniformly on bounded sets.

**Theorem 8.7.** 1.  $\mathcal{HP} \subset L^p(I + HS, \mu_t)$ , for any  $p > 1$ .

2.  $\|f\|_{t,n} \xrightarrow{n \rightarrow \infty} \|f\|_t$  for any  $f \in \mathcal{HP}$ .
3. The embedding of  $\mathcal{HP}$  into  $\mathcal{H}^t(G_\infty)$  can be extended to an isometry from  $\mathcal{HL}^2(I + HS, \mu_t)$  into  $\mathcal{H}^t(G_\infty)$ .

*Proof.* 1.

$$\begin{aligned} \int_{I+HS} |p_k^{m,l}(X)|^p \mu_t(dX) &= E|p_k^{m,l}(Y_t + I)|^p = E|(Y_t)_{ml} + \delta_{ml}|^{pk} \\ &\leq E2^{pk-1}(|(Y_t)_{ml}|^{pk} + 1) \leq 2^{pk-1}E(\|Y_t\|_{HS}^{pk} + 1) < \infty \end{aligned}$$

by Lemma 5.7. Thus  $\mathcal{HP} \subset L^p(I + HS, \mu_t)$  for any  $p > 1$ .

2. From the estimate on  $E\|Y_t\|_{HS}^p$  in Proposition 5.6 we can find  $C(p, t)$  such that  $\| |p|^2 \|_{t,n(k)} \leq C(p, t)$  for any  $k$ . Then apply Lemma 5.7 to  $f = |p|^2$ .

3. By part 1 of Theorem 8.7  $\mathcal{HP} \subset L^2(I + HS, \mu_t)$ . In addition, by part 2 of Theorem 8.7

$$\|p\|_{t,n} \xrightarrow{n \rightarrow \infty} \|p\|_t, \quad p \in \mathcal{HP}.$$

Therefore  $\|p\|_{t,\infty} = \|p\|_t$  and so the embedding is an isometry.

The space  $\mathcal{HL}^2(I + HS, \mu_t)$  is the closure of  $\mathcal{HP}$ , therefore the isometry extends to it from  $\mathcal{HP}$ .  $\square$

**Remark 8.8.** If  $f$  is an element of  $\mathcal{HL}^2(I + HS, \mu_t)$ ,  $g$  is its image under the isometry in this theorem and  $\tilde{f}$  is the skeleton of  $f$ , then  $\tilde{f}|_{G_\infty} = g$ .

**Corollary 8.9.** *The space  $\mathcal{H}^t(G_\infty)$  is an infinite dimensional Hilbert space.*

## 9. EXAMPLES

Denote  $HS_{n \times n} = \{A : \langle Af_m, f_k \rangle = 0 \text{ if } \max(m, k) > n\}$ . Take a basis  $\{e_k\}_{k=1}^\infty$  of  $HS$  such that  $\{e_k\}_{k=1}^{2n^2}$  is a basis of  $HS_{n \times n}$ . By  $B^T$  we will denote the transpose of the operator  $B$ , i. e.  $B^T = (ReB)^* + i(ImB)^*$ .

**Example 9.1.** We begin with the definition of the Hilbert-Schmidt complex orthogonal group.

**Definition 9.1.** *The Hilbert-Schmidt complex orthogonal group  $SO_{HS}$  is the connected component containing the identity  $I$  of the group  $O_{HS} = \{B : B - I \in HS, B^T B = BB^T = I\}$ . The Lie algebra of skew-symmetric Hilbert-Schmidt operators will be denoted by  $\mathfrak{so}_{HS} = \{A : A \in HS, A^T = -A\}$ .*

Let  $Q$  be a symmetric positive trace class operator on  $\mathfrak{so}_{HS}$  and let the inner product be defined by  $\langle A, B \rangle = \langle Q^{-1/2}A, Q^{-1/2}B \rangle_{HS}$ . In [7] we showed that if  $Q$  is the identity operator, then all Hilbert spaces we consider are isomorphic to  $\mathbb{C}$ ; that is, there are no nonconstant holomorphic functions. As in Section 5 we identify  $Q$  with its extension by 0 to the orthogonal complement of  $\mathfrak{so}_{HS}$ .

Define groups  $G_n = SO(n, \mathbb{C}) = \{B \in SO_{HS}, B - I \in HS_{n \times n}\}$ . These groups are isomorphic to the special complex orthogonal group of  $\mathbb{C}^n$ . Their Lie algebras are  $Lie(SO(n, \mathbb{C})) = \mathfrak{so}(n, \mathbb{C}) = \{A \in HS_{n \times n}, A^T = -A\}$  with an inner product  $\langle A, B \rangle_n = \langle (P_n Q P_n)^{-1/2} A, (P_n Q P_n)^{-1/2} B \rangle_{HS}$ . Here  $P_n = P_{\mathfrak{so}(n, \mathbb{C})}$ . We assume that all  $\mathfrak{so}(n, \mathbb{C})$  are invariant subspaces of  $Q$ . The groups  $SO(n, \mathbb{C})$  are not simply connected, therefore we have isometries from  $\mathcal{H}^t(SO_\infty)$  and  $\mathcal{HL}^2(I + HS, \mu_t)$  to  $J_t^0$ ,

but not an isomorphism between  $\mathcal{H}^t(SO_\infty)$  and  $J_t^0$ . In addition to the properties of the heat kernel measure described in this paper, we showed in [7] that the process  $Y_t + I$  actually lives in the group  $SO_{HS}$ .

**Example 9.2.** The Hilbert-Schmidt complex symplectic group is defined similarly to the Hilbert-Schmidt complex orthogonal group.

**Definition 9.2.** The Hilbert-Schmidt complex symplectic group  $Sp_{HS}$  is the group of operators  $X = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  such that  $A - I, D - I, B, C \in HS$  and  $X^T J X = J$ , where  $J = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ . The Lie algebra is  $\mathfrak{sp}_{HS} = \{X = \begin{pmatrix} A & B \\ C & D \end{pmatrix} : A, B, C, D \in HS, X^T J + J X = 0\}$ .

The corresponding finite dimensional groups are isomorphic to the classical symplectic complex groups  $Sp(n, \mathbb{C}) = \{X = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp_{HS}, A - I, D - I, B, C \in HS_{n \times n}\}$  with Lie algebras  $\mathfrak{sp}(n, \mathbb{C}) = \{X = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathfrak{sp}_{HS} : A, B, C, D \in HS_{n \times n}\}$ . An inner product on  $\mathfrak{sp}_{HS}$  and  $\mathfrak{sp}(n, \mathbb{C})$  is defined in the same way as in Example 9.1. Similarly to  $\mathfrak{so}_{HS}$  if  $Q$  is the identity operator, the corresponding  $J_0^t$  is trivial by Theorem 4.6. The groups  $Sp(n, \mathbb{C})$  are simply connected; therefore the isometry from  $\mathcal{H}^t(Sp_\infty)$  to  $J_t^0$  is surjective. Similar to  $SO_{HS}$  the process  $Y_t + I$  lives in  $Sp_{HS}$ .

**Statement 9.3.**  $Y_t + I$  lies in  $Sp_{HS}$  for any  $t > 0$  with probability 1.

*Proof.* We need to check that  $(Y_t + I)J(Y_t + I)^T = J$  with probability 1 for any  $t > 0$ . To do this we will apply Itô's formula to  $G(Y_t)$ , where  $G$  is defined as follows:  $G(Y) = \Lambda(YJY^T + YJ + JY^T)$ ,  $\Lambda$  is a linear real bounded functional from  $HS$  to  $\mathbb{R}$ .

In order to use Itô's formula we must verify several properties of the process  $Y_t$  and the mapping  $G$ :

1.  $B(Y_s)$  is an  $L_2^0$ -valued process stochastically integrable on  $[0, T]$ .
2.  $G$  and the derivatives  $G_t, G_Y, G_{YY}$  are uniformly continuous on bounded subsets of  $[0, T] \times HS$ .

*Proof of 1.* See 1 in the proof of Theorem 5.1.

*Proof of 2.* Let us calculate  $G_t, G_Y, G_{YY}$ . First of all,  $G_t = 0$ . For any  $S \in HS$

$$G_Y(Y)(S) = \Lambda(SJY^T + YJS^T + SJ + JS^T).$$

For any  $S, T \in HS$

$$G_{YY}(Y)(S \otimes T) = \Lambda(SJT^T + TJS^T).$$

Thus condition 2 is satisfied.

We will use the notation:

$$\begin{aligned} G_Y(Y)(S) &= \langle \bar{G}_Y(Y), S \rangle_{HS}, \\ G_{YY}(Y)(S \otimes T) &= \langle \bar{G}_{YY}(Y)S, T \rangle_{HS}, \end{aligned}$$

where  $\bar{G}_Y$  is an element of  $HS$  and  $\bar{G}_{YY}$  is an operator on  $HS$  corresponding to the functionals  $G_Y \in HS^*$  and  $G_{YY} \in (HS \otimes HS)^*$ .

Now we can apply Itô's formula to  $G(Y_t)$ :

$$(9.1) \quad G(Y_t) = \int_0^t \langle \bar{G}_Y(Y_s), B(Y_s) dW_s \rangle_{HS} + \int_0^t \frac{1}{2} \text{Tr}[G_{YY}(Y_s)(B(Y_s)Q^{1/2})(B(Y_s)Q^{1/2})^*] ds.$$

Let us calculate the two integrands in (9.1) separately.

The first integrand is

$$\begin{aligned} \langle \bar{G}_Y(Y_s), B(Y_s) dW_s \rangle_{HS} &= \langle \bar{G}_Y(Y_s), (Y_s + I) dW_s \rangle_{HS} \\ &= \Lambda(((Y_s + I) dW_s) J Y_s^T + Y_s J ((Y_s + I) dW_s)^T + ((Y_s + I) dW_s) J + J ((Y_s + I) dW_s)^T) \\ &= \Lambda((Y_s + I) dW_s J Y_s^T + Y_s J (dW_s^T (Y_s^T + I)) + (Y_s + I) dW_s J + J dW_s^T (Y_s^T + I)) \\ &= \Lambda((Y_s + I) dW_s J Y_s^T - Y_s dW_s J (Y_s^T + I) + (Y_s + I) dW_s J - dW_s J (Y_s^T + I)) = 0, \end{aligned}$$

since  $W_t$  is a  $\mathfrak{sp}_{HS}$ -valued process.

The second integrand is

$$\begin{aligned} &\frac{1}{2} \text{Tr}[G_{YY}(Y_s)(B(Y_s)Q^{1/2})(B(Y_s)Q^{1/2})^*] \\ &= \frac{1}{2} \sum_{n=1}^{\infty} \langle \bar{G}_{YY}(Y_s) B(Y_s) Q^{1/2} e_n, B(Y_s) Q^{1/2} e_n \rangle_{HS} \\ &= \frac{1}{2} \sum_{n=1}^{\infty} \Lambda(2((Y_s + I) \xi_n) J ((Y_s + I) \xi_n)^T) = \sum_{n=1}^{\infty} \Lambda((Y_s + I) \xi_n J \xi_n^T (Y_s^T + I)) \\ &= - \sum_{n=1}^{\infty} \Lambda((Y_s + I) \xi_n^2 J (Y_s^T + I)) = 0 \end{aligned}$$

by Lemma 5.2. This shows that the stochastic differential of  $G$  is zero, so  $G(Y_t) = 0$  for any  $t > 0$ .  $\square$

**Example 9.3.** Let  $G$  be a group of diagonal (infinite) complex matrices  $\text{diag}(1 + a_1, \dots, 1 + a_i, \dots)$ , where  $\sum_{i=1}^{\infty} |a_i|^2 < \infty, a_i \neq -1$ . Then  $G \cong \prod_{i=1}^{\infty} \mathbb{C} \setminus \{0\}$ . This group is abelian and is not simply connected. The Lie algebra  $\mathfrak{g}$  is an algebra of diagonal matrices  $\text{diag}(a_1, \dots, a_i, \dots)$ , where  $\sum_{i=1}^{\infty} |a_i|^2 < \infty$ . Note that if we take operator  $Q$  as before to define a new inner product on  $\mathfrak{g}$ , this inner product is  $Ad_{G_{\mathbb{R}}}$ -invariant. The process is a sum of processes in  $\mathbb{C}$

$$dY_t^i = B^i(Y_t^i) dW_t^i.$$

We can also write this equation in terms of the real and imaginary parts of  $Y_t^i$

$$\begin{aligned} dY_t^{i,1} &= (Y_t^{i,1} + 1) dW_t^{i,1} - Y_t^{i,2} dW_t^{i,2} \\ dY_t^{i,2} &= Y_t^{i,2} dW_t^{i,1} + (Y_t^{i,2} + 1) dW_t^{i,2}, \end{aligned}$$

where  $Y_t^{i,1} = \text{Re} Y_t^i$  and  $Y_t^{i,2} = \text{Im} Y_t^i$ .

**Example 9.4.** The following example shows that there exist Lie algebras with non trivial  $J_0^t$  for which the condition on the Lie bracket in Theorem 7.2 is satisfied. Let  $\mathfrak{g}$  be equal to  $Q^{1/2} HS$ , where the operator  $Q$  is defined as follows. We will view

the elements of  $HS$  as infinite matrices. Denote by  $e_{ij}$  an infinite matrix whose entries are all zero except the one equal to 1 at the intersection of the  $i$ th row and  $j$ th column. These matrices form an orthonormal basis of  $HS$ . We assume that  $Q$  is diagonal in this basis, namely,  $Qe_{ij} = e^{-(i+j)}e_{ij}$ . Note that this  $Q$  is a positive trace class operator, so we can construct the corresponding heat kernel measure. Thus the space  $\mathcal{H}^t(G_\infty)$  contains all holomorphic polynomials and therefore the space  $J_0^t$  is not trivial. Now let us verify that the condition on the Lie bracket is satisfied. For any  $x$  and  $y$  in  $\mathfrak{g}$

$$\begin{aligned} |xy|^2 &= \sum_{i,m} e^{i+m} \left( \sum_j x_{i,j} y_{j,m} \right)^2 \\ &= \sum_{i,m} \left( \sum_j e^{-j} e^{\frac{1}{2}(i+j)} x_{i,j} e^{\frac{1}{2}(j+m)} y_{j,m} \right)^2 \leq \sum_{i,m} \left( \sum_j e^{\frac{1}{2}(i+j)} x_{i,j} e^{\frac{1}{2}(j+m)} y_{j,m} \right)^2 \\ &\leq \sum_{i,m} \left( \sum_j e^{i+j} x_{i,j}^2 \right) \left( \sum_k e^{k+m} y_{k,m}^2 \right) = \sum_{i,j} e^{i+j} x_{i,j}^2 \sum_{m,k} e^{k+m} x_{m,k}^2 = |x|^2 |y|^2. \end{aligned}$$

Thus  $|[x, y]| \leq 2|x||y|$ . A similar construction can be done for the algebras considered in Example 9.1 and Example 9.2.

## REFERENCES

- [1] V.Bargmann, *On a Hilbert space of analytic functions and an associated integral transform, Part I*, Communications of Pure and Applied Mathematics, **24**, 1961, 187-214.
- [2] V.Bargmann, *Remarks on a Hilbert space of analytic functions*, Proc. of the National Academy of Sciences, **48**, 1962, 199-204.
- [3] G.DaPrato and J.Zabczyk, *Stochastic Equations in Infinite Dimensions*, Cambridge University Press, Cambridge, 1992.
- [4] B.Driver, *On the Kakutani-Itô-Segal-Gross and Segal-Bargmann-Hall isomorphisms*, J. of Funct. Anal., **133**, 1995, 69-128.
- [5] B.Driver and L.Gross, *Hilbert spaces of holomorphic functions on complex Lie groups*, Proceedings of the 1994 Taniguchi International Workshop (K. D. Elworthy, S. Kusuoka, I. Shigekawa, Eds.), World Scientific Publishing Co. Pte. Ltd, Singapore/New Jersey/London/Hong Kong, 1997.
- [6] E. B. Dynkin, *Calculation of the coefficients in the Campbell-Hausdorff formula*, Doklady Akad. Nauk SSSR (N.S.) **57**, 1947, 323-326.
- [7] M. Gordina, *Holomorphic functions and the heat kernel measure on an infinite dimensional complex orthogonal group*, to appear in Potential Analysis.
- [8] L. Gross and P. Malliavin, *Hall's transform and the Segal-Bargmann map*, Itô's Stochastic Calculus and Probability Theory, (M.Fukushima, N. Ikeda, H. Kunita and S. Watanabe, Eds.), Springer-Verlag, New York/Berlin, 1996.
- [9] B.Hall, *The Segal-Bargmann 'coherent state' transform for compact Lie groups*, J. of Funct. Anal., **122**, 1994, 103-151.
- [10] B. Hall and A. Sengupta, *The Segal-Bargmann transform for path-groups*, J. of Funct. Anal., **152**, 1998, 220-254.
- [11] O. Hijab, *Hermite functions on compact Lie groups I*, J. of Funct. Anal., **125**, 1994, 480-492.
- [12] O. Hijab, *Hermite functions on compact Lie groups II*, J. of Funct. Anal., **133**, 1995, 41-49.
- [13] Robert H. Martin, Jr., *Nonlinear operators and differential equations in Banach spaces*, Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1976.
- [14] I. Shigekawa, *Itô-Wiener expansions of holomorphic functions on the complex Wiener space*, in Stochastic Analysis, (E. Mayer et al, Eds.), Academic Press, San Diego, 1991, 459-473.

- [15] H.Sugita, *Properties of holomorphic Wiener functions-skeleton, contraction, and local Taylor expansion*, Probab. Theory Related Fields, **100**, 1994, 117–130.
- [16] H.Sugita, *Regular version of holomorphic Wiener function*, J. Math. Kyoto Univ., **34**, 1994, 849–857.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ON L8S  
4K1 CANADA

*E-mail address:* `gordina@math.McMaster.ca`