

involve

a journal of mathematics

Fair pricing and hedging under small perturbations of the
numéraire on a finite probability space

William Busching, Delphine Hintz,
Oleksii Mostovyi and Alexey Pozdnyakov



Fair pricing and hedging under small perturbations of the numéraire on a finite probability space

William Busching, Delphine Hintz,
Oleksii Mostovyi and Alexey Pozdnyakov

(Communicated by Robert B. Lund)

We consider the problem of fair pricing and hedging in the sense of Föllmer and Schweizer (1989) under small perturbations of the numéraire. We show that for replicable claims, the change of numéraire affects neither the fair price nor the hedging strategy. For nonreplicable claims, we demonstrate that is not the case. By reformulating the key stochastic control problem in a more tractable form, we show that both the fair price and optimal strategy are stable with respect to small perturbations of the numéraire. Further, our approach allows for explicit asymptotic formulas describing the fair price and hedging strategy's leading order correction terms. Mathematically, our results constitute stability and asymptotic analysis of a stochastic control problem under certain perturbations of the integrator of the controlled process, where constraints make this problem hard to analyze.

1. Introduction

In complete market models, the benchmark arbitrage-free pricing and hedging approach is based on replication, and it typically results in a unique price of a given security. It is proven in [Geman et al. 1995] that, in such settings, a change of numéraire affects neither pricing nor hedging. This result allows for more efficient pricing methodologies based on various changes of numéraire that are particularly evident for pricing and hedging of interest rate derivatives, where completeness of the underlying model is often embedded in the model assumption. We refer to [Brigo and Mercurio 2006] for more details.

MSC2020: primary 60G42, 60J74, 90C31, 91G20, 93E20; secondary 91G10.

Keywords: fair pricing, Föllmer–Schweizer decomposition, numéraire, stability, asymptotic analysis, conditional fair price.

This paper is a part of an REU project conducted in Summer 2021 at the University of Connecticut. The work was supported by the National Science Foundation under grants no. DMS-1950543 and DMS-1848339.

In incomplete markets, the situation is more complicated, in general. While there is still a class of derivative securities that is replicable, and the assertions of [Geman et al. 1995] apply to them, there are many other contingent claims that are nonreplicable and for which even the notion of a price becomes more complicated. As the arbitrage-free price is not unique for nonreplicable claims, other pricing methods have been introduced to overcome the nonuniqueness issue. Among them is fair pricing; see [Föllmer and Schweizer 1989]. It allows for regaining uniqueness of a (fair) price for a wide class of nonreplicable contingent claims.

As a numéraire is present in essentially every financial model, it is important to understand the response of the pricing and hedging methodologies to perturbations of the numéraire. In this work, we aim at understanding how fair pricing and hedging change under the small perturbations of numéraire. Working in settings with multiple stocks, we show that for replicable claims, fair pricing and hedging do not change, while for nonreplicable claims, we obtain explicit formulas for the first-order corrections for both the fair price and the hedging strategy. We work with a fairly general parametrization of perturbations of the numéraire. Also, regardless of the exact form of such perturbations, we show the stability of fair pricing and hedging with respect to small changes of numéraire.

Mathematically, we study the sensitivity of a solution of a stochastic control problem to small perturbations of the controlled process. In the settings of a finite probability space, we identify the conditions for the results to hold, which are represented via the invertibility of certain conditional covariance matrices. This condition is closely related to the nonredundancy of a given set of driving stochastic processes. In the core of our computations is a reformulation of this stochastic control problem from the one where we seek an optimal strategy among the ones satisfying the hard-to-deal with self-financing constraints to the one where these constraints become essentially vacuous. Our examples support the main results by showing that, for nonreplicable claims, both the fair price and hedging strategy change under a perturbation of the numéraire, as well as that the change of numéraire is fairly different from a seemingly related change of interest rate.

Our results complement the ones in [Monat and Stricker 1995], where the stability of fair pricing is established with respect to perturbations of the claim's payoff, as well as [Biagini and Pratelli 1999], by proving first-order corrections to the fair price and hedging strategy for nonreplicable claims in discontinuous stock-price settings. Historically, [Merton 1973] was one of the first to complete a change in numéraire, although the term numéraire was never formally defined in that work, and then [Margrabe 1978] focused on and clarified the term numéraire and its uses. The work [Geman et al. 1995] summarizes the history of the numéraire and gives convincing examples of its usages. We refer to [Karatzas and Kardaras 2021] for a recent literature overview and multiple contemporary results involving the concept of numéraire.

The remainder of this paper is organized as follows: In [Section 2](#), we specify the mathematical model, introduce the notions of fair pricing and hedging, and relate them to the Föllmer–Schweizer decomposition. In [Section 3](#), we present the notion of numéraire and establish some important related results that are subsequently used in [Section 4](#) to characterize fair pricing and hedging under a new tradable numéraire. We provide examples in [Section 5](#). In [Section 6](#), we consider the stability of our solution under small perturbations of the numéraire; in [Section 7](#), we obtain explicit formulas for the first-order correction terms of each component under small perturbations of the numéraire.

2. Fair pricing and hedging

This section aims to discuss the fair pricing and hedging in the settings with multiple stocks, making a digression and considering the case separately with one stock, where the computations become a bit simpler. Consider an economy indexed over discrete time, with uncertainty represented by a finite probability space $(\Omega, \mathbb{F}, \mathbb{P})$. The flow of information to all agents in this economy is represented by the filtration $\mathbb{F} = (\mathcal{F}_n)_{n=0,1,\dots,T}$ with fixed $T \in \mathbb{Z}^+$. Assume that \mathcal{F}_0 is trivial, containing only \emptyset and Ω , and that \mathcal{F}_T is the power set of Ω .

We begin by supposing there is a bank account S^0 with a price process equal to one at all times. We use this bank account as a numéraire in our introduction of the Föllmer–Schweizer decomposition, while noting that our subsequent analysis enables us to consider a more general tradable numéraire in the same circumstances. Let $S = (S_n)_{n=0,1,\dots,T}$ be a d -dimensional vector-valued \mathbb{F} -adapted process, i.e., each S_n is \mathcal{F}_n -measurable, and let S^i describe the evolution of the i -th stock, $i \in \{1, \dots, d\}$. We take S to describe the discounted price process of d stocks, and we denote the vector-valued increments of S by $\Delta S_n := S_n - S_{n-1}$ for $n = 1, \dots, T$.

Let $\xi = (\xi_n)_{n=1,\dots,T}$ represent a d -dimensional trading strategy corresponding to the number of shares of stocks held at any time n . We restrict the strategies to the ones that are *predictable* and *self-financing*, in the following senses: since our position in the stock market at time n must be chosen at time $n-1$, we say that ξ is *predictable*, namely, that ξ_n is \mathcal{F}_{n-1} -measurable for each $n \in \{1, \dots, T\}$. Let ξ^0 denote the positing in the bank account, and denoting $\bar{\xi} := (\xi^0, \xi)$, $\bar{S} := (1, S)$, we call a strategy $\bar{\xi}$ to be *self-financing* if

$$\bar{\xi}_n \cdot \bar{S}_n = \bar{\xi}_{n+1} \cdot \bar{S}_n, \quad n \in \{1, \dots, T-1\}, \quad (1)$$

where \cdot denotes the scalar product in \mathbb{R}^{d+1} ; below we also use the same symbol for the scalar product in \mathbb{R}^d .

Remark 2.1. Condition (1) implies that the accumulated gains and losses resulting from the asset price fluctuations are the only sources of changes in the portfolio

value. It can be restated in the equivalent forms

$$\bar{\xi}_{n+1} \cdot \bar{S}_{n+1} - \bar{\xi}_n \cdot \bar{S}_n = \bar{\xi}_{n+1} \cdot (\bar{S}_{n+1} - \bar{S}_n), \quad n \in \{1, \dots, T-1\}, \quad (2)$$

$$\bar{\xi}_n \cdot \bar{S}_n = \bar{\xi}_1 \cdot \bar{S}_0 + \sum_{k=1}^n \bar{\xi}_k \cdot (\bar{S}_k - \bar{S}_{k-1}), \quad n \in \{1, \dots, T\}, \quad (3)$$

where (3) follows from summation over (2) and $\bar{\xi}_1 \cdot \bar{S}_0$ is the initial wealth. We refer to [Föllmer and Schied 2016, p. 293] for more details. Below, we use the self-financing condition in the form (3).

Let Θ be the set of all predictable d -dimensional processes (that vacuously correspond to self-financing trading strategies). For $\xi \in \Theta$, we define the gains process

$$G_n(\xi) = \sum_{j=1}^n \xi_j \cdot \Delta S_j, \quad n \in \{1, \dots, T\}, \quad (4)$$

which can be thought of as gains in wealth up to time n for a self-financing trading strategy. Here and below, $\Delta W_n = W_n - W_{n-1}$ for every process W . An important observation is that, on the right-hand side of (4), as $S^0 \equiv 1$, one can equivalently use $\bar{\xi}_j \cdot \bar{\Delta S}_j$; however, the self-financing condition for such $\bar{\xi}$ must hold in this case.

Let V_0 denote the initial capital invested in the market at time $n = 0$. Then the total output from the trading process at time n is given by $V_0 + G_n(\xi)$. Now suppose there exists a derivative security that pays a value H at the final time T . To successfully hedge this security, we wish to trade in such a way that the trading output at time T is as close as possible to the payout of the derivative security. One way to ensure this is to minimize the expected quadratic cost incurred from hedging the security — namely, by solving the minimization problem

$$\min_{\substack{V_0 \in \mathbb{R} \\ \xi \in \Theta}} \mathbb{E}[(H - V_0 - G_T(\xi))^2]. \quad (5)$$

Following [Föllmer and Schweizer 1989], the V_0 in (5) is called *fair price*. We also call the optimal ξ the *fair price-based hedging strategy*. To characterize them, we introduce the discrete Föllmer–Schweizer decomposition, following [Föllmer and Schweizer 1989; Monat and Stricker 1995].

Theorem 2.2. *Let $S = M + A$ be the semimartingale decomposition of S into a martingale M and a predictable process A . Then every square-integrable and \mathcal{F}_N -measurable contingent claim H admits a decomposition*

$$H = V_0 + \sum_{k=1}^T \xi_k \cdot \Delta S_k + L_T \quad (6)$$

for some $V_0 \in \mathbb{R}$, a process $\xi \in \Theta$, and a martingale L , where

- (1) L and every component of M are orthogonal, meaning $\mathbb{E}[\Delta L_n \Delta M_n^i | \mathcal{F}_{n-1}] = 0$ for $n \in \{1, \dots, T\}$ and $i \in \{1, \dots, d\}$;
- (2) $\mathbb{E}[L_0] = 0$.

Following this decomposition, successful hedging of a contingent claim H requires the minimization of the unhedgeable L_T term. In turn, this is closely related to the concept of *local risk minimization*; see [Biagini and Pratelli 1999].

Remark 2.3. When S is a martingale, (6) is the Galtchouk–Kunita–Watanabe decomposition; see [Kunita and Watanabe 1967; Galtchouk 1975].

The Föllmer–Schweizer decomposition is crucial in obtaining optimal solutions ξ and V_0 to (5). Indeed, following [Föllmer and Schweizer 1989], for $d = 1$ (that is, with one risky asset), the recursive formula for ξ is given by

$$\xi_n = \frac{\text{Cov}_{\mathcal{F}_{n-1}}\left(H - \sum_{k=n+1}^T \xi_k \Delta S_k, \Delta S_n\right)}{\text{Var}_{\mathcal{F}_{n-1}}[\Delta S_n]}, \quad n \in \{1, \dots, T\}, \quad (7)$$

where $\text{Cov}_{\mathcal{F}_{n-1}}$ and $\text{Var}_{\mathcal{F}_{n-1}}$ denote the conditional covariance and variance, respectively.

It is this optimal strategy that we examine under a change of numéraire in the next section. Note that, in (7), we need the process $\text{Var}_{\mathcal{F}_{n-1}}[\Delta S_n]$, $n \in \{1, \dots, T\}$, to be *strictly* positive.

Remark 2.4. With multiple risky assets, this condition is replaced with the invertibility of conditional covariance matrices with probability 1. This condition is closely related to the nonredundancy of given stocks. However, the stability and asymptotic analysis are only imposed for the base model corresponding to $\epsilon = 0$.

3. Change of numéraire

To establish the machinery to change to a general numéraire, we begin by defining the set of general wealth processes starting from x to be

$$\mathcal{X}(x) := \left\{ x + G_n(\xi) = x + \sum_{k=1}^n \xi_k \cdot \Delta S_k \mid \xi \in \Theta, n \in \{0, \dots, T\} \right\}, \quad x \in \mathbb{R}. \quad (8)$$

A *numéraire*, most generally, can be defined as any *strictly* positive nondividend-paying asset. We focus on a tradable numéraire: that is, the ones where N is a strictly positive element of $\mathcal{X}(1)$. That is, N has the form

$$N_n = 1 + \sum_{k=1}^n \eta_k \cdot \Delta S_k, \quad n \in \{0, \dots, T\}, \quad (9)$$

for some $\eta \in \Theta$. The normalization condition $N_0 = 1$ is common in the literature.

The Föllmer–Schweizer decomposition discussed above utilizes an unchanging bank account as a numéraire; i.e., $N \equiv 1$ is constant at all times $[0, T]$. Indeed, a numéraire $N \equiv 1$ is implicit in many results in the field.

By a change in numéraire, we mean a change of units in which a price process of the traded securities, $\bar{S} = (1, S)$, is measured. Note that this process includes the bank account with value one across our time horizon as well as d stocks — all encoded in a vector \bar{S} . We denote the $(d+1)$ -dimensional price process of traded securities under a change of numéraire by

$$S^N := \left(\frac{1}{N}, \frac{S}{N} \right).$$

Under the new numéraire, it is natural to introduce the set of wealth processes analogous to (8); this is done in (10) below. To emphasize the self-financing constraints under a change of numéraire, first, we notice that one can rewrite (8) as

$$\mathcal{X}(x) = \left\{ x + \sum_{k=1}^n \bar{\xi}_k \cdot \Delta \bar{S}_k \mid \bar{\xi} \text{ is predictable and self-financing, } n \in \{0, \dots, T\} \right\}, \quad x \in \mathbb{R}.$$

This allows us to naturally extend (8) to the set of wealth processes under the numéraire N as

$$\mathcal{X}^N(x) := \left\{ x + \sum_{k=1}^n \bar{\xi}_k \cdot \Delta S_k^N \mid \bar{\xi} \text{ is predictable and self-financing, } n \in \{0, \dots, T\} \right\}, \quad x \in \mathbb{R}, \quad (10)$$

where the self-financing condition, analogous to (1), must now hold under the numéraire N , that is,

$$\bar{\xi}_n \cdot S_n^N = \bar{\xi}_{n+1} \cdot S_n^N, \quad 1 \leq n \leq T-1, \quad (11)$$

or, similarly, along the lines of Remark 2.1,

$$\bar{\xi}_n \cdot S_n^N = \bar{\xi}_1 \cdot S_0^N + \sum_{k=1}^n \bar{\xi}_k \cdot (S_k^N - S_{k-1}^N), \quad n \in \{1, \dots, T\}, \quad (12)$$

where $\bar{\xi}_1 \cdot S_0^N$ is the initial wealth.

We begin with a convenient result, analogous to Lemma 6.1 of [Mostovyi 2020], which demonstrates that wealth processes under a change of tradable numéraire adjust in an expected way. *In particular, the replicable claims stay replicable under a change of numéraire.* The proof of this result is similar to the proof of Lemma 4.2 below, and it is skipped.

Lemma 3.1. *Consider a stock price process under a change of numéraire $S^N = (1/N, S/N)$. Let us fix $x \in \mathbb{R}$ and consider the sets of wealth processes under the old and new numéraires $\mathcal{X}(x)$ and $\mathcal{X}^N(x)$ given by equations (8) and (10),*

respectively. Then, we have

$$\mathcal{X}^N(x) = \frac{\mathcal{X}(x)}{N} = \left\{ \frac{X}{N} = \left(\frac{X_n}{N_n} \right)_{n \in \{0, \dots, T\}} \mid X \in \mathcal{X}(x) \right\}. \quad (13)$$

4. Fair pricing and hedging and a numéraire change

We now turn to the question of applying the Föllmer–Schweizer decomposition-based hedging mechanism in an environment with a new numéraire. This has to be done with care. First, let us observe that the objective function in (5) can be rewritten as

$$\min_{X \in \bigcup_{x \in \mathbb{R}} \mathcal{X}(x)} \mathbb{E}[(H - X_T)^2]. \quad (14)$$

As the contingent claim H measured under the new numéraire to N is worth H/N , using the notation of the previous section, the natural formulation of (5) under N becomes

$$\min_{X^N \in \bigcup_{x \in \mathbb{R}} \mathcal{X}^N(x)} \mathbb{E} \left[\left(\frac{H}{N_T} - X_T^N \right)^2 \right]. \quad (15)$$

Recalling the definition of $\mathcal{X}^N(x)$ (including the self-financing condition (11)), the latter minimization problem (15) can be restated as

$$\begin{aligned} \text{minimize} \quad & \mathbb{E} \left[\left(\frac{H}{N_T} - \frac{V_0}{N_0} - \sum_{k=1}^T \bar{\xi}_k \cdot \Delta S_k^N \right)^2 \right], \\ \text{subject to} \quad & V_0 \in \mathbb{R}, \bar{\xi} \text{ is predictable and satisfies (11).} \end{aligned} \quad (16)$$

The optimal V_0 and $\bar{\xi}$ to (16) (whose existence is proven below) are defined to be the *fair price* and the *fair price-based hedging strategy* (or simply, the hedging strategy) under the numéraire N .

One can see that the self-financing constraint (11) enters problem (16) and makes it harder to analyze. Whereas in (5), we considered a risk-free asset, whose increments $\Delta S^0 \equiv 0$ at all times as a component of $\Delta \bar{S}$, and ΔS^N has no zero component, in general, since a tradable numéraire may change over time. In the change of numéraire case, the optimal $\bar{\xi}$ then additionally depends on the risk-free asset, which may evolve over time. The component of $\bar{\xi}$ corresponding to investment in the risk-free asset cannot, therefore, be chosen essentially after solving (16) in a way to make the optimal $\bar{\xi}$ self-financing, as the self-financing condition (11) is a constraint on our minimization problem (16). The following lemma demonstrates how we can bypass this issue by reformulating the minimization problem (16) in a more convenient way.

Let us consider

$$\min_{\substack{V_0 \in \mathbb{R} \\ \xi \in \Theta}} \mathbb{E} \left[\left(\frac{H - V_0 - \sum_{k=1}^T \xi_k \cdot \Delta S_k}{N_T} \right)^2 \right], \quad (17)$$

and let $W \in \bigcup_{x \in \mathbb{R}} \mathcal{X}(x)$ and $W^N \in \bigcup_{x \in \mathbb{R}} \mathcal{X}^N(x)$ denote the optimal wealth processes to (16) and (17), respectively.

Lemma 4.1. *Let N be a tradable numéraire. Then, (16) is equivalent to (17) in the sense that the objective functions are equal, and there is a one-to-one correspondence between the optimal wealth processes to (16) and (17), which are unique with probability 1, and we have*

$$W^N = \frac{W}{N}. \tag{18}$$

Proof. Observe that the expression $X^N := V_0 + \sum_{k=1}^T \bar{\xi}_k \cdot \Delta S_k^N$ is in $\mathcal{X}^N(V_0)$. Applying (13), we have $X^N = X/N$ for some X of the form $X_n = V_0 + \sum_{k=1}^n \xi_k \cdot \Delta S_k$, where $n \in \{0, \dots, T\}$, such that $X \in \mathcal{X}(V_0)$. Note that the ξ that gives rise to each wealth process may be different, but V_0 must be the same in both, due to the normalization condition $N_0 = 1$ and Lemma 3.1. Then (16), or rather the equivalent problem (15), becomes

$$\begin{aligned} \min_{X^N \in \bigcup_{V_0 \in \mathbb{R}} \mathcal{X}^N(V_0)} \mathbb{E} \left[\left(\frac{H}{N_T} - X_T^N \right)^2 \right] &= \min_{X \in \bigcup_{V_0 \in \mathbb{R}} \mathcal{X}(V_0)} \mathbb{E} \left[\left(\frac{H}{N_T} - \frac{X_T}{N_T} \right)^2 \right] \\ &= \min_{\substack{V_0 \in \mathbb{R} \\ \xi \in \Theta}} \mathbb{E} \left[\left(\frac{H - V_0 - \sum_{k=1}^T \xi_k \cdot \Delta S_k}{N_T} \right)^2 \right], \end{aligned} \tag{19}$$

which is (17). The chain of equalities above shows the objective functions in (16) and (17) are equal. Next, using the direct method from the calculus of variations and strict convexity of the function $x \mapsto x^2$, $x \in \mathbb{R}$, appearing in the objective, one can show the existence and uniqueness of the optimal self-financing wealth processes under the corresponding numéraires that are the minimizers to (16) and (17). The computations above, in particular (19) and Lemma 3.1, imply (18). \square

To emphasize the self-financing constraints, similarly to reformulation (16), one can restate (17) as

$$\begin{aligned} &\text{minimize} && \mathbb{E} \left[\left(\frac{H - V_0 - \sum_{k=1}^T \bar{\xi}_k \cdot \Delta \bar{S}_k}{N_T} \right)^2 \right], \\ &\text{subject to} && V_0 \in \mathbb{R}, \bar{\xi} \text{ is predictable and satisfies (1).} \end{aligned} \tag{20}$$

The following lemma establishes a relationship between the optimal hedging strategies for (16) and (20).

Lemma 4.2. *Let $W_n^N = V_0 + \sum_{k=1}^n \xi_k^{(1)} \cdot \Delta S_k^N$ and $W_n = V_0 + \sum_{k=1}^n \xi_k^{(2)} \cdot \Delta \bar{S}_k$, where $n \in \{0, \dots, T\}$, be the optimal self-financing wealth processes for (16)*

and (20), respectively. Then, we have

$$\xi_n^{(1)} \cdot \Delta \bar{S}_n = \xi_n^{(2)} \cdot \Delta \bar{S}_n, \quad n \in \{1, \dots, T\}, \quad (21)$$

$$\xi_n^{(1)} \cdot \Delta S_n^N = \xi_n^{(2)} \cdot \Delta S_n^N, \quad n \in \{1, \dots, T\}. \quad (22)$$

In particular, one can use the same strategy to optimize both (16) and (20).

Remark 4.3. Lemmas 4.1 and 4.2 assert that, for replicable claims (that is, the ones that are represented by a terminal value of an element of $\bigcup_{x \in \mathbb{R}} \mathcal{X}(x)$) change of numéraire affects neither the fair price nor the hedging strategy.

Proof of Lemma 4.2. As, by Lemma 4.1, $N_n W_n^N = W_n$, $n \in \{0, \dots, T\}$, we get

$$\Delta W_n = \Delta(W_n^N N_n),$$

that is,

$$\begin{aligned} \xi_n^{(2)} \cdot \Delta \bar{S}_n &= W_{n-1}^N \Delta N_n + N_{n-1} \Delta W_n^N + \Delta W_n^N \Delta N_n \\ &= W_{n-1}^N \Delta N_n + (N_{n-1} + \Delta N_n) \xi_n^{(1)} \cdot \Delta S_n^N \\ &= W_{n-1}^N \Delta N_n + \xi_n^{(1)} \cdot (N_{n-1} \Delta S_n^N + \Delta S_n^N \Delta N_n). \end{aligned} \quad (23)$$

As $\bar{S}_n = S_n^N N_n$, we get

$$\Delta \bar{S}_n = \Delta(S_n^N N_n) = S_n^N N_n - S_{n-1}^N N_{n-1} = S_{n-1}^N \Delta N_n + N_{n-1} \Delta S_n^N + \Delta S_n^N \Delta N_n,$$

and thus

$$N_{n-1} \Delta S_n^N + \Delta S_n^N \Delta N_n = \Delta \bar{S}_n - S_{n-1}^N \Delta N_n.$$

This allows us to rewrite (particularly, the last term in) (23) as

$$\begin{aligned} \xi_n^{(2)} \cdot \Delta \bar{S}_n &= W_{n-1}^N \Delta N_n + \xi_n^{(1)} \cdot (\Delta \bar{S}_{n-1} - S_{n-1}^N \Delta N_n) \\ &= \xi_n^{(1)} \cdot \Delta \bar{S}_n + (W_n^N - \xi_n^{(1)} \cdot S_n^N) \Delta N_n \\ &= \xi_n^{(1)} \cdot \Delta \bar{S}_n + \left(V_0 + \sum_{k=1}^n \xi_k^{(1)} \cdot \Delta S_k^N - \xi_n^{(1)} \cdot S_n^N \right) \Delta N_n. \end{aligned} \quad (24)$$

Notice that the self-financing condition for $\xi^{(1)}$, particularly in the form (12), implies

$$V_0 + \sum_{k=1}^n \xi_k^{(1)} \cdot \Delta S_k^N - \xi_n^{(1)} \cdot S_n^N = 0,$$

which is precisely the term in the parentheses in the last line of (24). This allows us to rewrite (24) as (21). We obtain (22) similarly. \square

Next, we apply Lemmas 4.1 and 4.2 to characterize Föllmer–Schweizer decomposition under a change of numéraire. For simplicity of notation, for random

variables X and Y , let us introduce

$$\mathcal{C}_{\mathcal{F}_n}^N(X, Y) := \mathbb{E}_{\mathcal{F}_n} \left[\left(\frac{X}{N_T} - \frac{\mathbb{E}_{\mathcal{F}_n}[X N_T^{-2}]}{N_T \mathbb{E}_{\mathcal{F}_n}[N_T^{-2}]} \right) \left(\frac{Y}{N_T} - \frac{\mathbb{E}_{\mathcal{F}_n}[Y N_T^{-2}]}{N_T \mathbb{E}_{\mathcal{F}_n}[N_T^{-2}]} \right) \right], \quad n \in \{1, \dots, T\}, \quad (25)$$

and consider the matrix-valued process

$$\mathcal{C}_n := (\mathcal{C}_{\mathcal{F}_{n-1}}^N(\Delta S_n^i, \Delta S_n^j))_{i,j=1,\dots,d}, \quad n \in \{1, \dots, T\}. \quad (26)$$

If \mathcal{C}_n is invertible with probability 1 for all n , we can define recursively, backward-in-time, vector-valued processes ξ and c as

$$\begin{aligned} \xi_n &:= [\mathcal{C}_n]^{-1} c_n, \quad n \in \{T, \dots, 1\}, \quad \text{where} \\ c_n &:= \begin{cases} (\mathcal{C}_{\mathcal{F}_{T-1}}^N(H, \Delta S_T^i))_{i=1,\dots,d} & \text{if } n = T, \\ (\mathcal{C}_{\mathcal{F}_{n-1}}^N(H - \sum_{k=n+1}^T \xi_k \cdot \Delta S_k, \Delta S_n^i))_{i=1,\dots,d} & \text{if } n \in \{T-1, \dots, 1\}. \end{cases} \end{aligned} \quad (27)$$

For ξ given by (27), let us also set

$$V_n = \mathbb{E}_{\mathcal{F}_n} \left[\left(H - \sum_{k=n+1}^T \xi_k \cdot \Delta S_k \right) \frac{N_T^{-2}}{\mathbb{E}_{\mathcal{F}_n}[N_T^{-2}]} \right], \quad n \in \{0, \dots, T-1\}. \quad (28)$$

Theorem 4.4. *Consider a model with T periods and d stocks. Let us consider \mathcal{C} , defined in (26), and assume that $[\mathcal{C}_n]^{-1}$ exists for every $n \in \{1, \dots, T\}$. Then V_0 , defined in (28), is the fair price, and ξ , defined in (27), is the (fair price-based) hedging strategy under the numéraire N , i.e., the optimizers to (17). We also have*

$$\frac{H}{N_T} = V_0 + \sum_{k=1}^T \xi_k \cdot \Delta S_k^N + L_T, \quad (29)$$

where L is the unhedgeable part that satisfies properties (1) and (2) in the statement of Theorem 2.2.

Remark 4.5. The process V defined in (28) can be thought as the *conditional fair price process under the numéraire N* , where V_n , $n \in \{0, \dots, T-1\}$, is the fair price under the information available up to time n . We can further extend V to T , by setting $V_T = H/N_T$. In particular, at $n = 0$, V_0 is the conditional fair price under trivial information represented by \mathcal{F}_0 . This is consistent with the (usual, or rather unconditional) fair price.

Remark 4.6. Invertibility of \mathcal{C}_n for every $n \in \{1, \dots, T\}$ is closely connected to the nonredundancy of d stocks.

Proof of Theorem 4.4. The proof proceeds recursively, backward in time. For brevity of the exposition, we focus on the main step and consider the minimization

problem at time $n \in \{0, \dots, T-1\}$, and if $n+1 < T$, assume we have already found ξ_{n+2}, \dots, ξ_T . Let us define

$$\tilde{V}_{n+1} := H - \sum_{k=n+2}^T \xi_k \cdot \Delta S_k. \quad (30)$$

Now, at time n , we want to minimize

$$\mathbb{E}_{\mathcal{F}_n} \left[\left(\frac{\tilde{V}_{n+1} - V_n - \xi_{n+1} \cdot \Delta S_{n+1}}{N_T} \right)^2 \right], \quad (31)$$

where the minimization is taken over all random variables V_n and ξ_{n+1} measurable with respect to \mathcal{F}_n .

Using the first-order conditions, we take the partial derivative of the objective function in (31) with respect to V_n , to obtain

$$\frac{\partial}{\partial V_n} \mathbb{E}_{\mathcal{F}_n} \left[\left(\frac{\tilde{V}_{n+1} - V_n - \xi_{n+1} \cdot \Delta S_{n+1}}{N_T} \right)^2 \right] = -2 \mathbb{E}_{\mathcal{F}_n} \left[\frac{\tilde{V}_{n+1} - V_n - \xi_{n+1} \cdot \Delta S_{n+1}}{N_T^2} \right] = 0,$$

and therefore, we get

$$V_n = \frac{\mathbb{E}_{\mathcal{F}_n} [\tilde{V}_{n+1} N_T^{-2}] - \xi_{n+1} \cdot \mathbb{E}_{\mathcal{F}_n} [\Delta S_{n+1} N_T^{-2}]}{\mathbb{E}_{\mathcal{F}_n} [N_T^{-2}]},$$

which, in view of (30), is exactly (28). Next, we substitute this V_n back into our objective function (31), and take partial derivatives with respect to each component of ξ to obtain, for $j \in \{1, \dots, d\}$,

$$\frac{\partial}{\partial \xi^j} \mathbb{E}_{\mathcal{F}_n} \left[\left(\frac{\tilde{V}_{n+1}}{N_T} - \frac{\mathbb{E}_{\mathcal{F}_n} [\tilde{V}_{n+1} N_T^{-2}]}{N_T \mathbb{E}_{\mathcal{F}_n} [N_T^{-2}]} - \xi_{n+1} \cdot \left(\frac{\Delta S_{n+1}}{N_T} - \frac{\mathbb{E}_{\mathcal{F}_n} [\Delta S_{n+1} N_T^{-2}]}{N_T \mathbb{E}_{\mathcal{F}_n} [N_T^{-2}]} \right) \right)^2 \right] = 0.$$

Computing these derivatives and using the notation specified in (25), we find that

$$\sum_{i=1}^d \xi_{n+1}^i c_{\mathcal{F}_n}^N(\Delta S^i, \Delta S^j) = c_{\mathcal{F}_n}^N(\tilde{V}_{n+1}, \Delta S^j), \quad j \in \{1, \dots, d\}.$$

Recalling the notation for \mathcal{C} in (26) and for \mathbf{c} in (27), we can rewrite the latter equation as

$$\mathcal{C}_{n+1} \xi_{n+1} = \mathbf{c}_{n+1}.$$

Now, using the assumed invertibility of \mathcal{C}_{n+1} , we have

$$\xi_{n+1} = [\mathcal{C}_{n+1}]^{-1} \mathbf{c}_{n+1},$$

which gives ξ in (27). Now, from Lemmas 4.1 and 4.2, one can show (29). \square




Figure 1. A one-period trinomial model.

5. Examples

To illustrate the results and to highlight some special features related to fair pricing under the change of numéraire, we consider the following examples, where we find the optimal trading strategy and the corresponding fair price using [Theorem 4.4](#). Key features of the results can be illustrated in a one-period trinomial model with one risky asset. Let the initial stock price S_0 equal 2. Also, let an increase in stock price happen by the factor of $u = 2$, a lack of movement be $c = 1$, and the down movement occur by the factor $d = \frac{1}{2}$ so that $uS_0 = 4$, $cS_0 = 2$, and $dS_0 = 1$ and so on; see [Figure 1](#). Let the probability of an up move occurring be $\frac{1}{6}$, the probability of a down move be $\frac{1}{3}$, and the probability of the stock staying steady be $\frac{1}{2}$.

The following example shows that for nonreplicable contingent claims, the fair price and the optimal strategy in the sense of optimization problem (17) change with N .

Example 5.1. We demonstrate the results of [Section 3](#) as applied to the trinomial model, obtaining the optimal hedging strategy and initial capital once a change of numéraire has been enacted.

(a) Consider a tradable numéraire given by $1 + \frac{1}{2}\Delta S$. Let H be a European call option which yields $\max\{0, K - S_1\}$, where S_1 is the value of the stock at time 1 and $K = 3$ is the strike price. Then, using [Theorem 4.4](#) for the one-period, one-stock case, we deduce

$$\xi_1 = \frac{C_{\mathcal{F}_0}^N(H, \Delta S_1)}{C_{\mathcal{F}_0}^N(\Delta S_1, \Delta S_1)} = \frac{\mathbb{E}\left[\frac{(H\mathbb{E}[N_1^{-2}] - \mathbb{E}[HN_1^{-2}])(\Delta S_1\mathbb{E}[N_1^{-2}] - \mathbb{E}[\Delta S_1N_1^{-2}])}{(N_1\mathbb{E}[N_1^{-2}])^2}\right]}{\mathbb{E}\left[\left(\frac{\Delta S_1\mathbb{E}[N_1^{-2}] - \mathbb{E}[\Delta S_1N_1^{-2}]}{N_1\mathbb{E}[N_1^{-2}]}\right)^2\right]}.$$

Computing the expectations using the values above gives

$$\mathbb{E}[N_1^{-2}] = \left(\frac{1}{1 + \frac{1}{2}(2)}\right)^2 \frac{1}{6} + \left(\frac{1}{1 + \frac{1}{2}(0)}\right)^2 \frac{1}{2} + \left(\frac{1}{\frac{1}{2}}\right)^2 \frac{1}{3} = \frac{45}{24}.$$

Substitution yields

$$\xi = \frac{75}{576} \approx 0.13021.$$

This would indicate that the optimal amount of stock to buy at time 0 is approximately 0.13021 of a share. The same process can be used to find V_0 . Calculating the expectations gives

$$V_0 = \frac{471}{4320} \approx 0.10903.$$

(b) *Comparison with optimal strategy when $N \equiv 1$.* For comparison, we now assume the numéraire $N \equiv 1$ and use the original one-period, one-stock formulas for the optimal trading strategy and initial value. The choice of $N \equiv 1$ simplifies the formulas for the optimal trading strategy ξ and the fair price V_0 to the following equations for the one-period, one-stock case:

$$\begin{aligned} \xi_1 &= \frac{\text{Cov}(H, \Delta S_1)}{\text{Var}(\Delta S_1)} = \frac{\mathbb{E}[(H - \mathbb{E}[H])(\Delta S_1 - \mathbb{E}[\Delta S_1])]}{\mathbb{E}[(\Delta S_1 - \mathbb{E}[\Delta S_1])^2]}, \\ V_0 &= \mathbb{E}[H] - \xi_1 \mathbb{E}[\Delta S_1]. \end{aligned} \quad (32)$$

Using the same method and information described above, the optimal trading strategy under the numéraire $N \equiv 1$ is $\frac{1}{3}$ and the fair price is $\frac{1}{6}$. Notice that this differs from the optimal strategy and fair price under a different numéraire. Thus, the choice of the numéraire affects the optimal strategy and the fair price.

The following example shows that changes of numéraire are different from the changes of interest rates, in general.

Example 5.2. There is the relatively common supposition that a change in the numéraire relates to a change in the interest rate, or at the very least, that perturbations of both act similarly. In this example, we include calculations on the interest rate in order to lay the groundwork for the understanding that perturbations of the interest rate and perturbations of the numéraire are not related, in general. In one-period settings, let us consider the model of the stock price as in [Example 5.1](#), and let us suppose that the interest rate is a constant $r > 0$ (instead of 0, as in part (b) of [Example 5.1](#)). Formulating the optimization problem similarly to (17), where the role of N is played by the bank account, leads to the computations performed in [\[Föllmer and Schweizer 1989, Section 2\]](#), which assert that the optimal ξ does not change compared to (32), whereas the fair price V_0 is given by

$$V_0 = \mathbb{E}\left[\frac{H}{1+r}\right] - \xi_1 \mathbb{E}\left[\frac{S_1}{1+r} - S_0\right],$$

which is different from V_0 specified through [Theorem 4.4](#), in general, even notationally, as r does not enter (17). This simple example shows that the perturbations of the interest rate are different from the perturbations of the numéraire.

6. Stability under perturbations of the numéraire

We now address the stability of the Föllmer–Schweizer decomposition under perturbations of the numéraire. Stability has already been shown for perturbations of V_T in [Monat and Stricker 1995] and for perturbations of S in [Boese et al. 2020]. We consider a family of \mathbb{F} -adapted strictly positive numéraire processes parametrized by ϵ , writing $(N^\epsilon)_{\epsilon \in (-\epsilon_0, \epsilon_0)}$ for some $\epsilon_0 > 0$. A specific example of such a family is given in the following section, but for now we only suppose that

$$\lim_{\epsilon \rightarrow 0} N^\epsilon_n(\omega) = N^0_n(\omega) = 1 \quad \text{for every } n \in \{0, \dots, T\} \text{ and } \omega \in \Omega. \quad (33)$$

Remark 6.1. We stress that, in (33) and below, the limits should be understood in the following sense: We fix n and ω , then we take a limit as $\epsilon \rightarrow 0$. Thus, the limit in (33) and other limits below hold for every $\omega \in \Omega$. In particular, the set of ω , for which (33) and other limits below exist, has probability 1.

Theorem 6.2. *For some $\epsilon_0 > 0$, let us consider a family of numéraire processes¹ $((N^\epsilon_n)_{n \in \{0, \dots, T\}})_{\epsilon \in (-\epsilon_0, \epsilon_0)}$ satisfying (33). Let us suppose that, for every $n \in \{1, \dots, T\}$, $(\text{Cov}_{\mathcal{F}_{n-1}}(\Delta S^i_n, \Delta S^j_n))_{i,j=1, \dots, d}$ is invertible with probability 1.² Then there exists $\bar{\epsilon}_0 \in (0, \epsilon_0]$ such that, for every random variable H and $\epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0)$, the assumptions of Theorem 4.4 are satisfied. Furthermore, the corresponding family of numéraire adjusted Föllmer–Schweizer decompositions³*

$$\frac{H}{N^\epsilon_T} = V_0^\epsilon + \sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} + L_T^\epsilon, \quad \epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0),$$

where ξ^ϵ are given via (27), satisfies

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \xi_n^\epsilon &= \xi_n^0, & n \in \{1, \dots, T\}, \\ \lim_{\epsilon \rightarrow 0} \Delta S_n^{N^\epsilon} &= \Delta \bar{S}_n, & n \in \{1, \dots, T\}, \\ \lim_{\epsilon \rightarrow 0} V_0^\epsilon &= V_0^0, \\ \lim_{\epsilon \rightarrow 0} L_n^\epsilon &= L_n^0, & n \in \{0, \dots, T\}. \end{aligned}$$

Proof. The proof goes recursively, backward in n . First, consider $n = T$. Since we are working on a finite probability space, via the definition of conditional expectation and (33), we get

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-1}}[N_T^\epsilon] = \mathbb{E}_{\mathcal{F}_{T-1}}[N_T^0] = 1,$$

¹Strict positivity for each $\epsilon \in (-\epsilon_0, \epsilon_0)$ is embedded in the definition of the numéraires.

²This conditional covariance matrix-valued process is \mathcal{C} defined in (26) for $N^0 \equiv 1$.

³These decompositions are given via (29) in Theorem 4.4.

and by continuity of $f(x) = x^{-2}$ on $(0, \infty)$, we obtain

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^\epsilon)^{-2}] = \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^0)^{-2}] = 1.$$

Moreover, this implies for random variables X and Y , both not depending on ϵ , that we have

$$\begin{aligned} & \lim_{\epsilon \rightarrow 0} \mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[X, Y] \\ &= \lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-1}} \left[\left(\frac{X}{N_T^\epsilon} - \frac{\mathbb{E}_{\mathcal{F}_{T-1}}[X(N_T^\epsilon)^{-2}]}{N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^\epsilon)^{-2}]} \right) \left(\frac{Y}{N_T^\epsilon} - \frac{\mathbb{E}_{\mathcal{F}_{T-1}}[Y(N_T^\epsilon)^{-2}]}{N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^\epsilon)^{-2}]} \right) \right] \\ &= \mathbb{E}_{\mathcal{F}_{T-1}} \left[\lim_{\epsilon \rightarrow 0} \left(\frac{X}{N_T^\epsilon} - \frac{\mathbb{E}_{\mathcal{F}_{T-1}}[X(N_T^\epsilon)^{-2}]}{N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^\epsilon)^{-2}]} \right) \left(\frac{Y}{N_T^\epsilon} - \frac{\mathbb{E}_{\mathcal{F}_{T-1}}[Y(N_T^\epsilon)^{-2}]}{N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^\epsilon)^{-2}]} \right) \right] \\ &= \mathbb{E}_{\mathcal{F}_{T-1}} \left[\left(\frac{X}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon} - \frac{\lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-1}}[X(N_T^\epsilon)^{-2}]}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon \lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^\epsilon)^{-2}]} \right) \right. \\ &\quad \left. \times \left(\frac{Y}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon} - \frac{\lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-1}}[Y(N_T^\epsilon)^{-2}]}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon \lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^\epsilon)^{-2}]} \right) \right] \\ &= \mathbb{E}_{\mathcal{F}_{T-1}} \left[\left(\frac{X}{N_T^0} - \frac{\mathbb{E}_{\mathcal{F}_{T-1}}[X(N_T^0)^{-2}]}{N_T^0 \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^0)^{-2}]} \right) \left(\frac{Y}{N_T^0} - \frac{\mathbb{E}_{\mathcal{F}_{T-1}}[Y(N_T^0)^{-2}]}{N_T^0 \mathbb{E}_{\mathcal{F}_{T-1}}[(N_T^0)^{-2}]} \right) \right] \\ &= \mathbb{E}_{\mathcal{F}_{T-1}}[(X - \mathbb{E}_{\mathcal{F}_{T-1}}[X])(Y - \mathbb{E}_{\mathcal{F}_{T-1}}[Y])] \\ &= \text{Cov}_{\mathcal{F}_{T-1}}[X, Y], \end{aligned} \tag{34}$$

which holds for every $\omega \in \Omega$. Thus, the continuity of $\mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[X, Y]$ in ϵ , and invertibility of $\mathcal{C}_n^{N_n^0}$, for every $n \in \{1, \dots, T\}$, imply that there exists $\bar{\epsilon}_0 \in (0, \epsilon_0)$ such that $\mathcal{C}_n^{N_n^\epsilon}$ is invertible for every $\epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0)$ and every $n \in \{1, \dots, T\}$. Such invertibility also implies continuity of $[(\mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[\Delta S_T^i, \Delta S_T^j])_{i,j=1}^d]^{-1}$ at $\epsilon = 0$, that is,

$$\lim_{\epsilon \rightarrow 0} [(\mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[\Delta S_T^i, \Delta S_T^j])_{i,j=1}^d]^{-1} = [(\mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^0}[\Delta S_T^i, \Delta S_T^j])_{i,j=1}^d]^{-1}.$$

Consequently, from (34), we deduce that

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \xi_T^\epsilon &= \lim_{\epsilon \rightarrow 0} [(\mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[\Delta S_T^i, \Delta S_T^j])_{i,j=1}^d]^{-1} (\mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[H, \Delta S_T^i])_{i=1}^d \\ &= [(\lim_{\epsilon \rightarrow 0} \mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[\Delta S_T^i, \Delta S_T^j])_{i,j=1}^d]^{-1} (\lim_{\epsilon \rightarrow 0} \mathcal{C}_{\mathcal{F}_{T-1}}^{N_T^\epsilon}[H, \Delta S_T^i])_{i=1}^d \\ &= [(\text{Cov}_{\mathcal{F}_{T-1}}[\Delta S_T^i, \Delta S_T^j])_{i,j=1}^d]^{-1} (\text{Cov}_{\mathcal{F}_{T-1}}[H, \Delta S_T^i])_{i=1}^d = \xi_T^0 \end{aligned} \tag{35}$$

for every $\omega \in \Omega$. If $T = 1$, this completes the proof for the stability of ξ . If $T > 1$, defining

$$A_n^\epsilon := H - \sum_{k=n+1}^T \xi_k^\epsilon \cdot \Delta S_k^{N_k^\epsilon}, \quad n \in \{0, \dots, T-1\}, \quad \epsilon \in (-\epsilon_0, \epsilon_0),$$

from (35), we get

$$\lim_{\epsilon \rightarrow 0} A_{T-1}^\epsilon = A_{T-1}^0, \quad \omega \in \Omega.$$

Consequently, as with (34), we obtain

$$\begin{aligned} & \lim_{\epsilon \rightarrow 0} \mathcal{C}_{\mathcal{F}_{T-2}}^{N_T^\epsilon} [A_{T-1}^\epsilon, \Delta S_{T-1}] \\ &= \lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-2}} \left[\left(\frac{A_{T-1}^\epsilon}{N_T^\epsilon} - \frac{\mathbb{E}_{\mathcal{F}_{T-2}}[A_{T-1}^\epsilon (N_T^\epsilon)^{-2}]}{N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-2}}[(N_T^\epsilon)^{-2}]} \right) \left(\frac{\Delta S_{T-1}}{N_T^\epsilon} - \frac{\mathbb{E}_{\mathcal{F}_{T-2}}[\Delta S_{T-1} (N_T^\epsilon)^{-2}]}{N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-2}}[(N_T^\epsilon)^{-2}]} \right) \right] \\ &= \mathbb{E}_{\mathcal{F}_{T-2}} \left[\left(\frac{\lim_{\epsilon \rightarrow 0} A_{T-1}^\epsilon}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon} - \frac{\lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-2}}[A_{T-1}^\epsilon (N_T^\epsilon)^{-2}]}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-2}}[(N_T^\epsilon)^{-2}]} \right) \right. \\ & \quad \times \left. \left(\frac{\Delta S_{T-1}}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon} - \frac{\lim_{\epsilon \rightarrow 0} \mathbb{E}_{\mathcal{F}_{T-2}}[\Delta S_{T-1} (N_T^\epsilon)^{-2}]}{\lim_{\epsilon \rightarrow 0} N_T^\epsilon \mathbb{E}_{\mathcal{F}_{T-2}}[(N_T^\epsilon)^{-2}]} \right) \right] \\ &= \mathbb{E}_{\mathcal{F}_{T-2}}[(A_{T-1}^0 - \mathbb{E}_{\mathcal{F}_{T-2}}[A_{T-1}^0])(\Delta S_{T-1} - \mathbb{E}_{\mathcal{F}_{T-2}}[\Delta S_{T-1}])] \\ &= \text{Cov}_{\mathcal{F}_{T-2}}[A_{T-1}^0, \Delta S_{T-1}], \quad \omega \in \Omega, \end{aligned}$$

and as with (35), we obtain the vector equation

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \xi_{T-1}^\epsilon &= \lim_{\epsilon \rightarrow 0} [(\mathcal{C}_{\mathcal{F}_{T-2}}^{N_T^\epsilon} [\Delta S_{T-1}^i, \Delta S_{T-1}^j])_{i,j=1}^d]^{-1} (\mathcal{C}_{\mathcal{F}_{T-2}}^{N_T^\epsilon} [A_{T-1}^\epsilon, \Delta S_{T-1}^i])_{i=1}^d \\ &= [(\lim_{\epsilon \rightarrow 0} \mathcal{C}_{\mathcal{F}_{T-2}}^{N_T^\epsilon} [\Delta S_{T-1}^i, \Delta S_{T-1}^j])_{i,j=1}^d]^{-1} (\lim_{\epsilon \rightarrow 0} \mathcal{C}_{\mathcal{F}_{T-2}}^{N_T^\epsilon} [A_{T-1}^\epsilon, \Delta S_{T-1}^i])_{i=1}^d \\ &= [(\text{Cov}_{\mathcal{F}_{T-2}} [\Delta S_{T-1}^i, \Delta S_{T-1}^j])_{i,j=1}^d]^{-1} (\text{Cov}_{\mathcal{F}_{T-2}} [A_{T-1}^0, \Delta S_{T-1}^i])_{i=1}^d \\ &= \xi_{T-1}^0, \quad \omega \in \Omega. \end{aligned}$$

Proceeding in this manner, one can show

$$\lim_{\epsilon \rightarrow 0} \xi_n^\epsilon = \xi_n^0, \quad n \in \{1, \dots, N\}, \quad \omega \in \Omega. \quad (36)$$

We also obtain

$$\lim_{\epsilon \rightarrow 0} \Delta S_n^{N^\epsilon} = \lim_{\epsilon \rightarrow 0} \left(\frac{\bar{S}_n}{N_n^\epsilon} - \frac{\bar{S}_{n-1}}{N_{n-1}^\epsilon} \right) = \frac{\bar{S}_n}{\lim_{\epsilon \rightarrow 0} N_n^\epsilon} - \frac{\bar{S}_{n-1}}{\lim_{\epsilon \rightarrow 0} N_{n-1}^\epsilon} = \bar{S}_n - \bar{S}_{n-1} = \Delta \bar{S}_n,$$

giving us, via (36), the equality

$$\lim_{\epsilon \rightarrow 0} \sum_{k=1}^n \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} = \sum_{k=1}^n \xi_k^0 \cdot \Delta \bar{S}_k, \quad n \in \{1, \dots, T\}, \quad \omega \in \Omega. \quad (37)$$

Therefore, by taking the expectation of

$$\frac{H}{N_T^\epsilon} = V_0^\epsilon + \sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} + L_T^\epsilon, \quad (38)$$

and using $\lim_{\epsilon \rightarrow 0} H/N^\epsilon = H$, $\mathbb{E}[L_T^\epsilon] = 0$, and (37), we get $\lim_{\epsilon \rightarrow 0} V_0^\epsilon = V_0^0$. Consequently, from convergence of the left-hand side in (38) to H , convergence of V_0^ϵ to V_0^0 , and (37), we obtain $\lim_{\epsilon \rightarrow 0} L_T^\epsilon = L_T^0$. Finally, using the martingale condition $\mathbb{E}_{\mathcal{F}_n}[L_T^\epsilon] = L_n^\epsilon$ on L^ϵ , we conclude the proof with

$$\lim_{\epsilon \rightarrow 0} L_n^\epsilon = L_n^0, \quad n \in \{0, \dots, T\}. \quad \square$$

7. Asymptotic analysis

In order to quantify how the fair price and trading strategy respond to numéraire perturbations, we introduce a (linear) parametrization of a tradable numéraire given by

$$N_n^\epsilon = 1 + \epsilon \sum_{k=1}^n \eta_k \cdot \Delta S_k, \quad \epsilon \in (-\epsilon_0, \epsilon_0), \quad (39)$$

where $\eta \in \Theta$, and ϵ_0 is chosen so that $N^\epsilon > 0$ for every $\epsilon \in (-\epsilon_0, \epsilon_0)$. Note that this satisfies (33), and so Theorem 6.2 is used throughout this section. Now we define the processes

$$N'_n := \lim_{\epsilon \rightarrow 0} \frac{N_n^\epsilon - N_n^0}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{1 + \epsilon \sum_{k=1}^n \eta_k \cdot \Delta S_k - 1}{\epsilon} = \sum_{k=1}^n \eta_k \cdot \Delta S_k, \quad n \in \{0, \dots, T\}.$$

For every $\epsilon \in (-\epsilon_0, \epsilon_0)$, we denote by \mathcal{C}^ϵ , \mathbf{c}^ϵ , and ξ^ϵ the processes defined in (26) and (27), respectively, corresponding to the numéraire N^ϵ . We also set

$$J'_n := -2N'_T + 2\mathbb{E}_{\mathcal{F}_n}[N'_T],$$

$$\begin{aligned} C'_n(X, Y) := & -\mathbb{E}_{\mathcal{F}_n}[X J'_n(Y - \mathbb{E}_{\mathcal{F}_n}[Y])] - \mathbb{E}_{\mathcal{F}_n}[Y J'_n(X - \mathbb{E}_{\mathcal{F}_n}[X])] \\ & - 2\mathbb{E}_{\mathcal{F}_n}[(X - \mathbb{E}_{\mathcal{F}_n}[X])N'_T(Y - \mathbb{E}_{\mathcal{F}_n}[Y])], \end{aligned}$$

$$\mathcal{C}'_n := (C'_{n-1}(\Delta S_n^i, \Delta S_n^j))_{i,j \in \{1, \dots, d\}}, \quad n \in \{1, \dots, T\}.$$

For $n = T$, we introduce

$$\mathbf{c}'_T := (C'_{T-1}(H, \Delta S_T^i))_{i \in \{1, \dots, d\}}, \quad (40)$$

$$\xi'_T := [\mathcal{C}'_T]^{\mathbf{c}'_T}(\mathbf{c}'_T - \mathcal{C}'_T \xi_T^0). \quad (41)$$

Continuing recursively, backward in time, for every $n \in \{T-1, \dots, 1\}$, we define

$$A'_{n+1} = - \sum_{k=n+1}^T \xi'_k \cdot \Delta \bar{S}_k + \sum_{k=n+1}^T \xi_k^0 \cdot \Delta (\bar{S}N')_k,$$

where $\bar{S}N'$ is a vector-valued stochastic process $(N', S^1 N', \dots, S^d N')$,

$$\tilde{C}'_n(X, Y) := C'_n(X, Y) + \mathbb{E}_{\mathcal{F}_n}[(A'_{n+1} - \mathbb{E}[A'_{n+1}])(Y - \mathbb{E}_{\mathcal{F}_n}[Y]),$$

$$\mathbf{c}'_n := \left(\tilde{\mathcal{C}}'_{n-1} \left(H - \sum_{k=n+1}^T \xi_k^0 \cdot \Delta \bar{S}_k, \Delta S_T^i \right) \right)_{i \in \{1, \dots, d\}}, \quad (42)$$

$$\xi'_n := [\mathcal{C}_n^0]^{-1} (\mathbf{c}'_n - \mathcal{C}'_n \xi_n^0), \quad n \in \{T-1, \dots, 1\}. \quad (43)$$

The following theorem gives the first-order corrections to the fair price, the hedging strategy, and the unhedgeable component under small perturbations of the numéraire.

Theorem 7.1. *Consider a family of numéraire processes $((N_n^\epsilon)_{n \in \{0, \dots, T\}})_{\epsilon \in (-\epsilon_0, \epsilon_0)}$ given by (39). Let us suppose that $(\text{Cov}_{\mathcal{F}_{n-1}}(\Delta S_n^i, \Delta S_n^j))_{i,j=1, \dots, d}$ is invertible for every $n \in \{1, \dots, T\}$ with probability 1.⁴ Then, for every H , there exists $\bar{\epsilon}_0 \in (0, \epsilon_0]$ such that for every $\epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0)$ with probability 1 we have*

$$\frac{H}{N_T^\epsilon} = V_0^\epsilon + \sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} + L_T^\epsilon, \quad (44)$$

where ξ^ϵ are given via (27) with $N = N^\epsilon$. The first-order corrections to the optimal trading strategy ξ_n , fair price V_0 , and unhedgeable component L_n are given by

$$\lim_{\epsilon \rightarrow 0} \frac{\xi_n^\epsilon - \xi_n^0}{\epsilon} = \xi'_n, \quad n \in \{1, \dots, T\}, \quad (45)$$

where ξ'_n , $n \in \{1, \dots, T-1\}$, is given by (43) and ξ'_T is specified in (41),

$$\lim_{\epsilon \rightarrow 0} \frac{V_0^\epsilon - V_0^0}{\epsilon} = \mathbb{E} \left[\sum_{k=1}^T \xi_k^0 \cdot \Delta (\bar{S} N')_k - H_T N'_T - \sum_{k=1}^T \xi'_k \cdot \Delta \bar{S}_k \right], \quad (46)$$

and

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \frac{L_n^\epsilon - L_n^0}{\epsilon} &= \mathbb{E}_{\mathcal{F}_n} \left[\sum_{k=1}^T \xi_k^0 \cdot \Delta (\bar{S} N')_k \right] - \mathbb{E} \left[\sum_{k=1}^T \xi_k^0 \cdot \Delta (\bar{S} N')_k \right] \\ &\quad - \left(\mathbb{E}_{\mathcal{F}_n} [H_T N'_T] - \mathbb{E} [H_T N'_T] \right. \\ &\quad \left. + \mathbb{E}_{\mathcal{F}_n} \left[\sum_{k=1}^T \xi'_k \cdot \Delta \bar{S}_k \right] - \mathbb{E} \left[\sum_{k=1}^T \xi'_k \cdot \Delta \bar{S}_k \right] \right), \quad n \in \{0, \dots, T\}. \end{aligned} \quad (47)$$

Proof. The proof parallels the proof of [Boese et al. 2020, Theorem 6.3], so, for brevity of the exposition, we only outline the main steps. We observe that invertibility of $[\mathcal{C}_n^0]^{-1}$, $n \in \{1, \dots, T\}$, and the argument in the proof of Theorem 6.2 imply that there exists $\bar{\epsilon}_0 \in (0, \epsilon_0]$ such that $[\mathcal{C}_n^\epsilon]^{-1}$ are invertible for every $n \in \{1, \dots, T\}$ and $\epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0)$. This implies that the assertions of 4.4 apply for every $\epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0)$, and therefore (44) holds. To show equation (45), we proceed recursively, backward in time, where (45) follows from direct computations.

⁴This condition is the same as in Theorem 6.2. Again, we only impose it for the base model corresponding to $\epsilon = 0$.

Now, we show (46). As $\mathbb{E}[L_T^\epsilon] = 0$, for every $\epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0)$, taking the expectation in (44), we deduce that

$$V_0^\epsilon = \mathbb{E} \left[\frac{H}{N_T^\epsilon} - \sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} \right], \quad \epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0). \quad (48)$$

One can see that

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \left(\mathbb{E} \left[\frac{H}{N_T^\epsilon} \right] - \mathbb{E} \left[\frac{H}{N_T^0} \right] \right) = \mathbb{E}[HN_T'], \quad (49)$$

and

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \mathbb{E} \left[\sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} - \sum_{k=1}^T \xi_k^0 \cdot \Delta S_k^{N^0} \right] = \mathbb{E} \left[\sum_{k=1}^T \xi_k' \cdot \Delta \bar{S}_k - \sum_{k=1}^T \xi_k^0 \cdot \Delta (\bar{S}N')_k \right], \quad (50)$$

Therefore, using (49) and (50), with (48), we deduce that (46) holds.

Finally, we show (47). Again, we start from (44), which we can rewrite as

$$L_T^\epsilon = \frac{H}{N_T^\epsilon} - V_0 - \sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon}. \quad (51)$$

Since L^ϵ is a \mathbb{P} -martingale, for every $\epsilon \in (-\bar{\epsilon}_0, \bar{\epsilon}_0)$, from (51), we obtain

$$L_n^\epsilon = \mathbb{E}_{\mathcal{F}_n}[L_T^\epsilon] = -V_0 + \mathbb{E}_{\mathcal{F}_n} \left[\frac{H}{N_T^\epsilon} - \sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} \right]. \quad (52)$$

For every $\omega \in \Omega$ and $n \in \{0, \dots, T\}$, one can see that

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \mathbb{E}_{\mathcal{F}_n} \left[\frac{H}{N_T^\epsilon} - \frac{H}{N_T^0} \right] = -\mathbb{E}_{\mathcal{F}_n}[HN_T'] \quad (53)$$

and

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \mathbb{E}_{\mathcal{F}_n} \left[\sum_{k=1}^T \xi_k^\epsilon \cdot \Delta S_k^{N^\epsilon} - \sum_{k=1}^T \xi_k^0 \cdot \Delta S_k^0 \right] = \mathbb{E}_{\mathcal{F}_n} \left[\sum_{k=1}^T \xi_k' \cdot \Delta \bar{S}_k - \sum_{k=1}^T \xi_k^0 \cdot \Delta (\bar{S}N')_k \right]. \quad (54)$$

Therefore, from (52), using (53) and (54), we obtain (47). \square

References

- [Biagini and Pratelli 1999] F. Biagini and M. Pratelli, “Local risk minimization and numéraire”, *J. Appl. Probab.* **36**:4 (1999), 1126–1139. [MR](#) [Zbl](#)
- [Boese et al. 2020] S. Boese, T. Cui, S. Johnston, G. Molino, and O. Mostovyi, “Stability and asymptotic analysis of the Föllmer–Schweizer decomposition on a finite probability space”, *Involve* **13**:4 (2020), 607–623. [MR](#) [Zbl](#)
- [Brigo and Mercurio 2006] D. Brigo and F. Mercurio, *Interest rate models: theory and practice*, 2nd ed., Springer, 2006. [MR](#) [Zbl](#)

- [Föllmer and Schied 2016] H. Föllmer and A. Schied, *Stochastic finance*, 4th ed., De Gruyter, Berlin, 2016. [MR](#)
- [Föllmer and Schweizer 1989] H. Föllmer and M. Schweizer, “Hedging by sequential regression: an introduction to the mathematics of option trading”, *ASTIN Bull.* **18** (1989), 147–160.
- [Galtchouk 1975] L. I. Galtchouk, “The structure of certain martingales”, pp. 9–33 in *Proceedings of the School and Seminar on the Theory of Random Processes, Part I* (Druskininkai, 1974), edited by B. Grigelionis and A. Širjaev, 1975. In Russian. [MR](#)
- [Geman et al. 1995] H. Geman, N. El Karoui, and J.-C. Rochet, “Changes of numéraire, changes of probability measure and option pricing”, *J. Appl. Probab.* **32**:2 (1995), 443–458. [MR](#) [Zbl](#)
- [Karatzas and Kardaras 2021] I. Karatzas and C. Kardaras, *Portfolio theory and arbitrage: a course in mathematical finance*, Graduate Studies in Mathematics **214**, Amer. Math. Soc., Providence, RI, 2021. [MR](#) [Zbl](#)
- [Kunita and Watanabe 1967] H. Kunita and S. Watanabe, “On square integrable martingales”, *Nagoya Math. J.* **30** (1967), 209–245. [MR](#) [Zbl](#)
- [Margrabe 1978] W. Margrabe, “The value of an option to exchange one asset for another”, *The Journal of Finance* **33**:1 (1978), 177–186.
- [Merton 1973] R. C. Merton, “Theory of rational option pricing”, *Bell J. Econom. and Management Sci.* **4** (1973), 141–183. [MR](#) [Zbl](#)
- [Monat and Stricker 1995] P. Monat and C. Stricker, “Föllmer–Schweizer decomposition and mean-variance hedging for general claims”, *Ann. Probab.* **23**:2 (1995), 605–628. [MR](#) [Zbl](#)
- [Mostovyi 2020] O. Mostovyi, “Asymptotic analysis of the expected utility maximization problem with respect to perturbations of the numéraire”, *Stochastic Process. Appl.* **130**:7 (2020), 4444–4469. [MR](#) [Zbl](#)

Received: 2021-08-10

Revised: 2022-01-07

Accepted: 2022-01-19

wbuschin@bowdoin.edu

Bowdoin College, Brunswick, ME, United States

hintzdelphine@gmail.com

Bethany Lutheran College, Mankato, MN, United States

oleksii.mostovyi@uconn.edu

Department of Mathematics, University of Connecticut, Storrs, CT, United States

alexey.pozdnyakov@uconn.edu

University of Connecticut, Storrs, CT, United States

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Chi-Kwong Li	College of William and Mary, USA
Martin Bohner	Missouri U of Science and Technology, USA	Robert B. Lund	Clemson University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Gaven J. Martin	Massey University, New Zealand
Scott Chapman	Sam Houston State University, USA	Steven J. Miller	Williams College, USA
Joshua N. Cooper	University of South Carolina, USA	Frank Morgan	Williams College, USA
Michael Dorff	Brigham Young University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joel Foisy	SUNY Potsdam, USA	Ken Ono	Univ. of Virginia, Charlottesville
Amanda Folsom	Amherst College, USA	Jonathon Peterson	Purdue University, USA
Stephan R. Garcia	Pomona College, USA	Vadim Ponomarenko	San Diego State University, USA
Anant Godbole	East Tennessee State University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Ron Gould	Emory University, USA	Józeph H. Przytycki	George Washington University, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Glenn H. Hurlbert	Virginia Commonwealth U, USA	Ann Trenk	Wellesley College, USA
Michael Jablonski	University of Oklahoma, USA	Ravi Vakil	Stanford University, USA
Nathan Kaplan	University of California, Irvine, USA	John C. Wierman	Johns Hopkins University, USA
David Larson	Texas A&M University, USA		

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2022 is US \$220/year for the electronic version, and \$290/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2022 Mathematical Sciences Publishers

involve

2022

vol. 15

no. 4

Convergence of random polygon sequences	547
INCHEOUL CHUNG, XINGPING SUN AND SONGFENG ZHENG	
Realizing Artin–Schreier covers with minimal a -numbers in positive characteristic	559
FIONA ABNEY-MCPEEK, HUGO BERG, JEREMY BOOHER, SUN MEE CHOI, VIKTOR FUKALA, MIROSLAV MARINOV, THEO MÜLLER, PAWEŁ NARKIEWICZ, RACHEL PRIES, NANCY XU AND ANDREW YUAN	
Ribbonlength of families of folded ribbon knots	591
ELIZABETH DENNE, JOHN CARR HADEN, TROY LARSEN AND EMILY MEEHAN	
A ternary additive problem involving fractional powers	629
RAYMOND LUO AND GANG YU	
Extensions of Rolle’s theorem	641
LAURA J. BATTS, MEGAN E. MORAN AND COURTNEY K. TAYLOR	
Fair pricing and hedging under small perturbations of the numéraire on a finite probability space	649
WILLIAM BUSCHING, DELPHINE HINTZ, OLEKSII MOSTOVYI AND ALEXEY POZDNYAKOV	
Family sizes for complete multipartite graphs	669
DANIELLE GREGG, THOMAS W. MATTMAN, ZACHARY PORAT AND GEORGE TODD	
Elliptic Harnack inequality for \mathbb{Z}^d	687
SIVA ATHREYA, NITYA GADHIWALA AND RITVIK R. RADHAKRISHNAN	
Rank bias for elliptic curves mod p	709
KIMBALL MARTIN AND THOMAS PHARIS	