

SMOOTH BUMPS, A BOREL THEOREM AND PARTITIONS OF SMOOTH FUNCTIONS ON P.C.F. FRACTALS.

LUKE G. ROGERS, ROBERT S. STRICHARTZ, AND ALEXANDER TEPLYAEV

ABSTRACT. We provide two methods for constructing smooth bump functions and for smoothly cutting off smooth functions on fractals, one using a probabilistic approach and sub-Gaussian estimates for the heat operator, and the other using the analytic theory for p.c.f. fractals and a fixed point argument. The heat semigroup (probabilistic) method is applicable to a more general class of metric measure spaces with Laplacian, including certain infinitely ramified fractals, however the cut off technique involves some loss in smoothness. From the analytic approach we establish a Borel theorem for p.c.f. fractals, showing that to any prescribed jet at a junction point there is a smooth function with that jet. As a consequence we prove that on p.c.f. fractals smooth functions may be cut off with no loss of smoothness, and thus can be smoothly decomposed subordinate to an open cover. The latter result provides a replacement for classical partition of unity arguments in the p.c.f. fractal setting.

1. INTRODUCTION

Recent years have seen considerable developments in the theory of analysis on certain fractal sets from both probabilistic and analytic viewpoints [1, 14, 26]. In this theory, either a Dirichlet energy form or a diffusion on the fractal is used to construct a weak Laplacian with respect to an appropriate measure, and thereby to define smooth functions. As a result the Laplacian eigenfunctions are well understood, but we have little knowledge of other basic smooth functions except in the case where the fractal is the Sierpinski Gasket [19, 5, 20]. At the same time the existence of a rich collection of smooth functions is crucial to several aspects of classical analysis, where tools like smooth partitions of unity, test functions and mollifications are frequently used. In this work we give two proofs of the existence of smooth bump functions on fractals, one taking the probabilistic and the other the analytic approach. The probabilistic result (Theorem 2.9) is valid provided the fractal supports a heat operator with continuous kernel and sub-Gaussian bounds, as is known to be the case for many interesting examples [1, 2, 3] that include non-post-critically finite (non-p.c.f.) fractals such as certain Sierpinski carpets. In this setting it can also be used to cut off functions of a certain smoothness class, with some loss of smoothness (Theorem 2.7). By contrast the analytic method (Theorem 3.8) is applicable to the smaller class of self-similar p.c.f. fractals with a regular harmonic structure and Dirichlet energy in the sense of Kigami [14].

For p.c.f. fractals we use our result on the existence of bump functions to prove a Borel-type theorem, showing that there are compactly supported smooth functions

2000 *Mathematics Subject Classification.* Primary 28A80; Secondary 31C45, 60J60.
Research supported in part by the National Science Foundation, Grant DMS-0140194.
Research supported in part by the National Science Foundation, Grant DMS-0505622.

with prescribed jet at a junction point (Theorem 4.3). This gives a very general answer to a question raised in [19, 5], and previously solved only for the Sierpinski Gasket [20]. We remark, however, that even in this special case the results of [20] neither contain nor are contained in the theorem proven here, as the functions in [20] do not have compact support, while those here do not deal with the tangential derivatives at a junction point.

Finally we apply our Borel theorem to the problem of partitioning smooth functions. Multiplication does not generally preserve smoothness in the fractal setting [4], so the usual partition of unity method is not available. As a substitute for this classical tool we show that a smooth function can be partitioned into smooth pieces with supports subordinate to a given open cover (Theorem 5.1). This result plays a major role in a forthcoming paper on a theory of distributions for p.c.f. self-similar fractals [22].

Setting. Let X be a self-similar subset of \mathbb{R}^d (or more generally any complete metric space) in the sense that there is a finite collection of contractive similarities $\{F_j\}_{j=1}^N$ of the space and X is the unique compact set satisfying $X = \cup_{j=1}^N F_j(X)$. The sets $F_j(X)$ are the 1-cells, and for a word $w = (w_1, w_2, \dots, w_m)$ of length m we define $F_w = F_{w_1} \circ \dots \circ F_{w_m}$ and call $F_w(X)$ an m -cell. If w is an infinite word then we define $[w]_m$ to be its length m truncation and let $F_w(X) = \cap_m F_{[w]_m}(X)$, which is clearly a point in X . The map from infinite words to X is surjective but not injective, and the points of non-injectivity play an important role in understanding the connectivity properties of the fractal (see Section 1.6 of [14]). In particular there are critical points of the cover by the F_j , namely those x and y for which there are $j \neq k$ in $\{1, \dots, N\}$ such that $F_j(x) = F_k(y)$ (so $F_j(x)$ is a critical value). We call an infinite word w critical if $F_w(X)$ is a critical value, and then call \tilde{w} post-critical if there is $j \in \{1, \dots, N\}$ such that jw is critical. The boundary ∂X of X consists of all points $F_{\tilde{w}}(X)$ for which \tilde{w} is post-critical. In the case that the set of post-critical words is finite the fractal is called post-critically finite (p.c.f.) and we also use the notations $V_0 = \partial X$ and $V_m = \cup_w F_w(V_0)$, where the union is over all words of length m . The points in $(\cup_m V_m) \setminus V_0$ are called *junction points*. We shall always assume that V_0 contains at least two elements.

We suppose that X comes equipped with a self-similar probability measure μ , meaning that there are μ_1, \dots, μ_N such that the cell corresponding to $w = (w_1, \dots, w_m)$ has measure $\mu(F_w(X)) = \prod_{j=1}^m \mu_{w_j}$. In order to do analysis on X we assume that X admits a Dirichlet form \mathcal{E} , so \mathcal{E} is a closed quadratic form on $L^2(\mu)$ with the (Markov) property that if $u \in \text{dom}(\mathcal{E})$ then so is $\tilde{u} = u\chi_{0 < u < 1} + \chi_{u \geq 1}$ and $\mathcal{E}(\tilde{u}, \tilde{u}) \leq \mathcal{E}(u, u)$, where χ_A is the characteristic function of A . We will work only with self-similar Dirichlet forms, having the property that

$$(1.1) \quad \mathcal{E}(u, v) = \sum_{m\text{-words } w} r_w^{-1} \mathcal{E}(u \circ F_w, v \circ F_w)$$

where the factors r_j are called resistance renormalization factors and as usual $r_w = r_{w_1} \cdots r_{w_m}$. For convenience we restrict to the case of regular harmonic structures, in which $0 < r_j < 1$ for all j . In addition we assume \mathcal{E} has the property that $C(X) \cap \text{dom}(\mathcal{E})$ is dense both in $\text{dom}(\mathcal{E})$ with \mathcal{E} -norm and in the space of continuous functions $C(X)$ with supremum norm. We often refer to \mathcal{E} as the energy. If X is a nested fractal in the sense of Lindström [17] then such a Dirichlet form may be

constructed using a diffusion or a harmonic structure [16, 7, 23]. Other approaches may be found in [21, 15, 18, 13].

Using the energy and measure we produce a weak Laplacian by defining $f = \Delta u$ if $\mathcal{E}(u, v) = -\int f v d\mu$ for all $v \in \text{dom}(\mathcal{E})$ that vanish on ∂X . Then $-\Delta$ is a non-negative self-adjoint operator on $L^2(\mu)$. When $\Delta u \in C(X)$ we write $u \in \text{dom}(\Delta)$; this notation is continued inductively to define $\text{dom}(\Delta^k)$ for each k and then $\text{dom}(\Delta^\infty) = \bigcap_k \text{dom}(\Delta^k)$. We say f is smooth if $f \in \text{dom}(\Delta^\infty)$. On a p.c.f. fractal the weak Laplacian admits an additional pointwise description in which \mathcal{E} is a renormalized limit of energies \mathcal{E}_m corresponding to the finite graph approximations V_m and the Laplacian Δ is a renormalized limit of the associated graph Laplacians Δ_m . Details are in Section 3.7 of [14].

By standard results, existence of the Dirichlet form \mathcal{E} implies existence of a strongly continuous semigroup $\{P_t\}$ with generator $-\Delta$. Conversely if there is such a semigroup and it is self-adjoint then there is a corresponding Dirichlet form, so we could equally well begin with $\{P_t\}$ and construct \mathcal{E} (see [6, 8]). The Markov property of \mathcal{E} ensures that if $0 \leq u \leq 1$ μ -a.e. then also $0 \leq P_t u \leq 1$ μ -a.e.

If X is p.c.f. then there is a definition of boundary normal derivatives of a function in $\text{dom}(\Delta)$ and a Gauss-Green formula relating these to the integral of the Laplacian on X . The usual definition uses resistance-renormalized limits of the terms of the graph Laplacian that exist at the boundary point. If q_i is the boundary point of X that is fixed by F_i and r_i is the resistance factor corresponding to F_i we may define a normal derivative ∂_n at q_i and have a Gauss-Green formula as in Section 3.7 of [14] by

$$(1.2) \quad \partial_n u(q_i) = - \lim_{m \rightarrow \infty} r_i^{-m} \Delta_{m, q_i} u(q_i)$$

$$(1.3) \quad \sum_{q \in \partial X} (v(q) \partial_n u(q) - u(q) \partial_n v(q)) = \int_X (v \Delta u - u \Delta v) d\mu$$

where in (1.2) the quantity $\Delta_{m, q_i} u(q_i)$ is the graph Laplacian at q_i . (We distinguish this from the graph Laplacian at interior points because it may involve a sum over fewer edges – for example in the case of the Sierpinski gasket the graph Laplacian at interior vertices has four terms, but at boundary points there are only two.)

Normal derivatives may also be localized to cells, so that $\partial_{n, F_w(X)} u(F_w(q_i))$ is given by the limit in (1.2) but with $\Delta_{m, w} u(F_w(q_i))$ denoting the graph Laplacian restricted to edges that lie inside $F_w(X)$. It is then easy to see that if Δu exists and is continuous on each of finitely many cells that meet at $F_w(q_i)$, then it is continuous on their union if and only if the following conditions hold: u is continuous, Δu has a unique limit at $F_w(q_i)$, and the normal derivatives at $F_w(q_i)$ sum to zero. We call these the *matching conditions* for the Laplacian.

We shall have need of two other pieces of information about a p.c.f. fractal with regular harmonic structure. The first is that there is a Green's function $g(x, y) \geq 0$ that is continuous on $X \times X$ and has self-similar structure related to the discrete Greens function $\Psi(x, y)$ on $(V_1 \setminus V_0)^2$. According to Section 3.5 of [14]

$$(1.4) \quad g(x, y) = \lim_{m \rightarrow \infty} \sum_{k=0}^{m-1} \sum_{w \in W_k} r_w \Psi(F_w^{-1}(x), F_w^{-1}(y))$$

where W_k is the collection of words of length k but the sum is only over those w such that $F_w^{-1}(x)$ and $F_w^{-1}(y)$ make sense. Integration against (the negative of)

this Green's function gives the Green's operator $Gf(x) = -\int g(x,y)f(y)d\mu(y)$ and solves $\Delta Gf = f$ with Dirichlet boundary conditions. In particular we will make use of pointwise estimates of $g(x,y)$ that follow from (1.4). The second thing we need to know is an estimate on the oscillation of a harmonic function on a cell $F_w(X)$, details of which are in Section 3.2 and Appendix A of [14]. A harmonic function $h(x)$ is determined by its values on the boundary V_0 , and its values may be computed using harmonic extension matrices A_i via $h|_{F_w(V_0)} = A_{w_m} \cdots A_{w_1} \cdot h|_{V_0}$. The A_i are positive definite, have eigenvalue 1 on the constant function and second eigenvalue at most r_i . It follows immediately that the oscillation of $A_i h|_{V_0}$ is at most r_i when the oscillation of $h|_{V_0}$ is bounded by 1, and similarly that the oscillation of $h|_{F_w(X)}$ is at most r_w .

More details about analysis on self-similar p.c.f. fractals may be found in [14], or [26] in the special case of the Sierpinski Gasket. The lecture notes of Barlow [1] cover the probabilistic approach that begins with a diffusion semigroup; non-p.c.f. examples include Sierpinski carpets [2, 3]. Some of the general theory connecting Dirichlet forms, heat semigroups and spectral theory of the Laplacian is covered in [6, 8].

Smooth bump functions. In classical analysis on Euclidean spaces the usual bump functions to consider are of the form $u \in C^\infty$ with support in a specified open set Ω , bounds $0 \leq u \leq 1$ and the property $u \equiv 1$ on a specified compact $K \subset \Omega$. In the fractal setting described above it is not usually the case that a product of smooth functions is itself smooth (see [4], or Section 5 below), so there is less practical benefit to asking that our bump functions be identically 1 on K and we will not always do so. Nor is it always essential that $0 \leq u \leq 1$, though this is sometimes useful. For this reason we will use the term *smooth bump function* to mean a function $u \in \text{dom}(\Delta^\infty)$ with support in a specified set Ω and a bound $|u - 1| \leq \epsilon$ on a specified compact $K \subset \Omega$.

Suppose X is a p.c.f. self-similar fractal and we have a function $u \in \text{dom}(\Delta^\infty)$ with $|u - 1| \leq \epsilon$ on $K \subset X$ and such that $\Delta^k u$ and $\partial_n \Delta^k u$ vanish on V_0 for all k . Then for any word w we see that

$$u_w = \begin{cases} u \circ F_w^{-1} & \text{on the cell } F_w(X) \\ 0 & \text{elsewhere} \end{cases}$$

is a smooth bump function with support in $F_w(X)$ by the matching conditions for the Laplacian. For this reason we also use the term *smooth bump function* to refer to $u \in \text{dom}(\Delta^\infty)$ on X with $|u - 1| \leq \epsilon$ on K and which *vanishes to infinite order* at all points $q \in V_0$, by which last phrase we mean $\Delta^k u(q) = \partial_n \Delta^k u(q) = 0$ for all k .

2. A SMOOTH BUMP FROM THE HEAT OPERATOR

In this section (X, dist) is a metric space with Borel measure μ and a self-adjoint Neumann Laplacian Δ . Let P_t denote the associated heat operator at time t , and $\|\cdot\|_{2,2}$ be the operator norm on L^2 . For later use we note from the spectral representation $P_t = \int_0^\infty e^{-xt} dE_\Delta(x)$ that there is an estimate

$$(2.1) \quad \|\Delta^k P_t\|_{2,2} \leq c_k t^{-k}.$$

Our main result here is a procedure for using the heat flow on X to cut-off a smooth function with only a small loss in smoothness. For this purpose we require an assumption on Δ .

Define

$$(D(t, d))^2 = \sup \left\{ \int_L |P_t f|^2 : \|f\|_2^2 = 1 \text{ and } L \subset X \text{ with } \text{dist}(\text{Sppt}(f), L) \geq d \right\},$$

so that for any set L and any $f \in L^2$ with $\text{dist}(\text{Sppt}(f), L) \geq d$ we have

$$(2.2) \quad \left(\int_L |P_t f|^2 \right)^{1/2} \leq D(t, d) \|f\|_2.$$

Our assumption is that there is ϵ_0 so that for any $0 < \epsilon < \epsilon_0$ there is a decreasing sequence t_j with all $t_j < \frac{2}{\lambda}$ and $\sum_j t_j = T < \infty$, and such that for every non-negative integer k ,

$$(2.3) \quad \sum_{j=1}^{\infty} t_{j+1}^{-k} D(t_j, \epsilon 2^{-j}) = C_k < \infty.$$

It is worth noting that if this condition is satisfied then we may make $T > 0$ as small as we like, because $D(t, d_1) \leq D(t, d_2)$ if $d_1 \geq d_2$ and therefore

$$\begin{aligned} \sum_{j=1}^{\infty} t_{j+m+1}^{-k} D(t_{j+m}, \epsilon 2^{-j}) &\leq \sum_{j=1}^{\infty} t_{j+m+1}^{-k} D(t_{j+m}, \epsilon 2^{-(j+m)}) \\ &= \sum_{j'=m}^{\infty} t_{j'+1}^{-k} D(t_{j'}, \epsilon 2^{-j'}) \leq C_k \end{aligned}$$

so we may simply take m so that $\sum_1^{\infty} t_{j+m}$ is as small as desired and use the sequence $t'_j = t_{j+m}$. At the same time, this allows us to reduce a finite number of the C_k to be as small as we desire. We suppose $C_0 \leq \frac{1}{2}$.

The assumption is given in this somewhat artificial form so as to emphasize the estimates needed in the proof. It is a non-trivial assumption, but is known to be true in many examples. For instance, if $P_t f$ is given by integration against a kernel $p(t, x, y)$ that satisfies a sub-Gaussian upper bound

$$(2.4) \quad p(t, x, y) \leq \frac{\gamma_1}{t^{\alpha/\beta}} \exp \left(-\gamma_2 \left(\frac{\text{dist}(x, y)^\beta}{t} \right)^{1/(\beta-1)} \right)$$

then (2.3) may readily be verified for sequences t_j that decrease sufficiently rapidly, for example $t_j = j^{-j}$ or $t_j = e^{-j^2}$. Heat kernel bounds like (2.4) have been the subject of a great deal of research, not only on fractals but also on manifolds and graphs (see [9, 24, 10, 11] and the references therein). Here we satisfy ourselves with noting that they are known for many interesting examples, including various p.c.f. fractals [16, 7, 23, 12] and certain highly symmetric generalized Sierpinski carpets (which are not p.c.f.) [2, 3]. They are also valid on products of some fractals [25]. Later we shall see that requiring $p(t, x, y)$ be continuous and satisfy (2.4) is an appropriate choice when using our cutoff procedure to produce a smooth function.

Definition 2.1. The space of heat-smoothed functions $\mathcal{A}(X)$ on X is

$$\mathcal{A}(X) = \bigcup_{T>0} P_T(L^2(X))$$

Functions in \mathcal{A} are characterized by the property that when expanded with respect to an orthonormal basis of Laplacian eigenfunctions, the coefficients decay exponentially with respect to the sequence of eigenvalues. In particular $\mathcal{A}(X)$ contains the eigenfunctions and is thus dense in $L^2(X)$.

Theorem 2.2. *Let $\phi \in \mathcal{A}(X)$ be $\phi = P_T f$ for some $f \in L^2(X)$ with $\|f\|_2 \leq 1$, and $T > 0$. Let K be a fixed compact set. If $0 < \epsilon < \epsilon_0$ then there is a function v that is equal to ϕ on K , that vanishes outside the ϵ -neighborhood of K , and has $\Delta^k v \in L^2$ for all k with $\|\Delta^k v\|_2 \leq 7c_1 c_k C_k t_1^{-k} e^{2\lambda T}$.*

On a first reading of the proof, we suggest thinking of the case where $\phi \equiv 1$ and X has finite measure, so that $u_j \equiv 1$ on K_j at each step. In this case the following heuristic may be helpful.

Our goal is a function with two properties, the first of which is L^2 -smoothness, meaning that $\Delta^k v \in L^2$ for all k , and the second is the property of being $\equiv 1$ on K and $\equiv 0$ outside the ϵ -neighborhood of K , which we call the *characteristic* property. Beginning with a function u_1 which has the characteristic property but is not L^2 -smooth, we recursively apply a two step method. The first step smoothes u_1 by applying the heat operator for a small time t_2 to obtain $v_2 = P_{t_2} u_1$, which is smooth but does not have the characteristic property. The second step splits X into a neighborhood K_2 of K , and the complement L_2 of a larger neighborhood, as well as the region A_2 between, and sets $u_2 \equiv 1$ on K_2 , $u_2 \equiv 0$ on L_2 , and $u_2 \equiv v_2$ on A_2 .

What we have gained in passing from u_1 to u_2 is replacing the original abrupt drop of the characteristic function with the improved piece on A_2 , as illustrated on Figure 1. This argument is repeated inductively on nested annulus-like regions A_j , each time applying the heat kernel for a shorter time t_j to get $v_j = P_{t_j} u_{j-1}$ and then cutting v_j off to be constant outside A_j to get u_j .

It is unsurprising that this process converges in L^2 . What is perhaps unexpected is that $\Delta^k v$ converges in L^2 for all k , and this is where the estimate (2.3) is crucial. Essentially what is going on is that the “steepness” of the $(j+1)$ -th interpolant depends both on the height it must interpolate and the “width scale” on which it interpolates. The “width scale” depends on t_{j+1} through the norm $\|\Delta^k P_{t_{j+1}}\|_{2,2} \leq c_k t_{j+1}^{-k}$ for the “steepness” measured by Δ^k , but the height to be interpolated depends instead on how much P_{t_j} changed the function during the smoothing step, which is controlled by $D(t_j, \epsilon 2^{-j})$ because that is how much L^2 -norm can “leak” across the annulus A_j when we apply the heat flow. The series of terms $\Delta^k (v_{j+1} - v_j)$ is therefore essentially just $t_{j+1}^{-k} D(t_j, \epsilon 2^{-j})$, which converges by (2.3).

Proof. Our assumption implies that we can choose a sequence t_j with $\sum_j t_j = T$ and such that (2.3) holds.

Define neighborhoods K_j shrinking dyadically to K , neighborhoods L_j shrinking dyadically to the complement of the ϵ -neighborhood of K , and annular regions A_j between them as follows:

$$\begin{aligned} K_j &= \{x : \text{dist}(x, K) < \epsilon 2^{-j}\}, \\ L_j &= \{x : \text{dist}(x, K) > \epsilon(1 - 2^{-j})\}, \\ A_j &= \{x : \epsilon 2^{-j} \leq \text{dist}(x, K) \leq \epsilon(1 - 2^{-j})\} = X \setminus (K_j \cup L_j). \end{aligned}$$




FIGURE 1. Initial steps of the proof of Theorem 2.2 for a constant function.

Using these regions, and writing $T_j = \sum_2^j t_i$, inductively define for $j \geq 2$

$$\begin{aligned} u_1 &= f\chi_{K_1} \\ v_j &= P_{t_j}u_{j-1} \\ u_j(x) &= \begin{cases} P_{T_j}f & x \in K_j \\ v_j(x) & x \in A_j \\ 0 & x \in L_j \end{cases} \end{aligned}$$

where χ_{K_1} is the characteristic function of the set K_1 . It will be notationally convenient to set $t_1 = 0$ and $v_1 \equiv 0$, so that the formula for u_j is valid for $j = 1$.

From Lemma 2.6 we see that v_j converges to a function v such that $\Delta^k v \in L^2$ for all k . However u_j and v_j have the same limit by Lemma 2.5. Since $u_j \equiv 0$ outside the ϵ -neighborhood of K the same is true of v . Moreover, on K we know that

$$u_j = P_{T_j}f \rightarrow \phi$$

and we conclude that the function v has the asserted properties. The estimate for $\Delta^k v$ then follows immediately from (??) and the estimate (2.1), because the latter gives

$$\|\Delta^k v_2\|_2 = \|P_{t_1}u_1\|_2 \leq c_1 t_1^{-k} \|u_1\| \leq c_1 t_1^{-k}.$$

□

Lemma 2.3. For $j \geq 2$,

$$\|(u_j - v_j)\chi_{K_j}\|_2 \leq D(t_j, \epsilon 2^{-j})(1 + \|v_{j-1}\|_2).$$

Proof. From the fact that $P_t \phi = e^{-\lambda t} \phi$ we have for $j \geq 2$:

$$\begin{aligned} &P_{t_j}u_{j-1} - P_{T_j}f \\ &= P_{t_j} \left((P_{T_{j-1}}f(x))\chi(x)_{K_{j-1}} + v_{j-1}\chi(x)_{A_{j-1}} \right) - P_{t_j}P_{T_{j-1}}f \\ &= P_{t_j} \left(- (P_{T_{j-1}}f(x))\chi(x)_{A_{j-1} \cup L_{j-1}} + v_{j-1}\chi(x)_{A_{j-1}} \right). \end{aligned}$$

On K_j the above is the difference between v_j and u_j . Since the support of the function to which we apply the heat operator is in $A_{j-1} \cup L_{j-1}$ and is therefore at least $\epsilon 2^{-j}$ distance from A_j we can use (2.2) to see

$$\int_{K_j} |u_j - v_j|^2 \leq D(t_j, \epsilon 2^{-j})^2 \left\| \left(P_{T_{j-1}} f(x) \right) \chi(x)_{A_{j-1} \cup L_{j-1}} + v_{j-1}(x) \chi(x)_{A_{j-1}} \right\|_2^2$$

so

$$\begin{aligned} \|(u_j - v_j) \chi_{K_j}\|_2 &\leq D(t_j, \epsilon 2^{-j}) \left\| \left(P_{T_{j-1}} f(x) \right) \chi(x)_{A_{j-1} \cup L_{j-1}} + v_{j-1}(x) \chi(x)_{A_{j-1}} \right\|_2 \\ &\leq D(t_j, \epsilon 2^{-j}) (1 + \|v_{j-1}(x) \chi(x)_{A_{j-1}}\|_2) \end{aligned}$$

because P_t contracts L^2 norm and $\|f\|_2 \leq 1$. \square

Lemma 2.4. For $j \geq 1$

$$\|v_{j+1}\|_2 \leq \|u_j\|_2 \leq 1 + 4 \sum_{i=2}^j D(t_i, \epsilon 2^{-i}) \leq 3$$

Proof. Since P_t is an L^2 contraction, the first inequality is immediate. Note that the second is true for $j = 1$ because $\|u_1\|_2 \leq \|f\|_2 \leq 1$. We also know from our assumption (2.3) that $\sum_j D(t_j, \epsilon 2^{-j}) = C_0 \leq \frac{1}{2}$. Induction and Lemma 2.3 then imply

$$\begin{aligned} \|u_j\|_2 &= \|v_j \chi_{A_j}\|_2 + \|u_j \chi_{K_j}\|_2 \\ &\leq \|v_j \chi_{A_j}\|_2 + \|v_j \chi_{K_j}\|_2 + \|(u_j - v_j) \chi_{K_j}\|_2 \\ &\leq \|v_j\|_2 + D(t_j, \epsilon 2^{-j}) (1 + \|v_{j-1}\|_2) \\ &\leq \|u_{j-1}\|_2 + D(t_j, \epsilon 2^{-j}) (1 + \|u_{j-2}\|_2) \\ &\leq 1 + 4 \sum_{i=2}^{j-1} D(t_i, \epsilon 2^{-i}) + D(t_j, \epsilon 2^{-j}) (2 + 4 \sum_{i=2}^{j-2} D(t_i, \epsilon 2^{-i})) \\ &\leq 1 + 4 \sum_{i=2}^j D(t_i, \epsilon 2^{-i}) \leq 3 \end{aligned}$$

for $j \geq 2$. Of course when $j = 2$ the $\|u_{j-2}\|$ term does not appear, and the estimate is trivial when $j = 1$. \square

Lemma 2.5. For $j \geq 2$ we have $\|u_j - v_j\|_2 \leq 7D(t_j, \epsilon 2^{-j})$.

Proof. Note first that $u_j \equiv 0$ on L_j , so

$$\|(u_j - v_j) \chi_{L_j}\|_2^2 = \|v_j \chi_{L_j}\|_2^2 = \int_{L_j} |P_{t_j} u_{j-1}|^2 \leq (D(t_j, \epsilon 2^{-j}))^2 \|u_{j-1}\|_2^2$$

because the support of u_{j-1} is in $K_{j-1} \cup A_{j-1}$, so is separated from L_j by a distance of at least $\epsilon 2^{-j}$, and therefore (2.2) is applicable.

Since u_j and v_j coincide on A_j we have

$$\begin{aligned} \|u_j - v_j\|_2 &= \|(u_j - v_j) \chi_{K_j}\|_2 + \|(u_j - v_j) \chi_{L_j}\|_2 \\ &\leq D(t_j, \epsilon 2^{-j}) (1 + \|v_{j-1} \chi_{A_{j-1}}\|_2) + (D(t_j, \epsilon 2^{-j})) \|u_{j-1}\|_2 \\ &\leq 7D(t_j, \epsilon 2^{-j}) \end{aligned}$$

by Lemma 2.3 and Lemma 2.4. \square

Lemma 2.6. *For each non-negative integer k the sequence $\{\Delta^k v_j\}$ is L^2 -Cauchy.*

Proof. Recall $T_m = \sum_2^m t_j$. Writing $v_{m+1} = P_{t_{m+1}}(u_m - v_m) + P_{t_{m+1}}v_m$ and substituting for the second term one may readily verify that

$$v_m = P_{T_m}u_1 + \sum_{l=2}^{m-1} P_{T_m - T_l}(u_l - v_l).$$

For notational convenience we set $v_1 = 0$ so the sum begins at 1, and compute

$$\begin{aligned} \Delta^k(v_{m+1} - v_m) &= \sum_{l=1}^{m-1} \Delta^k(P_{T_{m+1} - T_l} - P_{T_m - T_l})(u_l - v_l) + \Delta^k P_{t_{m+1}}(u_m - v_m) \\ &= \sum_{l=1}^{m-1} (P_{t_{m+1}} - I)(\Delta^k P_{T_m - T_l})(u_l - v_l) + \Delta^k P_{t_{m+1}}(u_m - v_m). \end{aligned}$$

Now using $\|(P_t - I)w\|_2 \leq t\|\Delta w\|_2$ and $\|\Delta^k P_t\|_{2,2} \leq c_k t^{-k}$ we estimate the sum. It is convenient (and essentially no loss) to estimate $T_m - T_l \geq t_{l+1}$. Inserting the estimate of Lemma 2.5 we obtain

$$\begin{aligned} \|\Delta^k(v_{m+1} - v_m)\|_2 &\leq t_{m+1} \sum_{l=1}^{m-1} \left\| \Delta^{k+1}(P_{T_m - T_l})(u_l - v_l) \right\|_2 + \|\Delta^k P_{t_{m+1}}(u_m - v_m)\|_2 \\ &\leq t_{m+1} \sum_{l=1}^{m-1} c_{k+1} t_{l+1}^{-(k+1)} \|u_l - v_l\|_2 + c_k t_{m+1}^{-k} \|u_m - v_m\|_2 \\ &\leq 7t_{m+1} \sum_{l=1}^{m-1} c_{k+1} t_{l+1}^{-(k+1)} D(t_l, \epsilon 2^{-l}) + 7c_k t_{m+1}^{-k} D(t_m, \epsilon 2^{-m}) \end{aligned}$$

Summing over m under the assumption that $\sum_{m=l}^{\infty} t_{m+1} \leq t_l$ we obtain by Fubini

$$\begin{aligned} \sum_{m=1}^n \|\Delta^k(v_{m+1} - v_m)\|_2 &\leq \sum_{m=1}^n 7t_{m+1} \left(\sum_{l=1}^{m-1} c_{k+1} t_{l+1}^{-(k+1)} D(t_l, \epsilon 2^{-l}) \right) + 7c_k \sum_{m=1}^n t_{m+1}^{-k} D(t_m, \epsilon 2^{-m}) \\ &= 7c_{k+1} \sum_{l=1}^{n-1} \left(\sum_{m=l+1}^n t_{m+1} \right) \left(t_{l+1}^{-(k+1)} D(t_l, \epsilon 2^{-l}) \right) + 7c_k \sum_{m=1}^n t_{m+1}^{-k} D(t_m, \epsilon 2^{-m}) \\ &\leq 14c_{k+1} \sum_{l=1}^n t_{l+1}^{-k} D(t_l, \epsilon 2^{-l}) \end{aligned}$$

because $c_k \leq c_{k+1}$. This sum converges by the assumption (2.3). \square

The construction in Theorem 2.2 produces a function v satisfying $\Delta^k v \in L^2$ for all k . The question of whether v is actually smooth (i.e. has $\Delta^k v \in C(X)$ for all k) is more difficult. Given that the only way we have to verify the assumptions for Theorem 2.2 in specific examples is to use an estimate of the form (2.4), and that in many instances the heat kernel is known to be continuous, the following is essentially as useful.

Theorem 2.7. *Suppose that the heat flow P_t is given by integration against a continuous kernel $p(t, x, y)$ satisfying (2.4). Let $\phi \in \mathcal{A}(X)$ and K be a fixed compact set. For any $0 < \epsilon < \epsilon_0$ there is a smooth function v that is equal to ϕ on K and equal to zero outside the ϵ -neighborhood of K .*

Theorem 2.7 is proved by applying Theorem 2.2 and the following lemma.

Lemma 2.8. *If P_t is given by integration against a continuous kernel $p(t, x, y)$ satisfying (2.4) then Δ^{-k} maps $L^2(X, \mu)$ into the continuous functions $C(X)$ for all $k > \frac{\alpha}{\beta} - 1$. In particular, if f is such that $\Delta^k f \in L^2$ for all k , then f is smooth.*

Proof. The spectral representation of Δ^{-k} implies

$$\begin{aligned} \Delta^{-k} f(x) &= \int \int_0^\infty t^k p(t, x, y) dt f(y) d\mu(y) \\ &= \int \int_0^1 t^k p(t, x, y) dt f(y) d\mu(y) + \int \int_0^\infty (t+1)^k p(t+1, x, y) dt f(y) d\mu(y) \end{aligned}$$

however we may rewrite the second term using

$$\begin{aligned} \int_0^\infty (t+1)^k p(t+1, x, y) dt &= \int_0^\infty (t+1)^k \int p(1, x, z) p(t, z, y) d\mu(z) dt \\ &= \int p(1, x, z) \int_0^\infty (t+1)^k p(t, z, y) dt d\mu(z) \\ &= (\Delta_y + I)^{-k} p(1, x, y). \end{aligned}$$

By the inequalities of Hölder and Minkowski

$$\begin{aligned} & \left| \Delta^{-k} f(x) - \Delta^{-k} f(x') \right| \\ & \leq \|f\|_2 \left\| \int_0^1 t^k (p(t, x, y) - p(t, x', y)) dt \right\|_{L^2(y)} \\ & \quad + \|f\|_2 \left\| (\Delta_y + I)^{-k} (p(1, x, y) - p(1, x', y)) \right\|_{L^2(y)} \\ & \leq \|f\|_2 \int_0^1 t^k \|p(t, x, y) - p(t, x', y)\|_{L^2(y)} dt + \|f\|_2 \left\| p(1, x, y) - p(1, x', y) \right\|_{L^2(y)} \end{aligned}$$

because $(\Delta_y + I)^{-k}$ is a contraction of L^2 . Now $\|p(t, x, y) - p(t, x', y)\|_{L^2(y)} \rightarrow 0$ as $x \rightarrow x'$, for any fixed t , because $p(t, x, y)$ is continuous. This deals with the second term and shows that the integrand in the first term converges pointwise to zero. The latter is bounded by $t^k (\|p(t, x, \cdot)\|_2 + \|p(t, x', \cdot)\|_2)$, which is an L^1 function when $k > \frac{\alpha}{\beta} - 1$, because

$$\|p(t, x, y)\|_{L^2(y)} = \int p(t, x, y) p(t, x, y) d\mu(y) = p(2t, x, x) \leq Ct^{-\alpha/\beta}$$

by (2.4). The result follows by dominated convergence. \square


Corollary 2.9. *If $\mu(X) < \infty$ then for any compact set K and neighborhood U of K , there is a smooth non-negative function that is 1 on K and vanishes outside U .*

Proof. Apply the theorem to the constant function 1 and note that all of the u_j and v_j are non-negative, so the limit function is also. \square

3. A SMOOTH BUMP AS A FIXED POINT OF AN OPERATOR

To understand why it is sometimes possible to construct a smooth bump function on a self-similar set as a fixed point of an operator, we invite the reader to consider an elementary situation. Let $I = [0, 1]$ be the unit interval in \mathbb{R} . We may view I as a p.c.f. self-similar set under the contractions $f_0 = x/2$ and $f_1 = (x + 1)/2$. If μ is Lebesgue measure and \mathcal{E} is defined using a limit of a regular self-similar harmonic structures with resistance factors $1/2$ then we obtain the usual Dirichlet energy and Laplacian, and the normal derivatives are the outward-directed one-sided derivatives at 0 and 1 (see [14, 26] for details).

The intuition for our construction is as follows. Consider a symmetric smooth bump function u on the interval $I = [0, 1]$, for which $u \equiv 1$ on $[L, 1 - L]$ as shown in Figure 2. If we look at the graph of $\Delta u = d^2u/dx^2$ we obtain something that

FIGURE 2. The smooth bump function u .

looks like a constant multiple of Figure 3, which appears as if it could be assembled from rescaled copies of u according to a rule like


$$(3.1) \quad \Phi u = \begin{cases} u\left(\frac{2x}{L}\right) & \text{if } 0 \leq x \leq \frac{L}{2} \\ -u\left(\frac{2x}{L} - 1\right) & \text{if } \frac{L}{2} < x \leq L \\ 0 & \text{if } L < x < 1 - L \\ -u\left(\frac{2x-2}{L} + 2\right) & \text{if } 1 - L \leq x < 1 - \frac{L}{2} \\ u\left(\frac{2x-2}{L} + 1\right) & \text{if } 1 - \frac{L}{2} \leq x \leq 1 \end{cases}$$

so that we might hope there is actually a smooth bump function u which has precisely this scaling behavior. If we let G denote the Green's operator for the operator Δ on I with Dirichlet boundary conditions, then this would be equivalent to asking that u be a fixed point of the operator

$$(3.2) \quad \Psi u(x) = \frac{G \circ \Phi u(x)}{G \circ \Phi u(1/2)}$$

It is a consequence of our general result Theorem 3.8 that the operator Ψ in (3.2) has a fixed point and that the fixed point is a smooth bump function. In fact more is true in the special case of I , where the fact that removing any interior point disconnects the set, along with the existence of an explicit formula for the Green's function, allows us to prove that the fixed point has values in $[0, 1]$ and is identically 1 on $[L, 1 - L]$. For reasons of brevity we do not include the proof of this result; it is a simpler version of the proof of Theorem 3.8.

Proposition 3.1. *If L is sufficiently small then the operator Ψ preserves the space of continuous functions on I that have values in $[0, 1]$, vanish at 0 and 1, and are*

FIGURE 3. The function $\Delta u = d^2u/dx^2 = \Phi(u)$.

identically 1 on $[L, 1-L]$. Furthermore Ψ is a contraction in the L^∞ norm on these functions, and its fixed point is a smooth function that vanishes to infinite order at 0 and 1.


Another example in which we can define operators Φ and Ψ that are similar to (3.1) and (3.2) is the Sierpinski gasket SG with its standard harmonic structure and measure, where for sufficiently large l we can set

$$(3.3) \quad \Phi u(x) = \begin{cases} 2u(F_i^{-(l+1)}(x)) & \text{if } x \in F_i^{(l+1)}(SG) \\ -u(F_j^{-1} \circ F_i^{-l}(x)) & \text{if } x \in F_i^l \circ F_j(SG), j \neq i \\ 0 & \text{otherwise} \end{cases}$$

and with p any vertex from V_1 let

$$(3.4) \quad \Psi u = \frac{G \circ \Phi u(x)}{G \circ \Phi u(p)}$$

as illustrated in Figure 4 for the case $l = 2$. Again we omit the variant of the proof

FIGURE 4. Φu in the case $l = 2$.

of Theorem 3.8 that establishes the following result

Proposition 3.2. *The operator Ψ of (3.4) is an L^∞ -contractive self-map of the set of functions that are continuous on SG , vanish at the boundary, are identically 1 on $SG \setminus \cup_i F_i^l$, and satisfy $|\int u - 1| \leq \frac{1}{2}$. The fixed point of Ψ is a smooth bump function.*

The method described for I and SG rely heavily on the symmetry of these sets and on the assumption that they are endowed with the symmetrical harmonic structures and measure. This assumption is unavoidable if we want to use the same operation Φ at all steps of the computation, as the natural linear combination of rescaled copies of the function will not otherwise have the desired properties, but it is very restrictive. Even some of the simplest of the nested fractals defined by Lindström [17] have insufficient symmetry for a fixed Φ to be used in the construction of a smooth bump by this method. Nonetheless the method can be adapted to general p.c.f. self-similar fractals with regular harmonic structure and self-similar measure.


Let X be p.c.f. self-similar with boundary $V_0 = \partial X$, measure μ that is self-similar with scaling factors μ_j and regular harmonic structure with factors r_j . We fix a scale l_1 with size to be determined later, and label the boundary l_1 cells by $Y_j = F_j^{l_1}(X)$. Their union is $Y = \cup Y_j$. For any $\epsilon > 0$ we will build a smooth function that satisfies $|u - 1| \leq \epsilon$ on $X \setminus Y$ by a construction that inductively determines its Laplacian on the cells Y_j , writing it as a fixed point of an operator Ψ on the following space of functions.

Definition 3.3. Let \mathcal{C} be the space of continuous functions u on X such that $u(q) = 0$ for $q \in \partial X$ and $\|u - 1\|_1 \leq \frac{1}{2}$. Note that this space is non-empty and closed in the continuous functions with supremum norm.

To define the operator Ψ we need a little more notation. Let $S \subset V_{l_1}$ consist of those points that lie in some Y_j and in at least one other l_1 -cell. If l_1 is sufficiently large then no two of the Y_j can intersect; we assume this and see that the connected components of $X \setminus S$ are the cells Y_j (less points of S) and the set $X \setminus Y$. Label those boundary points of the cell Y_j at which Y_j intersects another l_1 cell by $x_{i,j}$ for $i = 1, \dots, I_j$. Fixing a second scale l_2 , also with size to be determined, we associate to each $x_{i,j}$ the unique $(l_1 + l_2)$ -cell in Y_j containing $x_{i,j}$, calling it $Z_{i,j}$. We also set $Z_{0,j} = F_j^{l_1+l_2}(X)$, so it is the $(l_1 + l_2)$ -cell in Y_j that contains $q_j \in V_0$, and define $w_{i,j}$ to be the word such that $F_{i,j}(X) = F_{w_{i,j}}(X) = Z_{i,j}$. Figure 5 illustrates our labelling conventions in the case $X = SG$, $l_1 = 2$ and $l_2 = 1$. We identify a particular function that is in \mathcal{C} when l_1 is large enough. Let f be the piecewise harmonic function on X with values $f(x_{i,j}) = 1$ for $i = 1, \dots, I_j$ and $j = 1, \dots, N$ but $f(x_{0,j}) = 0$ for all j . It is clear that f is continuous, identically 1 on $X \setminus Y$ and harmonic on each of the sets Y_j . It fails to be harmonic only at the points $x_{i,j}$ with $i \geq 1$, and we readily compute that the Laplacian of f is a measure supported at these points. In fact if δ_x denotes the Dirac mass at x then

$$(3.5) \quad \Delta f = \sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j} \delta_{x_{i,j}} = - \sum_{j=1}^N \sum_{i=1}^{I_j} \partial_n f_j(x_{i,j}) \delta_{x_{i,j}}$$

with the second equality coming from the matching conditions for the Laplacian. Since $f_j \circ F_j^{l_1}$ is harmonic on X with boundary values 0 at q_j and 1 at all q_i , $i \neq j$, it has normal derivatives bounded by a constant $C(r)$ that may be chosen so as to

FIGURE 5. Notation if $X = SG$, $l_1 = 2$ and $l_2 = 1$.

depend only on the harmonic structure. Scaling of the normal derivatives under $F_j^{l_1}$ therefore implies $|a_{i,j}| = |\partial_n f_j(x_{i,j})| \leq C(r)r_j^{-l_1}$.

The smooth bump function we seek will actually be a perturbation of f , constructed by iteratively replacing the Dirac masses in (3.5) by rescaled copies of the stage k bump, correcting for the boundary normal derivatives, and applying the Dirichlet Green's operator to obtain the stage $k + 1$ bump. We will see that each stage gains one order of smoothness, so the limiting function will be in $\text{dom}(\Delta^\infty)$. Our first step is to estimate the effect of a perturbation of the type described.

Lemma 3.4. *For each $j = 1, \dots, N$ and $i = 1, \dots, I_j$ let $\nu_{i,j}$ be a finite, signed Borel measure with support in $Z_{i,j}$. If we use the coefficients in (3.5) to define*

$$\nu = \sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j} \nu_{i,j}$$

and let $u = G(\nu)$ be the result of applying the Dirichlet Green's operator, then

$$(3.6) \quad |u(x)| \leq C(r)N^2 \sup_{i,j} \|\nu_{i,j}\| \quad \text{for all } x \in X$$

where $\|\nu_{i,j}\|$ is the total variation of $\nu_{i,j}$. If in addition we have $\int \nu_{i,j} = 0$ for all i and j then

$$(3.7) \quad |u(x)| \leq C(r)N \left(\sup_{i,j} \|\nu_{i,j}\| \right) \left(\sum_{k=1}^N r_k^{l_2} \right) \quad \text{for all } x \in X \setminus Y$$

Proof. Recall that G may be represented as integration against the continuous kernel $-g(x, y)$, with sign chosen so $g(x, y) \geq 0$. The estimates we desire follow from (1.4) and the fact that $|a_{i,j}| \leq C(r)r_j^{-l_1}$. The former ensures both that $|g(x, y)| \leq C(r)r_j^{l_1}$ on each Y_j and that the oscillation of $g(x, y)$ on $Z_{i,j}$ is at most

$C(r)r_{w_{i,j}}$. We compute

$$\begin{aligned}
|u(x)| &= \left| \int_X g(x, y) d\nu(y) \right| \\
&\leq \sum_{j=1}^N \sum_{i=1}^{I_j} |a_{i,j}| \int_{Z_{i,j}} |g(x, y)| d\nu_{i,j}(y) \\
&\leq \sum_{j=1}^N \sum_{i=1}^{I_j} C(r)r_j^{-l_1} \int_{Z_{i,j}} C(r)r_j^{l_1} d|\nu_{i,j}(y)| \\
&\leq C(r)N^2 \sup_{i,j} \|\nu_{i,j}\|
\end{aligned}$$

which establishes the first inequality. To obtain the second we observe that $\Delta u = 0$ on $X \setminus Y$, so by the maximum principle we need only verify the inequality at the points $x_{i,j}$. Fix such a point $x_{i',j'}$, and use that each $\int d\nu_{i,j} = 0$ to subtract the appropriate constant from each integrand before estimating:

$$\begin{aligned}
|u(x_{i',j'})| &= \left| \int_X g(x_{i',j'}, y) d\nu(y) \right| \\
&= \left| \sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j} \int_{Z_{i,j}} g(x_{i',j'}, y) d\nu_{i,j}(y) \right| \\
&= \left| \sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j} \int_{Z_{i,j}} \left(g(x_{i',j'}, y) - g(x_{i',j'}, x_{i,j}) \right) d\nu_{i,j}(y) \right| \\
&\leq \sum_{j=1}^N \sum_{i=1}^{I_j} C(r)r_j^{-l_1} \int_{Z_{i,j}} C(r)r_{w_{i,j}} d|\nu_{i,j}| \\
&\leq C(r) \sum_{j=1}^N \sum_{i=1}^{I_j} r_{k(i,j)}^{l_2} \|\nu_{i,j}\|
\end{aligned}$$

because $r_{w_{i,j}} = r_j^{l_1} r_k^{l_2}$ for some $k = k(i, j)$. Each $k(i, j)$ occurs at most once for a fixed j , so we obtain

$$|u(x_{i',j'})| \leq C(r) \left(\sup_{i,j} \|\nu_{i,j}\| \right) \left(\sum_{j=1}^N \sum_{k=1}^N r_k^{l_2} \right) \leq C(r)N \left(\sup_{i,j} \|\nu_{i,j}\| \right) \left(\sum_{k=1}^N r_k^{l_2} \right)$$

□

There is an analogous but simpler estimate for the effect of introducing a mass supported on one of the small cells $Z_{0,j}$ at the boundary. If $\nu_{0,j}$ is a finite, signed Borel measure with support in $Z_{0,j}$ we use (1.4) to see $|g(x, y)| \leq C(r)r_j^{l_1+l_2}$ on $Z_{0,j}$ and therefore

$$(3.8) \quad |G(\nu_{0,j})| \leq \left| \int_{Z_{0,j}} |g(x, y)| d|\nu_{0,j}(y)| \right| \leq C(r)r_j^{l_1+l_2} \|\nu_{0,j}\|$$

The ideas discussed so far allow us to generalize the definition of the operator Ψ in (3.2) and (3.4). The idea is that to replace the Dirac mass terms on the boundary cells $Z_{i,j}$ in (3.5) with normalized rescaled copies of $u \in \mathcal{C}$ and apply the

Green's operator to obtain a function that is near 1 on $X \setminus Y$. By adding some terms on the cells $Z_{0,j}$ we can make the result have vanishing normal derivatives at the boundary without changing the value on $X \setminus Y$ very much. In consequence we will obtain an operator that smooths $u \in \mathcal{C}$ to be in $\mathcal{C} \cap \text{dom}(\Delta)$ with vanishing normal derivatives and is near 1 on $X \setminus Y$. Iterating the operator will then produce a sequence of smoother and smoother bump functions.

Let

$$(3.9) \quad u_{i,j}(x) = \begin{cases} \mu(Z_{i,j})^{-1} \left(\int_X u d\mu \right)^{-1} \left(u \circ F_{i,j}^{-1}(x) \right) & \text{if } x \in Z_{i,j} \\ 0 & \text{otherwise} \end{cases}$$

so that each $u_{i,j}$ is continuous and has integral 1. Since $u \in \mathcal{C}$ we also have that $\int |u_{i,j}| \leq \left(\int u \right)^{-1} \|u\| \leq 3$.

Definition 3.5. The operator Ψ on \mathcal{C} is $\Psi u = G(v)$, where

$$(3.10) \quad v(x) = \sum_{j=1}^N \sum_{i=0}^{I_j} b_{i,j} r_j^{-l_1} u_{i,j}(x)$$

and G is the Dirichlet Green's operator. In this expression the coefficients for $i \geq 1$ are given by $b_{i,j} = r_j^{l_1} a_{i,j}$ with $a_{i,j}$ as in (3.5), but the $b_{0,j}$ are yet to be determined.

Note that $|b_{i,j}| \leq C(r)$ when $i \geq 1$. It is immediate that $G(v) \equiv 0$ on ∂X and is continuous. Moreover $\Delta G(v) = v$ is a linear combination of continuous functions, so $\Psi u \in \text{dom}(\Delta)$. The next lemma uses the Gauss-Green formula to reduce finding the correct $b_{0,j}$ to a problem in linear algebra.

Lemma 3.6. *If l_1 and l_2 are sufficiently large then there are values $b_{0,j}$ such that $\partial_n \Psi u \equiv 0$ on ∂X . The minimal sizes of l_1 and l_2 depend only on the harmonic structure of X and the number of vertices N in ∂X .*

Proof. Let h_j be the function that is harmonic on X , equal to 1 at q_j and 0 at all other points of ∂X . Using $\Delta G(v) = v$, $G(v) \equiv 0$ on ∂X , and the Gauss-Green formula

$$\begin{aligned} \int h_j(y) v(y) d\mu(y) &= \int h_j(y) \Delta G(v)(y) d\mu(y) \\ &= \sum_{q_k \in \partial X} h_j(q_k) (\partial_n G(v))(q_k) \\ &= (\partial_n G(v))(q_j) \end{aligned}$$

from which $\partial_n G(v) \equiv 0$ on ∂X is simply

$$0 = \int h_j(y) v(y) d\mu(y) = \sum_{i',j'} b_{i',j'} r_{j'}^{-l_1} \int h_j(y) u_{i',j'}(y) d\mu(y)$$

for all $j = 1, \dots, N$. Moving the terms depending on the fixed values $b_{i',j'}$ for $i' \geq 1$ this may be reformulated as

$$(3.11) \quad \sum_{j'} b_{0,j'} r_{j'}^{-l_1} \int h_j(y) u_{0,j'}(y) d\mu(y) = - \sum_{j'=1}^N \sum_{i'=1}^{I_{j'}} b_{i',j'} r_{j'}^{-l_1} \int h_j(y) u_{i',j'}(y) d\mu(y)$$

which we recognize as a matrix equation $\sum_{j'} M(u)_{j',j} r_{j'}^{-l_1} b_{0,j'} = A(u)_j$ with

$$(3.12) \quad (M(ud\mu))_{j',j} = \int h_j(y) u_{0,j'}(y) d\mu(y)$$

$$(3.13) \quad (A(ud\mu))_j = - \sum_{j'=1}^N \sum_{i'=1}^{I_{j'}} b_{i',j'} r_{j'}^{-l_1} \int h_j(y) u_{i',j'}(y) d\mu(y)$$

It is clear that we need to know $M = M(ud\mu)$ is invertible, but rather than prove this directly we do so by proving a perturbation estimate similar to Lemma 3.4 that will be useful later. To this end consider replacing each of the measures $u_{0,j'} d\mu$ in (3.12) with a copy of a different probability measure $d\sigma$ scaled and translated to give $d\sigma_{i',j'}$ supported on $Z_{i',j'}$. We call the result $M(d\sigma)$. The difference of these measures has mass zero, so we can compute an estimate involving the total variation of the measures

$$(3.14) \quad \begin{aligned} \left| M(ud\mu - d\sigma)_{j',j} \right| &= \left| \int h_j(y) (u_{0,j'}(y) d\mu(y) - d\sigma_{0,j'}(y)) \right| \\ &\leq \left| \int (h_j(y) - h_j(x_{0,j'})) (u_{0,j'}(y) d\mu(y) - d\sigma_{0,j'}(y)) \right| \\ &\leq C(r) r_{j'}^{l_1+l_2} \|ud\mu - d\sigma\| \end{aligned}$$

because h_j is harmonic and therefore varies by at most $C(r) r_{j'}^{l_1+l_2}$ on each $Z_{0,j'}$. In particular if the measures $d\sigma_{i',j'}$ are Dirac masses at the points $x_{0,j'}$ then $M(d\sigma)$ is simply the identity, so (3.14) implies

$$\left| (I - M)_{j',j} \right| \leq C(r) r_{j'}^{l_1+l_2}$$

from which M is invertible when l_1+l_2 is large, with $\|I - M^{-1}\| \leq C(N, r) \sum_j r_j^{l_1+l_2}$.

A similar perturbation argument can be made for $A(d\mu - d\sigma)$, where $A(d\sigma)$ is obtained by replacing each $u_{i',j'} d\mu$ by $d\sigma_{i',j'}$ in (3.13). Estimating the integral terms and using the bound $|b_{i',j'}| \leq C(r)$ we obtain

$$(3.15) \quad \left| A(ud\mu - d\sigma)_j \right| \leq C(N, r) \left(\sum_i r_i^{l_2} \right) \|ud\mu - d\sigma\|$$

however this is not the most useful thing we can do here. Instead we recognize that the bounds $|h_j(y) - 1| \leq C(r) r_j^{l_1}$ on Y_j and $|h_j(y)| \leq r_{j'}^{l_1}$ on $Y_{j'}$ for $j' \neq j$ ensure

$$\left| A(ud\mu)_j + \sum_{i'=1}^{I_j} b_{i',j} r_j^{-l_1} \right| \leq C(N, r)$$

so that combining this with our bound on $I - M^{-1}$ we have

$$(3.16) \quad \left| b_{0,j} - \sum_{i'=1}^{I_j} b_{i',j} \right| \leq C(N, r) r_j^{l_1}$$

If we examine the function f in (3.5) it is clear that the normal derivative at each point $x_{0,j}$ is $\sum_{i=1}^{I_j} b_{i,j}$, so our choice of $b_{0,j}$ is a small perturbation of that which would be used to cancel the normal derivatives of f . We also remark that this shows all $|b_{0,j}| \leq C(N, r)$. \square

If l_1 and l_2 are large enough then the values $b_{0,j}$ from Lemma 3.6 may be used to complete Definition 3.5 for the operator Ψ . Some key properties of this operator are summarized in the following lemma.

Lemma 3.7. *If l_1 and l_2 are sufficiently large then $\Psi(u) \in \mathcal{C} \cap \text{dom}(\Delta)$ and*

$$(3.17) \quad \|\Psi u\|_\infty \leq C_1$$

$$(3.18) \quad |\Psi u(y) - 1| \leq C_2 \sum_{j=1}^N r_j^{l_2} \quad \text{for all } y \in X \setminus Y$$

where C_1 , C_2 and the minimal sizes of l_1 and l_2 are constants depending only on the harmonic structure of X , the measure μ and the number of vertices N in ∂X .

Proof. Since

$$\Psi(u) = G \left(\sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j} u_{i,j} + \sum_{j=1}^N b_{0,j} r_j^{-l_1} u_{0,j} \right)$$

we obtain (3.17) from (3.6) and (3.8), and the fact that $|b_{0,j}| \leq C(N, r)$ for all j . The estimate (3.18) is only a little more difficult. Using $f(x) = 1$ and (3.5) on the set $X \setminus Y$ we see that

$$\begin{aligned} |\Psi u(x) - 1| &= |\Psi u(x) - f(x)| \\ &= \left| G \left(\sum_{j=1}^N \sum_{i=0}^{I_j} b_{i,j} r_j^{-l_1} u_{i,j} \right) - G \left(\sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j} \delta_{x_{i,j}} \right) \right| \\ &\leq \left| G \left(\sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j} (u_{i,j} d\mu - \delta_{x_{i,j}}) \right) \right| + \sum_{j=1}^N |b_{0,j} r_j^{-l_1} G(u_{0,j} d\mu)| \\ &\leq C(N, r) \sup_{i,j} \|u_{i,j} d\mu - \delta_{x_{i,j}}\| \left(\sum_{k=1}^N r_k^{l_2} \right) + C(N, r) \left(\sum_{j=1}^N r_j^{l_2} \right) \|u_{0,j} d\mu\| \\ (3.19) \quad &\leq C(N, r) \left(\sum_{j=1}^N r_j^{l_2} \right) \end{aligned}$$

where the estimate for the $b_{0,j}$ terms came from (3.8) and that for the $a_{i,j}$ terms is from (3.7) because $\int u_{i,j} d\mu = 1 = \int \delta_{x_{i,j}}$ and both are supported on $Z_{i,j}$.

Finally we check that $\|\Psi u - 1\|_1 \leq \frac{1}{2}$. Using the results we have so far

$$\begin{aligned} \|\Psi u - 1\|_1 &\leq \int_{X \setminus Y} C(N, r) \left(\sum_{j=1}^N r_j^{l_2} \right) d\mu + \int_Y (1 + C_1) d\mu \\ (3.20) \quad &\leq C(N, r) \left(\sum_{j=1}^N r_j^{l_2} \right) + (1 + C_1) \mu(Y) \end{aligned}$$

so that we can be sure $\Psi u \in \mathcal{C}$ if both l_1 and l_2 are sufficiently large, because $\mu(Y) \rightarrow 0$ as $l_1 \rightarrow \infty$. It has already been observed that $u \in \text{dom}(\Delta)$ and $u \equiv 0$ on ∂X , so the lemma is proven. \square

Finally we come to the main result of this section. The following theorem implements the idea that motivated our definition of Ψ , namely that Ψ smoothes

functions in \mathcal{C} and therefore its recursive application gives a bump function in $\text{dom}(\Delta^\infty)$.

Theorem 3.8. *Given $\epsilon > 0$ there are l_1 and l_2 sufficiently large that Ψ has a fixed point u_0 in \mathcal{C} with $|u - 1| \leq \epsilon$ on $X \setminus Y$. The fixed point is a smooth bump function and every $u \in \mathcal{C}$ has $\|\Psi^k u - u_0\|_\infty \rightarrow 0$ as $k \rightarrow \infty$.*

Proof. Let $u, \tilde{u} \in \mathcal{C}$. We calculate $\Psi u - \Psi \tilde{u} = G(v - \tilde{v})$, where v and \tilde{v} are as in (3.10). Beginning with a variant of the computation (3.19) we have

$$\begin{aligned}
|G(v - \tilde{v})(x)| &\leq \left| G\left(\sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j}(u_{i,j} - \tilde{u}_{i,j})\right) \right| + \left| G\left(\sum_{j=1}^N r_j^{-l_1}(b_{0,j}u_{0,j} - \tilde{b}_{0,j}\tilde{u}_{0,j})\right) \right| \\
&\leq \left| G\left(\sum_{j=1}^N \sum_{i=1}^{I_j} a_{i,j}(u_{i,j} - \tilde{u}_{i,j})\right) \right| + \sum_{j=1}^N r_j^{-l_1} |\tilde{b}_{0,j}| |G(u_{0,j} - \tilde{u}_{0,j})| \\
(3.21) \quad &\quad + \sum_{j=1}^N r_j^{-l_1} |b_{0,j} - \tilde{b}_{0,j}| |G(u_{0,j})(x)|
\end{aligned}$$

which suggests we will need to know estimates for both $(u_{i,j} - \tilde{u}_{i,j})$ and $|b_{0,j} - \tilde{b}_{0,j}|$. Conveniently we can reduce the latter to the former using (3.14) and (3.15), because $b_{0,j}$ and $\tilde{b}_{0,j}$ are computed from equations of the form $\sum_j M(ud\mu)_{j,i} b_{0,j} r_j^{-l_1} = A(ud\mu)_i$. We easily see that

$$r_j^{-l_1}(b_{0,j} - \tilde{b}_{0,j}) = \left(M(ud\mu)^{-1} A(ud\mu - \tilde{u}d\mu) \right) + \left(M(ud\mu)^{-1} M(\tilde{u}d\mu - ud\mu)(r_j^{-l_1} b_{0,j}) \right)$$

however by (3.14) we have both $\|M(ud\mu - \tilde{u}d\mu)\| \leq C(N, r)\|u - \tilde{u}\|_1 \sum_i r_i^{l_1+l_2}$ and that $\|I - M^{-1}(ud\mu)\| \leq C(N, r)\|u\|_1 \sum_i r_i^{l_1+l_2}$, while (3.15) gives us that $\|A(ud\mu - \tilde{u}d\mu)\| \leq C(N, r) \sum_i r_i^{l_2} \|u - \tilde{u}\|_1$. In both cases we have used that the total variation of $u_{i,j} - \tilde{u}_{i,j}$ is bounded by $\|u_{i,j} - \tilde{u}_{i,j}\|_1$ and that writing $u_X = \int_X u$ we can calculate

$$\int |u_{i,j} - \tilde{u}_{i,j}| = \int_X \left| u_X^{-1} u(x) - \tilde{u}_X \tilde{u}(x) \right| \leq \tilde{u}_X^{-1} (1 + |u|_X u_X^{-1}) \int |u - \tilde{u}| \leq 8\|u - \tilde{u}\|_1$$

The conclusion is then that $r_j^{-l_1} |b_{0,j} - \tilde{b}_{0,j}| \leq C(N, r)\|u - \tilde{u}\|_1 \sum_i r_i^{l_2}$. Substituting this into (3.21) and using (3.7) and (3.8) we find that on $X \setminus Y$

$$\begin{aligned}
|G(v - \tilde{v})(x)| &\leq C(N, r) \left(\sum_{i=1}^N r_i^{l_2} \right) \sup_{i,j} \|u_{i,j}d\mu - \tilde{u}_{i,j}d\mu\| + C(N, r) \left(\sum_i r_i^{l_2} \right) \|u - \tilde{u}\|_1 \\
&\leq C(N, r) \left(\sum_i r_i^{l_2} \right) \|u - \tilde{u}\|_1
\end{aligned}$$

because the total variation $\|u_{i,j}d\mu - \tilde{u}_{i,j}d\mu\|$ was already computed to be at most $8\|u - \tilde{u}\|_1$. On the rest of X we must use (3.6) instead of (3.7). The weaker estimate is easily computed to be

$$(3.22) \quad |G(v - \tilde{v})(x)| \leq C(N, r)\|u - \tilde{u}\|_1$$

From our estimates on $G(v - \tilde{v}) = \Psi u - \Psi \tilde{u}$ we see that Ψ is a contraction on L^1 if l_1 and l_2 are sufficiently large, because

$$\begin{aligned} \|\Psi u - \Psi \tilde{u}\|_1 &\leq \mu(X \setminus Y)C(N, r) \left(\sum_i r_i^{l_2} \right) \|u - \tilde{u}\|_1 + \mu(Y)C(N, r) \|u - \tilde{u}\|_1 \\ &\leq C(N, r) \left(\mu(Y) + \sum_j r_j^{l_2} \right) \|u - \tilde{u}\|_1 \end{aligned}$$

It follows readily that Ψ has a unique fixed point in \mathcal{C} and $\Psi^k u$ converges to this fixed point in L^1 . From (3.22) this convergence is uniform, and we notice that the correct choice of l_2 provides $|u_0 - 1| = |\Psi u_0 - 1| \leq \epsilon$ on $X \setminus Y$ by (3.18).

It remains only to see that u_0 is a smooth bump function on X . Inductively suppose $\Psi^k u \in \text{dom}(\Delta^k)$ and both $\Delta^j \Psi^k u \equiv 0$ on ∂X for $0 \leq j \leq k$ and $\partial_n \Delta^j \Psi^k u \equiv 0$ on ∂X for $0 \leq j \leq k-1$. This is certainly true for $k=0$. By construction, $\Delta \Psi^{k+1} u$ is a linear combination of rescaled copies of $\Psi^k u$ that have been extended by zero as in (3.9). Each of these functions is in $\text{dom}(\Delta^k)$ by the matching conditions for the Laplacian, so we conclude that $\Psi^{k+1} u \in \text{dom}(\Delta^{k+1})$. It is immediate that $\Delta^j \Psi^{k+1} u \equiv 0$ on ∂X for $1 \leq j \leq k+1$ and $\partial_n \Delta^j \Psi^{k+1} u \equiv 0$ on ∂X for $1 \leq j \leq k$. By Lemma 3.7 we know also that $\Psi^{k+1} u$ and $\partial_n \Psi^{k+1} u$ vanish on ∂X , which closes the induction and establishes that $u_0 \in \text{dom}(\Delta^\infty)$ and vanishes to infinite order on ∂X . \square

4. A BOREL THEOREM ON P.C.F. FRACTALS

The classical Borel theorem tells us that given any neighborhood of $x_0 \in \mathbb{R}$ and any prescribed sequence of values for u and its derivatives at x_0 , we may construct a smooth function u with support in the neighborhood and the given sequence of derivatives at x_0 . Using the smooth bump functions we have constructed, we now show that the same result holds at junction points of certain p.c.f. fractals. In what follows X is p.c.f. and self-similar under $\{F_j\}_{j=1}^N$ and the measure μ is self-similar with factors $0 < \mu_j < 1$, $\sum_1^N \mu_j = 1$, so that $\mu(F_w(X)) = \prod_{j=1}^m \mu_{w_j}$ when w is the word $w_1 \dots w_m$. The Dirichlet form is that associated to a regular self-similar harmonic structure with resistance renormalization factors $0 < r_j < 1$ for $j = 1, \dots, N$. Our arguments depend on the existence of smooth bumps as previously constructed. The crucial step is the existence of smooth functions with finitely many prescribed normal derivative values, which is established in the following lemma.

Lemma 4.1. *Given a boundary point $q \in V_0$ there are smooth functions f_l such that*

$$\begin{aligned} \Delta^k f_l(p) &= 0 \quad \text{for all } k \geq 0 \text{ and all } p \in V_0 \\ \partial_n \Delta^k f_l(p) &= 0 \quad \text{for all } k \geq 0 \text{ and all } p \in V_0 \setminus \{q\} \\ \partial_n \Delta^k f_l(q) &= \delta_{lk} \end{aligned}$$

Proof. We begin with the case $l = 0$. If U is the smooth bump function on X produced in Theorem 3.8 we localize it near the boundary points of X at a scale m to be determined later. Define

$$U_j = \begin{cases} U \circ F_j^{-m} & \text{on } F_j^m(X) \\ 0 & \text{otherwise} \end{cases}$$

and observe from the matching conditions for the Laplacian that each U_j is smooth. Now apply the Dirichlet Green's operator G to these functions and form the linear combination

$$f = \sum_{j=1}^N a_j G(U_j)$$

with coefficients to be chosen. It is clear from the properties of U that $\Delta^k f = 0$ on V_0 for all $k \geq 0$ and that $\partial_n \Delta^k f = 0$ on V_0 if $k \geq 1$. Moreover the Gauss-Green formula yields values of the normal derivatives at the points $q_i \in V_0$

$$\partial_n f(q_i) = - \sum_{j=1}^N a_j \int_X h_i U_j$$

where h_i is the harmonic function on X with boundary values $h_i(q_j) = \delta_{ij}$. In order that there be coefficients a_j such that f has the properties asserted for f_0 it then suffices that we can invert the matrix with entries $A_{ij} = \int h_i U_j$. We use the fact that

$$\begin{aligned} |h_i| &\leq r_j^m \quad \text{on } F_j^m(X) \quad \text{for } j \neq i \\ |h_i - 1| &\leq r_i^m \quad \text{on } F_i^m(X) \end{aligned}$$

which follow from the estimates on the oscillation of a harmonic function that were mentioned at the end of the introduction. Using this we calculate

$$(4.1) \quad \begin{aligned} |A_{ij}| &= \left| \int h_i U_j \right| \leq r_j^m \mu_j^m \int_X |U| \quad \text{for } j \neq i \\ \left| A_{ii} - \mu_i^m \int_X U \right| &= \left| \int (h_i - 1) U_i \right| \leq r_i^m \mu_i^m \int_X |U| \end{aligned}$$

Let D be the diagonal matrix with entries $D_{ii} = \mu_i^m \int_X U$. Then we readily compute $(AD^{-1})_{ij} = \mu_j^{-m} (\int U)^{-1} A_{ij}$ is close to the identity if m is large. Indeed, by (4.1) we have $|(I - AD^{-1})_{ij}| \leq C\rho^m$ with $\rho = \max_i r_i$ and $C = (\int U)^{-1} (\int |U|)$, so that AD^{-1} is invertible provided m is sufficiently large.

We proceed by induction on l , with an almost unchanged argument. Suppose the functions f_l for $l \leq L - 1$ have been constructed as linear combinations of the form

$$(4.2) \quad f_l = \sum_{n=1}^l \sum_{j=1}^N a_{jn} G^{(n+1)}(U_j) \quad l \leq L - 1$$

and consider the function

$$f = \sum_{j=1}^N a_j G^{(L+1)}(U_j)$$

so that $\Delta^k f = 0$ on V_0 for all k and $\partial_n \Delta^k f = 0$ on V_0 for $k \geq L + 1$. When $k = L$ we have

$$\partial_n \Delta^L f(q_i) = - \sum_{j=1}^N a_j A_{ij}$$

where A_{ij} is as before, so we may select a_j to obtain $\partial_n \Delta^L f(q) = 1$ and $\partial_n \Delta^L f(p) = 0$ at other points $p \in V_0$. Subtracting an appropriate linear combination of the f_l for $l \leq L - 1$ we obtain the desired f_L in the form of (4.2). \square

With this in hand it is simple to deal with finitely many values of the Laplacian at a boundary point $q \in V_0$.

Lemma 4.2. *Given a boundary point $q \in V_0$ there are smooth functions g_l such that*

$$\begin{aligned} \partial_n \Delta^k g_l(p) &= 0 \quad \text{for all } k \geq 0 \text{ and all } p \in V_0 \\ \Delta^k g_l(p) &= 0 \quad \text{for all } k \geq 0 \text{ and all } p \in V_0 \setminus \{q\} \\ \Delta^k g_l(q) &= \delta_{lk} \end{aligned}$$

Proof. Let h be the harmonic function which is 1 at q and 0 at all other points of V_0 . Clearly $\Delta^k h \equiv 0$ for all $k \geq 1$ and therefore also $\partial_n \Delta^k h(p) = 0$ for all $k \geq 1$ and $p \in V_0$. For each $p \in V_0$ let $f_{0,p}$ be the function constructed in Lemma 4.1 with non-vanishing normal derivative at p . It is clear that $g_0 = h - \sum_{p \in V_0} (\partial_n h(p)) f_{0,p}$ has the desired properties, so we have found the first of our functions. To obtain the others we simply apply the Dirichlet Green's operator G . Notice that $\Delta^k G^l h(p) = \delta_{k,l} \delta_{p,q}$ for all k and all $p \in V_0$, and also $\partial_n \Delta^k G^l h(p) = 0$ for all $k \geq l + 1$. To obtain g_k it remains only to subtract off all normal derivatives that occur for $0 \leq k \leq l$ using the functions from Lemma 4.1. \square

The proof of a Borel-type theorem from the above lemmas is standard. All that is needed is information about how scaling the support of a function changes its Laplacian and normal derivatives. Recall that for a Dirichlet form associated to a regular self-similar resistance, both the Laplacian and the normal derivative may be obtained as renormalized limits of corresponding quantities defined on the approximating graphs (Section 3.7 of [14]). In particular, pre-composition with the map F_i^{-1} rescales the k -th power of the Laplacian by $(\mu_i r_i)^{-k}$ and its normal derivative by $\mu_i^{-k} r_i^{-k-1}$. For this reason, if $q = q_i$ is the boundary point of interest we define

$$\begin{aligned} f_{l,m} &= \begin{cases} \mu_i^{ml} r_i^{m(l+1)} f_l \circ F_i^{(-m)} & \text{on } F_i^m \\ 0 & \text{otherwise} \end{cases} \\ g_{l,m} &= \begin{cases} (\mu_i r_i)^{ml} g_l \circ F_i^{(-m)} & \text{on } F_i^m \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

so that we have for all k

$$(4.3) \quad \begin{aligned} \Delta^k f_{l,m}(q) &= 0 & \Delta^k g_{l,m}(q) &= \delta_{lk} \\ \partial_n \Delta^k f_{l,m}(q) &= \delta_{lk} & \partial_n \Delta^k g_{l,m}(q) &= 0 \end{aligned}$$

but the L^∞ norms of the lower order derivatives have decreased and those of the higher order derivatives have increased.

$$(4.4) \quad \|\Delta^k f_{l,m}\|_\infty = \mu_i^{m(l-k)} r_i^{m(l+1-k)} \|\Delta^k f_l\|_\infty$$

$$(4.5) \quad \|\Delta^k g_{l,m}\|_\infty = (\mu_i r_i)^{m(l-k)} \|\Delta^k g_l\|_\infty$$

With this in hand we can easily prove our version of the Borel theorem.

Theorem 4.3. *Let $q \in V_0$ be fixed, and Ω be an open neighborhood of q . Given a jet $\rho = (\rho_0, \rho_1, \dots)$ of values for powers of the Laplacian and $\sigma = (\sigma_0, \sigma_1, \dots)$ of values for their normal derivatives, there is a smooth function f with support in Ω and both $\Delta^k f(q) = \rho_k$ and $\partial_n \Delta^k f(q) = \sigma_k$ for all k .*

Proof. We give the usual proof that it is possible to define f by the series

$$(4.6) \quad f = \sum_l \left(\rho_l g_{l, m_l} + \sigma_l f_{l, n_l} \right)$$

for an appropriate choice of m_l and n_l .

Let $m_0 = n_0$ be sufficiently large that $F_i^{m_0}(X) \subset \Omega$. For each l choose $m_l \geq m_0$ so large that

$$\|\rho_l \Delta^k g_{l, m_l}\|_\infty \leq 2^{k-l-1} \quad \text{for } 0 \leq k \leq l-1$$

using the scaling estimate (4.5). Similarly use the scaling relation (4.4) to take $n_l \geq m_0$ such that

$$\|\sigma_l \Delta^k f_{l, n_l}\|_\infty \leq 2^{k-l-1} \quad \text{for } 0 \leq k \leq l-1$$

Then for fixed k we have

$$\left\| \sum_{l=k+1}^{\infty} \Delta^k \left(\rho_l g_{l, m_l} + \sigma_l f_{l, n_l} \right) \right\|_\infty \leq \sum_{k+1}^{\infty} 2^{k-l} \leq 1$$

so that all powers of the Laplacian applied to (4.6) produce L^∞ convergent series. It follows that f as defined in (4.6) is smooth and has support in Ω . By (4.3) it has the desired jet, so the result follows. \square

We remark that for any $\epsilon > 0$ we could replace the bounds 2^{k-l-1} in the proof with $\epsilon 2^{k-l-1}$. It follows that we can define f by (4.6) and have the estimate

$$(4.7) \quad \|\Delta^k f\|_\infty \leq C(k, \Omega) \sum_{l=0}^k (|\rho_l| + |\sigma_l|) + \epsilon$$

where $C(k, \Omega)$ does not depend on the jet we prescribe.

It is also possible to estimate the effect of the size of the support $F_i^{m_0}(X)$ on a finite collection of jet terms. Take $m_l = m_0$ for $0 \leq j \leq L$, $n_l = m_0$ for $0 \leq j \leq L-1$ and thereafter choose both so large that the estimates $\epsilon 2^{k-l-1}$ hold. From (4.4) and (4.5) we deduce for $0 \leq k \leq L$,

$$(4.8) \quad \|\Delta^k f\|_\infty \leq C(k) (r_i \mu_i)^{-m_0 k} \left(\sum_{l=0}^L (r_i \mu_i)^{m_0 l} |\rho_l| + \sum_{l=0}^{L-1} r_i^{m_0(l+1)} \mu_i^{m_0 l} |\sigma_l| \right) + \epsilon.$$

Similar estimates hold for other norms and seminorms, provided only that their scaling is understood. For example, if we consider the seminorm $\mathcal{E}(\Delta^k f)^{1/2}$, then the scaling in (4.4) and (4.5) is reduced by $r_i^{-m/2}$, so the above construction would give

$$\mathcal{E}(\Delta^k f)^{1/2} \leq C(k) r_i^{-m_0(k+1/2)} \mu_i^{-m_0 k} \left(\sum_{l=0}^L (r_i \mu_i)^{m_0 l} |\rho_l| + \sum_{l=0}^{L-1} r_i^{m_0(l+1)} \mu_i^{m_0 l} |\sigma_l| \right) + \epsilon.$$

The result of the theorem may be transferred to any junction point $F_w(q)$ and cell $F_w(X)$ in X , simply by modifying the desired jet to account for the effect of composition with F_w , solving for f on X , and defining the new function to be $f \circ F_w^{-1}$ on the cell. We record a version of this that will be useful later; note that in the following we use the notation ∂_n for the normal derivative with respect to the cell $F_w(X)$.

Corollary 4.4. *Let $F_w(q)$ be a junction point in X . Given a jet (ρ_0, ρ_1, \dots) , $(\sigma_1, \sigma_2, \dots)$ there is a smooth function f on $F_w(X)$ that has $\Delta^k f(p) = \partial_n \Delta^k f(p) = 0$ at all points $p \in \partial F_w(X)$ such that $p \neq F_w(q)$, and satisfies $\Delta^k f(F_w(q)) = \rho_k$ and $\partial_n \Delta^k f(F_w(q)) = \sigma_k$ for all k .*

5. ADDITIVE PARTITIONS OF FUNCTIONS

The results of [4] show that multiplication is not generally a good operation on functions in $\text{dom}(\Delta)$. In particular, for X a p.c.f. fractal with self-similar measure and regular self-similar harmonic structure it is generically the case that if $u \in \text{dom}(\Delta)$ then $u^2 \notin \text{dom}(\Delta)$. In such a situation there is no hope of using a smooth partition of unity to localize problems in the classical manner. Instead we provide a simple method for making a smooth decomposition of $f \in \text{dom}(\Delta^\infty)$ using Theorem 4.3. Throughout this section we make the same assumptions on X as were made in Section 4.

Theorem 5.1. *Let $\cup_\alpha \Omega_\alpha$ be an open cover of X and $f \in \text{dom}(\Delta^\infty)$. There is a decomposition $f = \sum_{k=1}^K f_k$ in which each k has a corresponding α_k such that f_k is smooth on X and supported in Ω_{α_k} .*

Proof. Compactness of X allows us to reduce to the case of a finite cover $\cup_1^K \Omega_{\alpha_k}$ for which there is no sub-collection that covers X . We write Ω_k for Ω_{α_k} , and construct the functions f_k inductively. At the k -th stage we suppose there are functions f_1, \dots, f_{k-1} with the properties asserted in the lemma, and that function $g_{k-1} = f - \sum_{l=1}^{k-1} f_l$ is smooth on X and vanishes identically on a neighborhood Π_{k-1} of $X \setminus \left(\cup_{j=k}^K \Omega_j\right)$. In the base case $k = 1$ this assumption is trivial, and it is clear that the theorem follows immediately from the induction. We have therefore reduced to the case where our cover consists of the two sets Ω_k and $\tilde{\Omega}_k = \cup_{j=k+1}^K \Omega_j$, because the induction is complete once we have f_k as in the lemma such that g_k is identically zero on a neighborhood Π_k of $X \setminus \tilde{\Omega}_k$.

For a scale m and $x \in X$, define the m -scale open neighborhood of x to be the interior of the unique m -cell containing x if $x \notin V_m$, and to be the union $\{x\} \cup (\cup_w F_w(X) \setminus \partial F_w(X))$ if $x = F_w(q_i)$ is a junction point. By Section 1.3 of [14], the m -scale open neighborhoods form a fundamental system of open neighborhoods of x . At each x in Ω_k there is a largest m such that the m -scale neighborhood of x is contained in Ω_k . The collection of all such largest neighborhoods of points of Ω_k is an open cover of the compact set $\text{Sppt}(g_{k-1}) \setminus \tilde{\Omega}_k$. We use Λ_k to denote the union over a finite subcover.

Clearly Λ_k has finitely many boundary points. Let those boundary points that are also in $\tilde{\Omega}_k$ be x_1, \dots, x_J , and take at each a finite collection of cells $\{C_{i,j}\}_{i=1}^{I_j}$ having x_j in their boundary. We require that $\Lambda_k \cup \left(\cup_{i=1}^{I_j} C_{i,j}\right)$ contains a neighborhood of x_j , that all of the $C_{i,j}$ lie entirely within Ω_k and none intersect Λ_k , and that $C_{i,j} \cap C_{i',j'}$ is empty unless $j = j'$, in which case it contains only x_j . On each cell we apply Corollary 4.4 to find functions $h_{i,j}$ that match $g_{k-1}(x_j)$ and all powers of its Laplacian at x_j , and such that the sum $\sum_i h_{i,j}$ has normal derivatives that cancel $\partial_n \Delta^n g_{k-1}(x_j)$ at x_j for all n . Thus $\sum_{i=1}^{I_j} h_{i,j}$ matches g_{k-1} in the sense of the matching condition and vanishes to infinite order at the other boundary points of $\cup_{i=1}^{I_j} C_{i,j}$. The matching condition implies that $f_k = g_{k-1}|_{\Lambda_k} + \sum_j \sum_i h_{i,j}$ is smooth. It is clearly supported on Ω_k and equal to f on the closure $\bar{\Lambda}_k$ of Λ_k ,

so $g_k = g_{k-1} - f_k$ is zero on a neighborhood Π_k of $X \setminus \tilde{\Omega}_k$, which completes the induction and the proof. \square

Remark 5.2. Theorem 5.1 is stronger than the partitioning results we obtained directly from the heat kernel smoothing procedure of Section 2, but only applies to the more limited case of p.c.f. self-similar sets with regular Dirichlet form. We note, however, that it provides a stronger version of Theorem 3.8, because by smoothly cutting off the constant function 1 we obtain a smooth bump that is identically 1 on a compact K and 0 outside an ϵ -neighborhood of K . In contrast to Corollary 2.9 it does not imply that there is a positive smooth bump of this type.

REFERENCES

- [1] Martin T. Barlow, *Diffusions on fractals*, Lectures on probability theory and statistics (Saint-Flour, 1995), Lecture Notes in Math., vol. 1690, Springer, Berlin, 1998, pp. 1–121. MR 1668115 (2000a:60148)
- [2] Martin T. Barlow and Richard F. Bass, *The construction of Brownian motion on the Sierpiński carpet*, Ann. Inst. H. Poincaré Probab. Statist. **25** (1989), no. 3, 225–257. MR 1023950 (91d:60183)
- [3] ———, *Brownian motion and harmonic analysis on Sierpinski carpets*, Canad. J. Math. **51** (1999), no. 4, 673–744. MR 1701339 (2000i:60083)
- [4] Oren Ben-Bassat, Robert S. Strichartz, and Alexander Teplyaev, *What is not in the domain of the Laplacian on Sierpinski gasket type fractals*, J. Funct. Anal. **166** (1999), no. 2, 197–217. MR 1707752 (2001e:31016)
- [5] Nitsan Ben-Gal, Abby Shaw-Krauss, Robert S. Strichartz, and Clint Young, *Calculus on the Sierpinski gasket. II. Point singularities, eigenfunctions, and normal derivatives of the heat kernel*, Trans. Amer. Math. Soc. **358** (2006), no. 9, 3883–3936 (electronic). MR 2219003 (2007h:28009)
- [6] E. B. Davies, *Heat kernels and spectral theory*, Cambridge Tracts in Mathematics, vol. 92, Cambridge University Press, Cambridge, 1989. MR 990239 (90e:35123)
- [7] Pat J. Fitzsimmons, Ben M. Hambly, and Takashi Kumagai, *Transition density estimates for Brownian motion on affine nested fractals*, Comm. Math. Phys. **165** (1994), no. 3, 595–620. MR 1301625 (95j:60122)
- [8] Masatoshi Fukushima, Yōichi Ōshima, and Masayoshi Takeda, *Dirichlet forms and symmetric Markov processes*, de Gruyter Studies in Mathematics, vol. 19, Walter de Gruyter & Co., Berlin, 1994. MR 1303354 (96f:60126)
- [9] A. A. Grigor'yan, *The heat equation on noncompact Riemannian manifolds*, Mat. Sb. **182** (1991), no. 1, 55–87. MR 1098839 (92h:58189)
- [10] Alexander Grigor'yan and Andras Telcs, *Sub-Gaussian estimates of heat kernels on infinite graphs*, Duke Math. J. **109** (2001), no. 3, 451–510. MR 1853353 (2003a:35085)
- [11] Alexander Grigor'yan and András Telcs, *Harnack inequalities and sub-Gaussian estimates for random walks*, Math. Ann. **324** (2002), no. 3, 521–556. MR 1938457 (2003i:58068)
- [12] B. M. Hambly and T. Kumagai, *Transition density estimates for diffusion processes on post critically finite self-similar fractals*, Proc. London Math. Soc. (3) **78** (1999), no. 2, 431–458. MR 1665249 (99m:60118)
- [13] P. Edward Herman, Roberto Peirone, and Robert S. Strichartz, *p-energy and p-harmonic functions on Sierpinski gasket type fractals*, Potential Anal. **20** (2004), no. 2, 125–148. MR 2032945 (2004k:31023)
- [14] Jun Kigami, *Analysis on fractals*, Cambridge Tracts in Mathematics, vol. 143, Cambridge University Press, Cambridge, 2001. MR 1840042 (2002c:28015)
- [15] ———, *Harmonic analysis for resistance forms*, J. Funct. Anal. **204** (2003), no. 2, 399–444. MR 2017320 (2004m:31010)
- [16] Takashi Kumagai, *Short time asymptotic behaviour and large deviation for Brownian motion on some affine nested fractals*, Publ. Res. Inst. Math. Sci. **33** (1997), no. 2, 223–240. MR 1442498 (98k:60130)
- [17] Tom Lindstrøm, *Brownian motion on nested fractals*, Mem. Amer. Math. Soc. **83** (1990), no. 420, iv+128. MR 988082 (90k:60157)

- [18] Robert Meyers, Robert S. Strichartz, and Alexander Teplyaev, *Dirichlet forms on the Sierpiński gasket*, Pacific J. Math. **217** (2004), no. 1, 149–174. MR 2105771 (2005k:31028)
- [19] Jonathan Needleman, Robert S. Strichartz, Alexander Teplyaev, and Po-Lam Yung, *Calculus on the Sierpinski gasket. I. Polynomials, exponentials and power series*, J. Funct. Anal. **215** (2004), no. 2, 290–340. MR 2150975
- [20] Kasso A. Okoudjou, Luke G. Rogers, and Robert S. Strichartz, *Generalized eigenfunctions and a Borel theorem on the Sierpinski gasket*, To Appear in Canad. Math. Bull.
- [21] Roberto Peirone, *Convergence and uniqueness problems for Dirichlet forms on fractals*, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) **3** (2000), no. 2, 431–460. MR 1769995 (2001i:31016)
- [22] Luke G. Rogers and Robert S. Strichartz, *Distributions on p.c.f. fractafolds*, In preparation.
- [23] C. Sabot, *Existence and uniqueness of diffusions on finitely ramified self-similar fractals*, Ann. Sci. École Norm. Sup. (4) **30** (1997), no. 5, 605–673. MR 1474807 (98h:60118)
- [24] L. Saloff-Coste, *A note on Poincaré, Sobolev, and Harnack inequalities*, Internat. Math. Res. Notices (1992), no. 2, 27–38. MR 1150597 (93d:58158)
- [25] Robert S. Strichartz, *Analysis on products of fractals*, Trans. Amer. Math. Soc. **357** (2005), no. 2, 571–615 (electronic). MR 2095624 (2005m:31016)
- [26] ———, *Differential equations on fractals*, Princeton University Press, Princeton, NJ, 2006, A tutorial. MR 2246975 (2007f:35003)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269-3009, U.S.A.

E-mail address: `rogers@math.uconn.edu`

DEPARTMENT OF MATHEMATICS, MALOTT HALL, CORNELL UNIVERSITY, ITHACA, NY 14853-4201, U.S.A.

E-mail address: `str@math.cornell.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269-3009, U.S.A.

E-mail address: `teplyaev@math.uconn.edu`